

SERIES PUBLICATION,
THE FAUNA OF BRITISH INDIA,

INCLUDING

CEYLON AND BURMA.

PUBLISHED UNDER THE AUTHORITY OF THE SECRETARY OF
STATE FOR INDIA IN COUNCIL

EDITED BY LIEUT.-COL. R. B. S. SEWELL, C.I.E., M.C.D., F.R.S.

ASSISTED BY F. W. EDWARDS, M.A., Sc.D.

DIPTERA
VOL. V.


Family CULICIDÆ.

Tribes MEGARHININI and CULICINI.

BY

CAPT. P. J. BARRAUD, F.E.S., F.Z.S., F.L.S.

*Entomologist to the Malaria Survey of India.
Formerly Entomologist to the Kala-azar Commission in India,
and Officer-in-Charge of Inquiry on Indian Culicidæ*



LONDON:

TAYLOR AND FRANCIS, RED LION COURT, FLEET STREET.

March, 1934.

2687
591.954
N20-5



(19)



PRINTED BY TAYLOR AND FRANCIS,
RED LION COURT, FLEET STREET.

2687

571

CONTENTS.

	Page
ASSISTANT-EDITOR'S PREFACE	v
AUTHOR'S PREFACE	ix
BIBLIOGRAPHY	xi ,
SYSTEMATIC INDEX	xxii
ABBREVIATIONS USED IN TEXT	xxvii
ERRATA	xxviii
INTRODUCTION	1
SYSTEMATIC :	
Tribe Megarhinini	8
Tribe Culicini	28
APPENDIX (by F. W. EDWARDS)	427
ALPHABETICAL INDEX	455

6

7

8

9

ASSISTANT-EDITOR'S PREFACE.

THE proposal for the preparation of a volume on Mosquitoes in the ' Fauna of British India ' series was first made in 1920, when the late Editor (Sir Arthur Shipley), with the sanction of the Secretary of State for India, arranged with Col. (now Sir) S. R. Christophers and myself to undertake the work. However, at about the same time Capt. P. J. Barraud had received a commission under the Indian Research Fund Association to make a general survey of the mosquitoes of the Indian Empire. In view of this appointment, it appeared undesirable that the proposed volume should be proceeded with until the results of Capt. Barraud's survey were available for inclusion, and publication was therefore deferred. Meanwhile some work which I had already done was published in the form of a synopsis of the Culicine mosquitoes of the Oriental region, and the results of Capt. Barraud's survey were published by him in a series of papers revising the Culicine mosquitoes of India, all of which work appeared in the ' Indian Journal of Medical Research.'

During the course of Capt. Barraud's survey a surprising wealth of new forms was discovered, and the decision to delay publication thus proved amply justified. Moreover, it soon became obvious that it would hardly be possible to include all the new information available in a volume of the ordinary size, at least without undue condensation; it was, therefore,

decided to divide the work, issuing one volume on Anopheline and one on Culicine mosquitoes. The former, by Col Sir Rickard Christophers, was published on November 27th, 1933

Partly because of pressure of other work, and partly because the far greater share undertaken by Capt. Barraud entitled him to the fullest recognition as author, I preferred to withdraw from joint authorship of the volume on Culicini. Nevertheless my position as assistant-editor has enabled me to take some share in its production, and at the request of the author I have carefully checked, and in a few cases revised, his keys, and, where necessary, supplemented his account from a study of additional material available in the collections of the British Museum. In view of the fact that the volume will no doubt have a wide circulation outside India, it seemed desirable to include some reference to Oriental species which have not yet been found within the Indian area, indicating briefly their distinctions from known Indian forms. This information is usually given in the form of footnotes.

All interpolations made by me in the Author's manuscript are indicated by enclosure within square brackets

Owing to the necessity of compressing a descriptive account of the adults and early stages of nearly 250 species of Indian Culicine mosquitoes within the limits of a volume of 500 pages, it was decided not to attempt to include an extended introduction, however much this might be desirable. Reference may, however, be made to Sir Rickard Christophers' companion volume on Anophelini, the Introduction to which is largely applicable also to Culicini.

The present volume concludes the account of the Indian Culicinae, the subfamily which, according to our present classification, includes all the true mosquitoes. For the sake of completeness I have given in an Appendix a brief account of the distinguishing characters of the family Culicoidae and its relationships with other families of Nematocerous Diptera, and also descriptions of the very few Indian members of the other two subfamilies, the Dixinae and Chaoborinae.

Nearly all the line drawings, and a large number of the photographic illustrations published by Capt Barraud in his 'Revision of the Culicine Mosquitoes of India,' are reproduced here on a smaller scale, permission to copy having been kindly granted by the Secretary of the Scientific Advisory Board of the Indian Research Fund Association

In addition to previously published illustrations, a large number of new figures have been specially prepared for this volume. Some of these are the work of the Author, but the majority are by Mr A. J. E. Terzi, whose fine work is its own recommendation

Mr S. Maulik has kindly assisted in reading the proofs, and has checked the spelling of the place-names so as to ensure uniformity with that adopted in the Indian Post Office Guide.

In spite of the vast increase in our knowledge of Indian mosquitoes within the past twenty years—an increase through which the Culicidæ have become one of the best-known groups of insects in the Indian fauna—a great deal of work remains to be done before our knowledge even approaches completeness. As will be noted from Capt Barraud's account, the larvæ of 96 of the 239 described Indian Culicini are still unknown, and regarding the distribution of these and others within the area our information is very meagre. Little or no collecting has yet been done in some areas, notably Tenasserim and parts of central India, and even in the areas where most work has been accomplished there can be little doubt that new species await discovery.

It has sometimes been remarked that mosquitoes can be divided into two main groups: those which are found only as adults, and those found only as larvæ. The truth behind this remark is that the breeding-places of some of the common domestic species are easily overlooked, whereas many species of which the larvæ are easily found are not aggressive blood-suckers. The adult mosquitoes which

obtrude themselves on our notice are by now fairly well known, but this cannot yet be said of the many "wild" species which seldom enter dwellings, whose breeding-places are, nevertheless, not difficult to discover. The most likely habitats in which to search for larvæ of hitherto undescribed species are the small collections of water to be found in various plants. Much attention has been paid to the tree-hole and bamboo fauna, particularly in the Himalayan region, and many remarkable discoveries have been made. Some other types of breeding-place, however, have been but little investigated, partly, perhaps, because few or no Anophelini are to be found in them. For example, in the Malayan region many Culicine mosquitoes are found breeding in the water contained in the leaves of pitcher-plants (*Nepenthes*), in leaf-axils of *Colocasia*, *Crinum*, *Pandanus*, and *Susum*, and in flowers or inflorescences of *Cyrtandra*, *Hymenocallis*, and *Rafflesia*. Many species of mosquitoes are more or less confined to such situations, the adults being rarely seen unless the larvæ are collected and reared. No doubt the number of water-bearing plants is higher in the Malayan than the Indian region, but the genera *Colocasia*, *Crinum*, and *Nepenthes* are widely distributed in India and Ceylon, and should repay investigation.

F. W. EDWARDS

London,
February 1934

AUTHOR'S PREFACE.

THIS volume embodies the results of several years' work on the Megarhinine and Culicine mosquitoes of India which the writer has been able to carry out whilst employed under the Indian Research Fund Association. The Author wishes to express his sincere thanks to Colonel Sir Rickard Christophers, Kt, C.I.E., O.B.E., F.R.S. (I.M.S., retired), for initiating an inquiry into this subject, and for much help and encouragement over a period of many years. The Author is also greatly indebted to Dr. F. W. Edwards, of the British Museum (Natural History), for kind assistance in many ways, especially in editing the manuscript of this work, arranging for a number of the illustrations, and in lending his MSS. of the 'Genera Insectorum, Family Culicidæ,' before publication. The Author's thanks are also due to Lieut.-Colonel J. A. Sinton, V.C., O.B.E., I.M.S., Director of the Malaria Survey of India, for much kind assistance and advice, and to many others, both in India and other countries, for the gift or loan of valuable material.

Full use has been made of publications by other writers, the titles of which will be found in the list of references.

Endeavour has been made to bring the subject up to date, but there is little doubt that many additional species of the Culicini still await discovery, especially in the south-western part of the Peninsula and in Burma. It is interesting to look back to 1900, when Colonel G. M. Giles, I.M.S., in reading a paper before the Bombay Natural History Society, stated that "Two years ago, when I took up the task of collecting

the literature of the Culicidæ, it is an actual fact that no more than four species were recorded as having been found in all India. There was, in fact, hardly any other known country with such scanty records of the subject. The subjoined list includes 32 species, and I have little doubt the final total of species will be found to be not far off a hundred, as new species are constantly turning up "

The number (including 43 Anophelini) has now reached a total of 288, of the three tribes included in the subfamily Culicinae, not counting named varieties, some of which may eventually be shown to be distinct species. The larvæ of about 40 species of the Culicini are here described for the first time, but those of many others remain unknown.

With very few exceptions the Author has been able to examine adult specimens of all species. In the records of distribution the localities from which the Author has examined specimens are marked with an asterisk.

P J BARRAUD

Kasauli,
July 1933

BIBLIOGRAPHY.

- ACTON, H W., and RAO, S. S. 1931.—The Diagnosis of Lymphatic Obstruction of Filarial Origin. *Ind Med Gaz* lxi, pp. 11-17.
- ANNANDALE, N 1911.—A new Genus of Short-beaked Gnats from Ceylon. *Spolia Zeylanica*, vii, pp 187-193.
- BACOT, A. W 1916.—Yellow Fever Commission (W Africa) Ent. Rept., pp 130 & 142.
- BARRAUD, P. J 1920.—Notes on some Culicids collected in Lower Mesopotamia. *Bull Ent. Res* x, pp 323-325.
- 1921.—Mosquitoes collected in Palestine and Adjacent Territories. *Ibid* xi, pp. 387-395.
- 1923 a.—A Revision of the Culicine Mosquitoes of India.—I. *Stegomyia* and *Christophersomyia*. *Ind. Journ Med. Res* x, pp 772-788.
- 1923 b.—Some new Culicine Mosquitoes found in India, and a Note on *Finlaya assamensis* (Theo.) *Bull Ent Res* xii, pp. 405-408.
- 1923 c.—A Revision of the Culicine Mosquitoes of India.—II. The Larvæ of some Indian Species of *Culex*. *Ind Journ. Med. Res* x, pp 934-942.
- 1923 d.—A Revision of the Culicine Mosquitoes of India.—III. Notes on certain Species of the Genus *Finlaya* and Descriptions of new Species. *Ibid* xi, pp 214-219.
- 1923 e.—A Revision of the Culicine Mosquitoes of India.—IV. The Larvæ of some Indian Species of *Finlaya*. *Ibid* xi, pp 220-223.
- 1923 f.—A Revision of the Culicine Mosquitoes of India.—V. Further Notes on the Genera *Stegomyia* and *Finlaya*, with Descriptions of new Species. *Ibid.* xi, pp 224-228.
- 1923 g.—A Revision of the Culicine Mosquitoes of India.—VI. Some Indian Species of the Genus *Finlaya*. *Ibid* xi, pp 475-493.
- 1923 h.—A Revision of the Culicine Mosquitoes of India.—VII. The Larvæ of some Species of *Stegomyia*, *Christophersomyia*, *Mimomyia*, and *Alldomyia*. *Ibid.* xi, pp. 495-505.
- 1923 i.—Two new Species of *Culex* from Assam. *Ibid.* xi, pp. 507-509.
- 1924 a.—A Revision of the Culicine Mosquitoes of India.—VIII Further Descriptions of Indian Species of *Finlaya*. *Ibid.* xi, pp. 845-865.
- 1924 b.—A new Mosquito from Kashmir. *Ibid.* xi, pp. 967-968.
- 1924 c.—A Revision of the Culicine Mosquitoes of India.—IX. The Indian Species of *Lutzia*. *Ibid.* xi, pp 971-976.

- BARRAUD, P. J. 1924 *d*—A Revision of the Culicine Mosquitoes of India.—X The Larvæ of Indian Species of *Lutzia* Ind Journ Med. Res xi, pp. 977-978
- 1924 *e*—A Revision of the Culicine Mosquitoes of India.—XI Some Indian Species of *Culex* Ibid. xi, pp. 979-998
- 1924 *f*—Four new Mosquitoes from the Western Himalayas Ibid. xi, pp. 999-1006
- 1924 *g*—A Revision of the Culicine Mosquitoes of India.—XII Further Descriptions of Indian Species of *Culex*, including two new Species Ibid. xi, pp. 1259-1274
- 1924 *h*—A Revision of the Culicine Mosquitoes of India.—XIII Further Descriptions of Indian Species of *Culex*, including three new Species. Ibid. xi, pp. 1275-1282
- 1924 *i*—A Revision of the Culicine Mosquitoes of India.—XIV The Indian Species of the Subgenus *Culisomomyia* (Theo.) Edw., including one new Species Ibid. xi, pp. 15-22
- 1924 *j*. A Revision of the Culicine Mosquitoes of India.—XV The Indian Species of the Subgenus *Lophoceratomyia* (Theo.) Edw., including two new Species Ibid. xi, pp. 39-46
- 1924 *k*—A new Mosquito from Kashmir and the North-West Frontier Province Ibid. xi, pp. 73-74.
- 1924 *l*—A Revision of the Culicine Mosquitoes of India.—XVI The Indian Species of *Theobaldia*. Ibid. xi, pp. 139-143
- 1924 *m*—A Revision of the Culicine Mosquitoes of India.—XVII Further descriptions of the Larvæ of Indian Species of *Culex* Ibid. xi, pp. 427-434
- 1926.—A Revision of the Culicine Mosquitoes of India.—XVIII The Indian Species of *Uranotania* and *Harpagomyia*, with Descriptions of five new Species. Ibid. xiv, pp. 331-350
- 1927 *a*—A Revision of the Culicine Mosquitoes of India.—XIX. The Indian Species of *Aïdomyia* and *Orthopodomyia*, with Descriptions of two new Species. Ibid. xiv, pp. 523-532
- 1927 *b*.—A Revision of the Culicine Mosquitoes of India.—XX The Indian Species of *Armigeres* (including *Leicesteria*), with Descriptions of two new Species Ibid. xiv, pp. 533-548.
- 1927 *c*—A Revision of the Culicine Mosquitoes of India.—XXI Descriptions of new Species of *Aedimorphus* and *Finlaya*, and Notes on *Stegomyia albolineata*. Ibid. xiv, pp. 549-554
- 1927 *d*—A Revision of the Culicine Mosquitoes of India.—XXII. The Indian Species of *Tæniotrypanus* (including *Mansonioides*), with a Description of one new Species. Ibid. xiv, pp. 555-563
- 1928 *a*—A Revision of the Culicine Mosquitoes of India.—XXIII. The Genus *Aedes* (sens lat.) and Classification of Subgenera. Descriptions of Indian Species of *Aedes* (*Aëdimorphus*), *Aedes* (*Ochlerotatus*), and *Aedes* (*Banksimella*), with Notes on *Aedes* (*Stegomyia*) *variegatus* Ibid. xv, pp. 653-669.
- 1928 *b*.—A Revision of the Culicine Mosquitoes of India.—XXIV. The Indian Species of the Subgenera *Skusea* and *Aedes*, with Descriptions of eight new Species and Remarks on a new Method for identifying the Females of the Subgenus *Aedes* Ibid. xvi, pp. 357-375
- 1928 *c*—The Distribution of "*Stegomyia fasciata*" in India, with Remarks on Dengue and Yellow Fever Ibid. xvi, pp. 377-388.

- BARRAUD, P. J. 1929 *a*—A Revision of the Culicine Mosquitoes of India—XXV The Genera *Mucidus*, *Mimomyia*, *Picalbia*, *Rachnotomyia*, and *Hodgesia*. Ind Journ Med Res xvi, pp 1052-1063
- 1929 *b*—A Revision of the Culicine Mosquitoes of India—XXVI The Genera *Herzmannia*, *Hæmagogus*, *Topomyia*, and *Megarhinus*. *Ibid* xvii, pp 259-280
- 1931 *a*—The Early Stages of some Indian Mosquitoes *Meqarhinus*. *Ibid* xviii, pp 1127-1132
- 1931 *b*—Notes on some Indian Mosquitoes of the Subgenus *Siegomyia*, with Descriptions of new Species. *Ibid* xix, pp. 221-227
- 1931 *c*—Descriptions of Eight new Species of Indian Culicine Mosquitoes. *Ibid* xix, pp 609-616
- 1932.—The Early Stages of some Indian Mosquitoes *Orthopodomyia*. *Ibid*. xix, pp. 1013-1017
- BARRAUD, P. J., and COVELL, G. 1927.—The Morphology of the Buccal Cavity of the Mosquito. Trans Far East Assoc Trop Med, 7th Congress, iii, pp 98-102
- 1928.—The Morphology of the Buccal Cavity in Anopheline and Culicine Mosquitoes. Ind Journ Med Res xv, pp 671-679.
- BEDFORD, G. A. H. 1910.—New Culicine Larva from the Transvaal. U.S. Afr. Dept. Agric, 5th & 6th Repts. Vet. Res. pp. 739-743
- 1928.—South African Mosquitoes. U.S. Afr. Dept. Agric., 13th & 14th Repts. Vet. Res. pp. 883-990
- BONNE-WEPSTER, J. 1930.—The Genus *Tæniorhynchus* in the Dutch East Indies. Meded. Volks. Ned.-Ind. xix. Part I. *Mansonioides*, pp. 106-212. Part II. *Coquillettidia*, pp. 387-399. [Also in Dutch in Geneesk. Tijds. Ned.-Ind. (9) lxx.]
- 1932 *a*.—A Mosquito with Larval Habits like *Tæniorhynchus*. Bull. Ent. Res. xxiii, pp 60-72
- 1932 *b*.—Over symbiosen tusschen muskieton-larven in Waterplanten. Tijds. v. Ent. lxxv, pp. 254-258.
- BONNE-WEPSTER, J., and BRUG, S. L. 1932.—The Subgenus *Siegomyia* in Netherland India. Geneesk. Tijds. v. Ned.-Ind. Bijblad ii, pp. 39-119
- BORDEL, E. 1926.—Les moustiques de la Cochinchine et du Sud-Annam. Arch. Inst. Pasteur d'Indochine, nos. 3 & 4, pp. 75-121
- 1927.—Nomenclature des moustiques de la Cochinchine et du Sud-Annam. Trans. Far East Assoc Trop. Med., 7th Congress, iii, p. 165
- 1928 *a*.—Les moustiques de la Cochinchine et du Sud-Annam *Aedes*. Arch. Inst. Pasteur d'Indochine, no. 7, pp 75-106.
- 1928 *b*.—Les moustiques de la Cochinchine et du Sud-Annam.—III. *Aedes (Fulaya)*. *Ibid*. no. 8, pp. 41-76.
- BORDEL, E., and LABERNADIE, V. G. F. 1929.—A List of Species of Mosquitoes collected in the French Settlements in India. Ind Med. Gaz. lxiv, p. 495.
- BRODEMANN, H. W. 1919.—Sur quelques *Oulex* des Pyrénées.—II. Campagne 1918. Ann. Soc. Ent. France, lxxxviii, pp. 65-103
- BROOKS, G. E. 1922.—Tentative Table for the Diagnosis of Malayan Culicids. Govt. Health Dept., Straits Settlements, Serv. Publ. no. 2.

- BRUG, S L 1924 *a* —Notes on Dutch East Indian Mosquitoes. Bull. Ent Res xiv, pp 433-442.
- 1924 *b* —De Voornaamste Ned-Indische Culicinen Weltevreden
- 1926 —The Geographical Distribution of Mosquitoes in the Malayan Archipelago Meded Volks Ned.-Ind (Foreign Ed), pp 471-482
- 1931 *a* —Filariasis in the Dutch East Indies Proc Roy Soc, Med Sec., Trop Dis. & Parasit xxiv, pp 23-33
- 1931 *b* —Culiciden der Deutschen Limnologischen Sunda-Expedition Arch Hydrobiol Suppl-Bd ix, pp 1-42
- 1932 *a* —Notes on Dutch East Indian Mosquitoes Bull Ent Res. xxiii, pp. 73-83.
- 1932 *b* —Berichtigung zu meiner Arbeit "Culiciden" der Deutschen Limnologischen Sunda-Expedition Arch. Hydrobiol Suppl-Bd xi, pp 399-400
- BRUNETTI, E 1907.—Annotated Catalogue of Oriental Culicidæ. Rec. Ind Mus i, pp 297-377.
- 1912 —Annotated Catalogue of Oriental Culicidæ Suppl Rec Ind Mus. iv, pp 403-516.
- 1914 —Critical Review of the Genera in Culicidæ. Rec. Ind Mus. x, pp. 15-73
- 1920 —Catalogue of Oriental and South Asiatic Nemocera. *Ibid.* xvii, pp 1-300
- BUNTON, P A., and HOPKINS, G H E 1925 —The Early Stages of Samoan Mosquitoes Bull Ent Res xv, pp 295-301.
- 1927 —Researches in Polynesia and Melanesia —Parts I-IV. Pp 1-260. London Sch Hyg & Trop Med
- CARTER, H F. 1910 —Report on a Collection of Culicidæ from Cochun China Entomologist, xlii, pp 274-276.
- CHRISTOPHERS, S R 1906 —On the Importance of Larval Characters in the Classification of Mosquitoes. Sci Mem Med & San. Depts India, New Series, no. 25, pp 1-18
- 1911 —A new Culicine *Lesireomyia taeniorhynchoides*, nov gen et sp. Paludism, ii, pp. 68-72
- 1922 —The Development and Structure of the Terminal Abdominal Segments and Hypopygium of the Mosquito, with Observations on the Homologies of the Terminal Segments of the Larva Ind Journ Med Res x, pp. 530-572
- 1923 —The Structure and Development of the Female Genital Organs and Hypopygium of the Mosquito *Ibid* x, pp 698-720
- CHRISTOPHERS, S R, and BARRAUD, P J. 1923.—Descriptive Terminology of Male Genital Characters of Mosquitoes. *Ibid.* x, pp. 827-835
- CONNAL, S L M S 1927 —On the Variations occurring in *Aedes argenteus* in Lagos, Nigeria Bull. Ent Res. xviii, pp 5-11.
- COOLING, L E. 1924 —The Larval Stages and Biology of the commoner Species of Australian Mosquitoes. Commonwealth Austr. Dept. Health, Serv Publ. (Trop. Div) no. 8, pp 1-40
- CRUIKSHANK, J. A., and WRIGHT, R E 1914 —Filariasis in Cochun. Ind. Journ Med. Res. i, pp. 741-785.

- DOVER, C. 1928 — Notes on the Fauna of Pitcher-plants from Singapore Island Journ. Mal. Br. R. As. Soc. vi, pt. 3, pp. 1-27
- DYAR, H. G. 1920 — A Collection of Mosquitoes from the Philippine Islands Insec. Insc. Mens. viii, pp. 175-186
- 1921 — Two new Mosquitoes from China. *Ibid.* ix, pp. 147-148
- 1925 — Note on the Male of *Aedes punctifemora* Ludlow *Ibid.* xiii, p. 217
- 1928 — The Mosquitoes of the Americas Carnegie Inst., Washington
- DYAR, H. G., and SHANNON, R. C. 1925 — The Types of Philippine Mosquitoes described by Ludlow, and other Notes on the Fauna. Insec. Insc. Mens. xiii, pp. 66-89
- EDWARDS, F. W. 1911 — The African Species of *Culex* and Allied Genera Bull. Ent. Res. ii, pp. 241-266
- 1912 a — A Synopsis of the Species of African Culexidae other than *Anopheles*. *Ibid.* iii, pp. 1-53.
- 1912 b — Revised Keys to the known Larvae of African Culexidae *Ibid.* iii, pp. 373-385
- 1913 a — Further Notes on African Culexidae *Ibid.* iv, pp. 47-59
- 1913 b — New Synonymy in Oriental Culexidae *Ibid.* iv, pp. 221-242
- 1913 c — Tipulidae and Culexidae from the Lake of Tiberias and Damascus Journ. Proc. Asiatic Soc. Bengal, ix, pp. 47-61.
- 1914 a — On the Oriental Culexid Genus *Leicesteria*. Bull. Ent. Res. iv, pp. 255-263
- 1914 b — New Species of Culexidae in the British Museum, with Notes on the Genitalia of some African *Culex*. *Ibid.* v, pp. 63-81.
- 1914 c — New Culexidae from Borneo and Hong Kong *Ibid.* v, pp. 125-128
- 1915 a — New and little-known East African Culexidae *Ibid.* v, pp. 273-281.
- 1915 b — Diagnoses of new Bornean Culexidae *Ibid.* v, pp. 283-285.
- 1916 — Eight new Mosquitoes in the British Museum Collection. *Ibid.* vi, pp. 357-364.
- 1917 — Notes on Culexidae, with Descriptions of new Species *Ibid.* vii, pp. 201-220
- 1920 a — Mosquito Notes *Ibid.* x, pp. 129-137.
- 1920 b — The Nomenclature of the Parts of the Male Hypopygium of Diptera Nematocera, with special reference to Mosquitoes Ann. Trop. Med. & Parasit. xiv, pp. 23-40.
- 1921 a — H. Sauter's Formosan collections. Culexidae. Ann. & Mag. Nat. Hist. (9) viii, pp. 629-632.
- 1921 b — Mosquito Notes.—II. Bull. Ent. Res. xi, pp. 69-80
- 1921 c — A Revision of the Mosquitoes of the Palearctic Region. Bull. Ent. Res. xi, pp. 263-351.
- 1922 a — A Revision of the Genus *Harpagomyia*. Trans. Ent. Soc. London, January, pp. 496-503
- 1922 b — Mosquito Notes.—II. Bull. Ent. Res. xiii, pp. 75-102.
- 1922 c — A Synopsis of Adult Oriental Culexine Mosquitoes.—Part 1. Ind. Journ. Med. Res. x, pp. 249-293.
- 1922 d — A Synopsis of Adult Oriental Culexine Mosquitoes.—Part 2. *Ibid.* x, pp. 430-475

- EDWARDS, F W 1923 —Mosquito Notes —IV Ind Journ Med Res. xiv, pp 1-9
- 1924 —A Synopsis of the Adult Mosquitoes of the Australian Region *Ibid* xiv, pp 351-401
- 1925 a —Mosquito Notes —V. *Ibid* xv, pp 257-270
- 1925 b —Diptera Nematocera from the Dutch East Indies Treubia, vi, pp 154-172
- 1925 c —Fauna Buruana Culicidæ *Ibid* vii, p 40.
- 1926 a —Mosquito Notes —VI Bull Ent Res xvii, pp. 101-131
- 1926 b —Una revisione delle zanzare delle regioni Paleartiche Riv Malar. v, pp 253-285, 392-466, 613-652
- 1928 a —Mosquito Notes —VII Bull Ent. Res. xviii, pp 267-284
- 1928 b —Diptera Nematocera from the Federated Malay States Museums. Journ F M S Mus xiv, pp. 1-139
- 1929 a —Philippine Nematocerous Diptera —Part 2 Not Ent ix, pp 1-14
- 1929 b —Mosquito Notes —VIII Bull Ent. Res. xx, pp 321-343
- 1930 a —Mosquito Notes.—IX *Ibid* xxi, pp. 287-306
- 1930 b —Mosquito Notes —X *Ibid*. xxi, pp 541-545
- 1930 c —Notes on Exotic Chaoborinae, with Descriptions of new Species (Diptera, Culicidæ). Ann & Mag Nat Hist (10) vi, pp 527-540
- 1931 —Mosquitoes breeding in Plant Pitchers Nat Hist. Mag iii, pp. 25-28.
- 1932 —Genera Insectorum Fasc cxciv.—Family Culicidæ. Brussels
- EDWARDS, F W, and GIVEN, D H. C. 1928 —The early Stages of some Singapore Mosquitoes Bull Ent Res xviii, pp 337-357
- EVANS, A. M 1929 —Descriptions of the early Stages of two further Mosquitoes collected in Southern Nigeria by Mr L H Dunn Ann Trop. Med & Parasit. xxiii, pp 407-413.
- FARQUHARSON, C O 1918-19 —*Harpagomyia* and other Diptera fed by *Cremastogaster* Ants in Southern Nigeria Proc Ent. Soc London, 1918, pp xxix-xxxix.
- FICALBI, E 1897 —Revisione sistematica delle specie Europee della Famiglia delle Culicidæ. Bull. Soc Ent. Ital xxviii, pp 197-313
- 1899.—Venti specie di zanzare (Culicidæ) Italiane *Ibid*. xxxi, pp 46-234
- GALLIARD, H 1927 —Note sur les Culicines de Corse Ann. Parasit hum & comp pp 97-104.
- GIL COLLADO, J 1930 —Datos actuales sobre la distribución geográfica de los culicoides españoles Eos, Rev Españ Ent vi, pp 329-347
- GILES, G M 1900 —A Handbook of Gnats or Mosquitoes 1st Ed London.
- 1901 a —A Plea for the Collective Investigation of Indian Culicidæ, with suggestions as to moot points for enquiry, and a Prodrömus of Species known to the Author. Journ Bomb. Nat. Hist. Soc. xiii, pp 592-610
- 1901 b.—Six new Species of Culicidæ from India Entomologist, xxxiv, pp. 192-197.
- 1902 —A Handbook of Gnats or Mosquitos. 2nd Ed. London,

- GILES, G. M. 1903.—Notes on the Culicids of Dehra Dun, with a Description [by Giles] of a new *Mansonia* which mimics *Anopheles*. By Major F. Wyville Thomson, communicated by G. M. Giles. Journ. Trop. Med. vi, p. 315.
- 1904.—Notes on some Mosquitoes, etc., received from the Philippine Islands and Angola, with some incidental remarks upon Classification. *Ibid.* vii, pp. 366-369.
- GRAHAM, W. M. 1911.—Results obtained from a Monthly Examination of the Native Domestic Water Receptacles at Lagos, Southern Nigeria, in 1910-1911. Bull. Ent. Res. ii, pp. 127-139.
- GRUBB, E. E. 1905.—On *Toxorhynchites immanis*, the Elephant Mosquito. Spol. Zeyl. ii, pp. 158-164.
- GUENTHER, K. 1913.—Die lobenden Bewohner der Kannen der insekten-fressenden Pflanze *Nepenthes distillatoria* auf Ceylon. Zeitschr. für wissenschaftl. Insekten-Biol. ix, pp. 198-207 & 259-270.
- HAGA, J. 1924.—Aantockinging omtrent muskieten (II). Geneesk. Tijds. Ned.-Ind. (5) lxiv, pp. 1-20 (sep.).
- HILL, G. F. 1925.—The Distribution of Anopheline Mosquitoes in the Australian Region, with Notes on some Culicine Species. Proc. Roy. Soc. Vic. xxxvii, pp. 61-77.
- HINDLE, E. 1929.—An experimental Study of Yellow Fever. Trans. Roy. Soc. Trop. Med. & Hyg. xxii, pp. 405-434.
- HOWARD, L. O., DYAR, H. G., and KNAB, F. 1912.—The Mosquitoes of North and Central America and the West Indies. Washington (I. Structure & Bionomics, II. Plates).
- 1915.—The Mosquitoes of North and Central America and the West Indies—III. (Systematic Part 1).
- 1917.—The Mosquitoes of North and Central America and the West Indies—IV. (Systematic Part 2).
- INGRAM, A. 1912.—Notes on the Mosquitoes observed at Bole, Northern Territories, Gold Coast. Bull. Ent. Res. ii, pp. 73-78.
- INGRAM, A., & MACFIE, J. W. S. 1917 a.—Notes on some distinctive Points in the Pupae of West African Mosquitoes. *Ibid.* viii, pp. 73-91.
- 1917 b.—The early Stages of certain West African Mosquitoes. *Ibid.* viii, pp. 135-154.
- 1919.—The early Stages of West African Mosquitoes—IV. *Ibid.* x, pp. 59-69.
- INGRAM, A., & DE MEILLON, B. 1927.—A Mosquito Survey of certain Parts of Southern Africa, with special reference to the Carriers of Malaria and their Control. Pub. S. Afr. Inst. Med. Res. no. 22, pp. 1-81.
- IVENGAB, M. O. T. 1932.—Filariasis in North Travancore. Ind. Journ. Med. Res. xx, pp. 671-672.
- JACOBSON, E. 1909 a.—Ameisen aus Java und Krakatau—II. Biologischer Theil. Notes. Leyden Museum, xxxi, p. 246.
- 1909 b.—Ein Moskito als Gast. Tijds. v. Ent. iii, pp. 158-164.
- 1911.—Nahere Mitteilungen über die myrmecophilen Culiciden *Harpagomyia splendens*. *Ibid.* liv, pp. 158-161.
- JAMES, S. P. 1900.—On the Metamorphosis of the *Filaria sanguinis hominis* in Mosquitoes etc. Ind. Med. Gaz. Sept. pp. 340-343.
- 1914.—Summary of a Year's Mosquito Work in Colombo. Ind. Journ. Med. Res. ii, pp. 227-266.

- JOYEUX, C 1918—Note sur les Culicoides de Macédoine Bull. Soc. Path. Exot. xi, pp. 530-547
- KIRKPATRICK, T. W. 1925.—The Mosquitoes of Egypt Pp. 1-224 Cairo
- KORKE, V. T. 1928.—Observations on Filariasis in some Areas in British India Ind. Journ. Med. Res. xvi, pp. 187-198 & 695-715
- 1929.—Observations on the Atypical Variety of *bancrofti* and its Significance. *Ibid.* xvi, pp. 1023-1031.
- 1930 a.—Observations on the Correlation between the Incidence of Filarial Infection in the Human Host and in the Carrier in relation to the Terrain. *Ibid.* xvii, pp. 319-331
- 1930 b.—Observations on the Characters of Filarial Endemic Areas in Bihar and Orissa. *Ibid.* xvii, pp. 333-336
- 1930 c.—Observations on the Natural History of *Filaria bancrofti* in Dwellings in relation to the Systems of Drainage. *Ibid.* xvii, pp. 427-442.
- 1932.—Observations on Filariasis in some Areas in British India. *Ibid.* xx, pp. 335-339.
- KUMM, H. W. 1931 a.—Studies on *Aedes* Larvæ in South-Western Nigeria, etc. Bull. Ent. Res. xxii, pp. 65-74
- 1931 b.—Geographical Distribution of Yellow Fever Vectors Amer. Journ. Hyg. Mono. ser. no. 12, pp. 1-110
- LANG, W. D. 1920.—A Handbook of British Mosquitoes British Museum
- LANGERON, M. 1918.—Morphologie et biologie de la larve de *Theobaldia spathopalpos* Bull. Soc. Path. Exot. xi, pp. 98-103.
- LEICESTER, G. F. 1904 (in F. V. Theobald).—New Culicidæ from the Federated Malay States Entomologist, xxxvii, pp. 238-239
- 1908.—The Culicidæ of Malaya Studies from Inst. Med. Res. F.M.S. iii, pp. 18-261
- LUDLOW, C. S. 1904 a.—Mosquito Notes Can. Ent. xxxvi, pp. 233-236
- 1904 b.—Mosquito Notes.—2 *Ibid.* xxxvi, pp. 297-301.
- 1905 a.—Mosquito Notes.—3 *Ibid.* xxxvii, pp. 94-102
- 1905 b.—Mosquito Notes.—3 (cont.) *Ibid.* xxxvii, pp. 129-135.
- 1905 c.—Mosquito Notes.—4 *Ibid.* xxxvii, pp. 385-388
- 1906.—Mosquito Notes.—5 *Ibid.* xxxviii, pp. 360-368
- 1907.—Mosquito Notes.—5 (cont.) *Ibid.* xxxix, pp. 413-414
- 1910.—Mosquito Observations *Ibid.* xlii, pp. 193-196
- MACFARLANE, H. 1915.—The *Stegomyia* Survey in Hong Kong. Bull. Ent. Res. vi, pp. 67-68
- MACFIE, J. W. S. 1914.—A Note on the Action of Common Salt on the Larvæ of *Stegomyia fasciata* *Ibid.* iv, pp. 339-344.
- 1915.—Observations on the Bionomics of *Stegomyia fasciata* *Ibid.* vi, pp. 205-229
- 1917.—Morphological Changes observed during the Development of the Larvæ of *Stegomyia fasciata* Bull. Ent. Res. vii, pp. 297-307
- 1920 a.—The Chatotaxy of the Pupa of *Stegomyia fasciata*. *Ibid.* x, pp. 161-169
- 1920 b.—The Early Stages of West African Mosquitoes—V. (contains Observations on Chatotaxy of Pupa of *Culex*). *Ibid.* xi, pp. 105-112.

- MACFIE, J. W. S., and INGRAM, A. 1916 *a*—Now Culicoides Larva from the Gold Coast. *Ibid.* vii, pp 1-18
- 1916 *b*—The Domestic Mosquitoes of Accra. *Ibid.* vii, pp. 161-177.
- 1922—On the Genital Armature of the Female Mosquito. *Ann Trop Med & Parasit* xvi, pp 167-188.
- 1923—The Early Stages of West African Mosquitoes—VI. *Bull. Ent Res* xiii, pp 409-442
- MACGREGOR, M. E. 1916.—Resistance of the Eggs of *Stegomyia fasciata* to Conditions adverse to Development. *Ibid.* vii, pp 81-85.
- MARTINI, E. 1923 —Über einige für das System bedeutungsvolle Merkmale der Stechmücken (Haarstellung der Larven, mannliche Geschlechtsorgane, Kiefertaster) *Zool Jahrb. Abt. f. System.* no. 6, pp 517-590.
- 1929-31 (in Erwin Lindner).—Die Fliegen der paläarktischen Region. *Culicoides*. Pp. 1-144 (1929), 145-320 (1930), 321-398 (1931).
- MEGAW, J. W. D. 1923—The Dengue-Sandfly-Fever Problem. *Ind. Med. Gaz.* lviin, pp 401-406.
- MEGAW, J. W. D., and GUPTA, J. S. 1927 —The Geographical Distribution of some of the Diseases of India. *Ibid.* lxi, pp 299-313
- DE MEILLERE, J. C. H. 1909 —Die myrmecophile Dipteren aus Java. *Tijd. v. Ent.* li, pp 165-174
- 1910.—*Nepenthes*-Tiere—I. Systematik. *Ann. Jard. Bot. Buitenzorg* (2), Supp. iii, pp 917-940
- 1911 —Zur metamorphose der myrmecophilen Culicoides *Harpagomyia splendens*. *Tijd. v. Ent.* liv, pp 162-167
- DE MEILLON, 1928 —Notes on some Mosquitoes found in South Africa.—I. *S. Afr. Journ. Sci.* xxv, pp 316-324
- MONTCHADSKY, H. S. 1928.—Larva of *Aedes (O) pulchritarsis*, Rond., var. *stegomyia*, Stack. & Montch., nov., from Turkestan. *Bull. Ent. Res.* xvii, pp 151-157.
- MONTCHADSKY, A. 1930 —Die Stigmalplatten der Culicoides-Larven. *Zool. Jahrb. Syst.* lviii, pp 541-636
- NEVEU-LEMAIRE, M. 1905 —Mission du bourg de Bozas. Description d'une nouvelle espèce de *Stegomyia* recueillie par le Dr Brumpt à Harar. *Bull. Soc. Zool. Franco.* xxx, pp. 8-11
- NOË, G. 1899 —Contribuzioni allo studio dei Culicoidi. *Bull. Soc. Ent. Ital.* xxxi, pp 235-262.
- PAINÉ, R. W., and EDWARDS, F. W. 1929.—Mosquitoes from the Solomon Islands. *Bull. Ent. Res.* xx, pp. 303-316.
- PAIVA, C. A. 1910.—Notes on the Larva of *Toxorhynchites unimacricornis*, Rec. Ind. Mus. v, pp 187-190
- 1912.—Materials for a Survey of the Mosquitoes of Calcutta. *Ibid.* vii, pp. 93-98.
- PATTON, W. S. 1905—The Culicoid Fauna of the Aden Hinterland, their Haunts and Habits. *Journ. Bomb. Nat. Hist. Soc.* xvi, pp. 623-637.
- PEUS, F. 1933.—Zur Kenntnis der *Aedes*-Arten des deutschen Faunengebietes (Dipt., Culicoid.). Die Weibchen der *Aedes communis*-Gruppe. *Konowia*, xii, pp 145-159.
- PRASHAD, BAJNI. 1915.—Male Generative Organs of some Indian Mosquitoes. *Ind. Journ. Med. Res.* iii, pp 497-502.

- PRASAD, BAINI 1918 —Contributions to the Anatomy of Aquatic Diptera—I Larval and Pupal Stages of an Indian *Chaoborus* and *Dixa*. Rec Ind Mus. xv, pp 153-158.
- RAO, S S, and IYENGAR, M O T. 1930 —Studies on the Influence of Season on the Development of *Filaria bancrofti* in *Culex fatigans* Ind. Journ Med Res xvii, pp. 759-768
- . 1932 —Experimental Infection of some Indian Mosquitoes with *Wuchereria (Filaria) bancrofti*. *Ibid* xx, pp 25-34
- ROY, S. K, and BOSU, S. C 1922.—Filariasis in Puri Ind. Med Gaz lvi, pp. 281-286.
- SEGUY, E 1924 —Les Moustiques de l'Afrique Mineure, de l'Egypte et de la Syrie Pp. 1-257 Encycl Ent, Paris.
- 1925 a —Faune de France —12 Dipt., Culicidae Pp. 13-82 Paris
- 1925 b —Notes sur les Moustiques de l'Afrique Mineure, de l'Egypte et de la Syrie —I Encycl. Ent., Diptera, i, pp 13-21
- 1925 c —Notes sur les Moustiques —I & II *Ibid*. pp. 120, 157-158
- SENIOR-WHITE, R 1919-1927 See WHITE, R SENIOR
- STRICKLAND, C. 1917 —A curious Adaptation of Habit to its Environment of a Malayan Mosquito. Journ Straths Branch Roy Asiatic Soc no 75.
- THEOBALD, F V. 1901 a.—A Monograph of the Culicidae of the World—1. British Museum
- 1901 b.—A Monograph of the Culicidae of the World.—2. British Museum.
- 1903 a —A Monograph of the Culicidae of the World —3 British Museum
- 1903 b —New Culicidae from the Federated Malay States. Entomologist, xxxvi, pp 256-259
- 1904 a —A new Culicid Genus from Uganda Journ Trop. Med. vii, pp 17-18
- 1904 b.—The Mosquitoes of Egypt, the Sudan, and Abyssinia. First Rept Well. Trop. Res. Lab pp 62-83 Dept Educ. Sudan Govt, Khartoum
- 1904 c —New Culicidae from the Federated Malay States. Entomologist, xxxvii, pp 77-78.
- 1904 d —New Culicidae from the Federated Malay States *Ibid* xxxvii, pp. 163-165
- 1904 e.—New Culicidae from the Federated Malay States. *Ibid*. xxxvii, pp 236-239.
- 1905 a.—A Catalogue of the Culicidae in the Hungarian National Museum, with Descriptions of new Genera and Species. Ann. Mus Nat Hung. iii, pp. 61-120.
- 1905 b.—A new Genus of Culicidae Entomologist, xxxviii, pp. 52-56.
- 1905 c.—Genera Insectorum Fasc xxvi —Family Culicidae.
- 1905 d —New Culicidae from India, Africa, British Guiana, and Australia. Journ Econ. Biol. i, pp 17-36.
- 1905 e.—Some new Mosquitoes from Ceylon. Journ. Bomb. Nat. Hist Soc. xvi, pp. 237-250.
- 1906 —Second Report Wellcome Tropical Research Laboratories. Report on Economic Entomology, pp. 67-83. Dept. Educ. Sudan Govt., Khartoum.
- 1907.—A Monograph of the Culicidae of the World.—4. British Museum.

- THEOBALD, F. V. 1908.—First Report on the Collection of Culicidae and Colethridae in the Indian Museum, Calcutta, with Descriptions of new Genera and Species. *Rec. Ind. Mus.* ii, pp. 287-302.
- 1909.—Descriptions of new Mosquitoes collected by Dr. Graham in Ashanti. *Col. Off. Rept. Miscell.* no. 237. [Unpublished.]
- 1910 a.—The Culicidae of Fiji, including two new Species. *Entomologist*, xliii, pp. 155-159.
- 1910 b.—A Monograph of the Culicidae of the World.—5. British Museum.
- 1910 c.—Second Report on the Collection of Culicidae in the Indian Museum, Calcutta, with Descriptions of new Genera and Species. *Rec. Ind. Mus.* iv, pp. 1-33.
- 1911.—Culicidae of the R. Zool. Soc. "Natura Artis Magistra," Amsterdam, and Descriptions of three new Species. *Tijds. v. Ent.* liv, pp. 233-240.
- THOMBOR, O. 1924.—Pupa of some Palestine Culicines. *Bull. Ent. Res.* xiv, pp. 341-345.
- VOGEL, R. 1920.—*Aedes pulchritarsis* ein Baumhohlenbrutor. *Int. Rev. Hydrobiol.* xxi, pp. 161-170.
- WATERSTON, J. 1918.—On the Mosquitoes of Macodoma. *Bull. Ent. Res.* ix, pp. 1-12.
- WESCOMB, W. 1910.—On the Larval and Pupal Stages of West African Culicidae. *Bull. Ent. Res.* i, pp. 7-50.
- WESSENBURG-LUND, C. 1921.—Contributions to the Biology of the Danish Culicidae. *Mém. Acad. R. Sci. & Lettres, Copenhagen, Sec. Sci.* 8th Series, vii, no. 1, pp. 1-210.
- WHITE, R. SENIOR 1919.—*Toxorhynchites minimus*. *Spol. Zeyl.* xi, pp. 189-191.
- 1920.—A Survey of the Culicidae of a Rubber Estate. *Ind. Journ. Med. Res.* vii, pp. 304-325.
- 1922.—Notes on Indian Diptera. *Mem. Dept. Agric. India, Ent. Series*, vii, pp. 83-169.
- 1923.—Catalogue of Indian Insects.—Part 2. Culicidae. Pp. 1-124. Calcutta.
- 1925.—Notes on Ceylon Mosquitoes.—I. *Spol. Zeyl.* xii, pp. 213-222.
- 1927.—Notes on Ceylon Mosquitoes.—II. The Larva of the commoner non-*Anopheles* Mosquitoes. *Spol. Zeyl.* xiv, pp. 61-76.
- WIGGLESWORTH, V. B. 1929.—The early Stages of some West African Mosquitoes. *Bull. Ent. Res.* xx, pp. 59-68.
- YAMADA, S. 1921.—Descriptions of Ten new Species of *Aedes* found in Japan, with Notes on the Relation between some of these Mosquitoes and the Larva of *Phlebotomus bancrofti*. *Ann. Zool. Jap.* x, Article 6, pp. 45-81.

SYSTEMATIC INDEX.

	Page	Uranotomia (cont.)	Page
Subfam CULICINÆ		23. unguiculata <i>Edw.</i> . . .	67
Tribe MEGARHININI . . .	8	24. campastis <i>Leic.</i> . . .	68
1. <i>Megarhinus R.-D.</i> . . .	8	25. macfarlanei <i>Edw.</i> . . .	70
1. <i>albipes Edw.</i> . . .	10	26. longirostris <i>Leic.</i> . . .	71
2. <i>kempii Edw.</i> . . .	18	27. atra <i>Theo.</i> . . .	72
3. <i>gravelyi Edw.</i> . . .	20	28. testacea <i>Theo.</i> . . .	74
4. <i>minimus Theo.</i> . . .	22	29. annandalei <i>Barr.</i> . . .	75
5. <i>edwardsi Barr.</i> . . .	23	30. myrpleura <i>Leic.</i> . . .	76
6. <i>splendens (Wied.)</i> . . .	24	31. maxima <i>Leic.</i> . . .	77
Tribe CULICINI . . .	28	32. bimaculata <i>Leic.</i> . . .	78
2. <i>Tripteroides Giles.</i> . . .	34	33. luteola, sp. n. . . .	80
7. <i>similis (Leic.)</i> . . .	38	34. stricklandi <i>Barr.</i> . . .	80
8. <i>powelli (Ludl.)</i> var. <i>indicus Barr.</i> . . .	39	35. hebes <i>Barr.</i> . . .	81
9. <i>edwardsi (Barr.)</i> . . .	41	36. maculipleura <i>Leic.</i> . . .	82
10. <i>araneoides (Theo.)</i> . . .	42	37. bicolor <i>Leic.</i> . . .	83
var. <i>serrata Barr.</i> . . .	42	38. recondita <i>Edw.</i> . . .	83
11. <i>affinis (Edw.)</i> . . .	44	39. novobscura, sp. n. . .	84
12. <i>dofleini (Günther)</i> . . .	45	7. <i>Theobaldia N.-L.</i> . . .	86
13. <i>Topomyia</i> . . .	45	40. longiareolata <i>Mey.</i> . . .	88
13. <i>aureoventer (Theo.)</i> . . .	46	41. mveitannata (<i>Theo.</i>) . .	91
4. <i>Harpagomyia de Meij.</i> . . .	47	42. indica <i>Edw.</i> . . .	93
14. <i>genurostris (Leic.)</i> . . .	48	8. <i>Orthopodomyia Theo.</i> . . .	94
15. <i>jacobsoni Edw.</i> . . .	51	43. anopheloides (<i>Stiles</i>) . .	98
5. <i>Hodgesia Theo.</i> . . .	52	var. <i>maculata Theo.</i> . . .	101
16. <i>malayi Leic.</i> . . .	54	var. <i>maculipes Theo.</i> . . .	101
17. <i>baillyi Barr.</i> . . .	55	var. n. <i>andamanensis</i> . . .	102
6. <i>Uranotomia L. A.</i> . . .	56	44. <i>albipes Leic.</i> . . .	102
18. <i>alboannulata (Theo.)</i> . .	62	45. <i>flavithorax Barr.</i> . . .	103
19. <i>rutherfordi Edw.</i> . . .	63	46. <i>flavicoxa Barr.</i> . . .	104
20. <i>christophersi Barr.</i> . . .	64	9. <i>Picalbia Theo.</i> . . .	105
21. <i>edwardsi Barr.</i> . . .	65	Subg. <i>Mimomyia Theo.</i> . . .	108
22. <i>orientalis Barr.</i> . . .	66	47. <i>chamberlaini (Ludl.)</i> . .	108
		48. <i>hybrida (Leic.)</i> . . .	111
		Subg. <i>Etoleptomyia Theo.</i> . .	113
		49. <i>luzonensis (Ludl.)</i> . . .	113
		50. <i>fusca (Leic.)</i> . . .	115

	Page	Subg. Finlaya (cont)	Page
Subg. Ficalbia, s. str.	116	94. annulensis <i>Edw.</i>	198
51. minima (<i>Theo</i>)	116	95. pulchrisenter (<i>Giles</i>)	199
10. <i>Mansonia</i> <i>Blanch</i>	118	96. antoni <i>Barr</i>	200
Subg. Coquillettia <i>Dyar</i>	119	97. auronitens <i>Edw.</i>	202
52. crassipes (<i>v d W</i>)	120	98. sub similis <i>Barr</i>	203
53. novochracea (<i>Barr</i>)	121	99. dissimilis (<i>Leic</i>)	204
54. ochracea (<i>Theo.</i>)	122	var. karwan <i>Barr</i>	204
Subg. <i>Mansonioides</i> <i>Theo.</i>	123	100. albolateralis (<i>Theo</i>)	205
55. annulifera (<i>Theo</i>)	127	101. niveus (<i>Ludl</i>)	208
56. longipalpis (<i>v d W</i>)	128	102. alboniveus, sp. n. . . .	210
57. uniformis (<i>Theo</i>)	129	103. niveoides, sp. n. . . .	211
58. indiana <i>Edw.</i>	130	104. novoniveus, sp. n. . . .	211
11. <i>Aedomyia</i> <i>Theo</i>	131	Subg. <i>Christophersomyia</i> <i>Barr.</i>	212
59. venustipes (<i>Stuese</i>)	132	105. thomsoni (<i>Theo</i>)	213
12. <i>Aedes</i> <i>Meig.</i>	134	106. annulirostris (<i>Theo</i>)	215
Sub <i>Mucidae</i> <i>Theo</i>	144	107. ibis <i>Barr.</i>	215
60. scatophagoides <i>Theo</i>	145	Subg. <i>Rhinoskeusa</i> <i>Edw.</i>	216
61. laniger (<i>Wied</i>)	147	108. longirostris (<i>Leic</i>)	216
Subg. <i>Ochlerotatus</i> (<i>Theo</i>)	147	Subg. <i>Stegomyia</i> <i>Theo</i>	217
62. caspius (<i>Pall.</i>)	148	109. egypti (<i>Lam</i>)	221
63. pulchritarsis (<i>Rond</i>)	150	110. desmotus <i>Giles</i>	225
var. asiaticus <i>Edw.</i>	151	111. annandalei <i>Theo.</i>	227
var. versicolor <i>Barr.</i>	151	var. quadrimacatus	
64. pullatus (<i>Cog</i>)	152, 442	<i>Barr.</i>	228
Subg. <i>Finlaya</i> <i>Theo.</i>	153	112. craggi <i>Barr.</i>	229
65. poecilus <i>Theo.</i>	157	113. mediopunctatus <i>Theo.</i>	230
66. gubernatoris (<i>Giles</i>)	159	var. perplexus <i>Leic.</i>	231
67. deccanus <i>Barr.</i>	163	var. submediopunctatus <i>Barr.</i>	231
68. fessgraderi, sp. n. . . .	164	var. n. surensis	231
69. cogilli <i>Edw.</i>	165	114. edwardsi <i>Barr.</i>	232
70. asameusis (<i>Theo.</i>)	166	115. w-albus <i>Theo.</i>	232
71. caelharanus <i>Barr.</i>	166	116. albopictus (<i>Stuese</i>)	233
72. lophoventralis (<i>Theo.</i>)	167	117. pseudalbopictus <i>Borel</i>	235
73. khazani <i>Edw.</i>	168	118. novalbopictus <i>Barr.</i>	237
74. prominens <i>Barr.</i>	169	119. subalbopictus <i>Barr.</i>	238
75. inquinatus <i>Edw.</i>	169	120. flavopictus <i>Yam.</i>	239
76. uncinatus <i>Edw.</i>	170	121. scutellaris (<i>Wall</i>)	240
77. albocinctus <i>Barr.</i>	172	122. umineatus (<i>Theo.</i>)	242
78. stevensoni <i>Barr.</i>	174	123. albolineatus (<i>Theo</i>)	243
79. albetennatus (<i>Leic.</i>)	174	124. vittatus (<i>Big.</i>)	245
var. mikimnus <i>Edw.</i>	176	Subg. <i>Asdimorphus</i> <i>Theo.</i>	246
80. macdougalli <i>Edw.</i>	177	125. jamesi (<i>Edw</i>)	249
81. pseudotennatus (<i>Giles</i>)	178	126. lowisi <i>Theo.</i>	250
82. elaine <i>Barr.</i>	180	127. alboscuteallatus (<i>Theo.</i>)	250
83. shortti <i>Barr.</i>	183	128. niveoscuteallum <i>Theo</i>	251
84. greeni (<i>Theo</i>)	184, 442	129. culicinus <i>Edw.</i>	252
85. chrysolinatus (<i>Theo.</i>)	185	130. vexans (<i>Meig</i>)	253
86. harveyi <i>Barr.</i>	188	131. stenotus (<i>Theo.</i>)	255
87. formosensis <i>Yam</i>	189	132. syntheticus <i>Barr.</i>	256
88. pallirostus <i>Edw.</i>	190	133. caesus (<i>Theo</i>)	257
89. saxicola <i>Edw.</i>	191	134. pipersulatus (<i>Giles</i>)	258
90. oreophilus (<i>Edw.</i>)	192	135. teniorhynchoides (<i>Chr.</i>)	260
91. suffusus <i>Edw.</i>	194	136. pallidostriatus (<i>Theo.</i>)	261
92. christophersi <i>Edw.</i>	195	137. nigrostriatus <i>Barr.</i>	262
93. gilli <i>Barr.</i>	196		

Subg. Aedimorphus (cont)	Page	Subg. Armingeres (cont)	Page
138. mediocinctus (Theo.)	263	183. kuchingensis Edw.	317
139. trunculatus (Theo.)	264	184. theobaldi, nom. n.	319
140. nummatus Edw.	265	185. amecolus (Leic.)	319
141. ostentatio (Leic.)	267	Subg. Leicesteria Theo.	320
Subg. Indusius, nov.	268, 445	186. flavus (Leic.)	321
142. pulverulentus Edw.	268	187. magnus (Theo.)	324
Subg. Baucanella Theo.	269	188. annulitarsis Leic.	325
143. lineatopennis (Ludl.)	269	189. annulipalpis (Theo.)	327
Subg. Diceromyia Theo.	271	190. inchoatus Barr.	328
144. periscaletus (Giles)	272	191. longipalpis (Leic.)	329
145. punctipes Edw.	273	192. digitatus Edw.	330
146. iyengui Edw.	273	193. omnesus Edw.	330
147. micropterus Giles	275	194. dentatus Barr.	331
148. regine Edw.	277	16. Culex Linn.	332
Subg. Aodes Meig.	277	Subg. Lutzia Theo.	338
149. uniformis (Theo.)	281	195. fuscus Wied.	341
150. abditus Barr.	282	196. raptor Edw.	343
151. yusak Barr.	283	197. vorax Edw.	344
152. indicus (Theo.)	283	198. halifax Theo.	344
153. pseudomedioscutus Theo.	286	Subg. Beauveridia Edw.	345
154. cautus Barr.	288	199. modestus Flo.	346
155. ceylonicus Edw.	288	Subg. Neoculex Dyar.	347
156. andamanensis Edw.	290	200. brevipalpis (Giles)	348
157. vallistris Barr.	290	201. tenuipalpis Barr.	351
158. hirsutipleura Barr.	291	Subg. Mochtiogenes Edw.	352
159. conatus Barr.	292	202. iphis Barr.	354
160. agrestis Barr.	293	203. castrensis Edw.	355
161. yerburyi Edw.	293	204. pluvialis Barr.	356
162. atrius Barr.	293	205. khazam Edw.	357
163. lugubris Barr.	294	206. malayi (Leic.)	358
164. olivatus Barr.	294	Subg. Lophoceratomyia Theo.	359
165. pseudolumus Theo.	295	207. minutissimus (Theo.)	363
166. rami Barr.	295	208. semoni, sp. n.	365
167. sigmoides Barr.	295	209. cinctellus Edw.	366
168. butleri Theo.	296	210. rubithoracis Leic.	367
Subg. Canacraea Edw.	297	211. fraudatrix (Theo.)	368
169. canericomes Edw.	298	212. minor Leic.	370
170. simplex (Theo.)	298	var. bengalensis, nov.	371
13. Heuzmannia Ludl.	299	213. plantaginis Barr.	372
171. complex (Theo.)	302	214. uniformis Theo.	373
172. funerea (Leic.)	302	215. mamillifer Leic.	374
173. indica (Theo.)	303	216. flavicornis Barr.	375
174. covelli Barr.	304	Subg. Onchomyia Theo.	376
175. viridis Barr.	306	217. viridiventer Giles	378
176. himalaysensis Edw.	300	218. shebbearei Barr.	380
177. metallica (Leic.)	307	219. pallidothorax Theo.	381
178. chandi Edw.	308	220. bailvi, sp. n.	382
179. greeni (Theo.)	308	221. nigropunctatus Edw.	383
14. Hæmagogus Will.	309	222. fragilis Ludl.	385
180. discrepans Edw.	309	223. bahri Edw.	386
181. tripunctatus (Theo.)	310	Subg. Culex Linn.	387
15. Armigeres Theo.	310	224. epidesmus (Theo.)	389
Subg. Armigeres Theo.	313	225. bitæmorhynchus Giles	391
182. obturbans (Walk.)	314	var. ambiguus Theo.	393
		var. tsunax Theo.	398

Subg. <i>Culex</i> (cont.)	Page	Subg. <i>Culex</i> (cont.)	Page
226. <i>suenensis</i> Theo	394	237. <i>minulus</i> Edw.	412
227. <i>cornutus</i> Edw	396	238. <i>ulgaricus</i> Edw.	413
228. <i>edwardsi</i> Barr	397	239. <i>thenlei</i> Theo	414
229. <i>sitonsi</i> Wied.	398	240. <i>vagens</i> Wied	416
230. <i>vishnu</i> Theo	400	241. <i>univittatus</i> Theo	418
231. <i>whitei</i> Barr.	402	242. <i>fatigans</i> Wied	420
232. <i>barraudi</i> Edw.	403	243. <i>hutchinsoni</i> Barr. . . .	423
233. <i>tritaniorhynchus</i> Giles. .	404	244. <i>fuscocephalus</i> Theo. . .	424
234. <i>whitmorei</i> (Giles)	406	245. <i>fuscitarsis</i> Barr . . .	426
235. <i>gelidus</i> Theo.	407		
236. <i>mimeticus</i> Noël.	409, 451		
var <i>minuloides</i> Barr.	411		

APPENDIX.

Subfam. DIXINÆ	Page	Subg. <i>Sayomyia</i> Coq	Page
1. <i>Dixa</i> Meig.	430	2. <i>asiaticus</i> Giles	440
Subg. <i>Dixa</i> , s. str.	431	3. <i>annandalei</i> Edw	441
1. <i>maculipennis</i> Brun.	431	4. <i>indicus</i> Giles	441
2. <i>trinitata</i> , sp. n.	432		
3. <i>montana</i> Brun.	433	Subfam. CULICINÆ.	
4. sp.	433	Tribe CULICINI.	
5. sp.	434	(Addenda.)	
6. <i>ochroleuca</i> Brun	434	<i>Aedes</i> Meig.	442
7. <i>zeylanica</i> White	434	Subg. <i>Aedimorphus</i> Theo. . . .	443
8. <i>biatriata</i> Brun.	435	128 <i>a punctifemore</i> Ludl. . . .	443
9. <i>bifasciata</i> Brun.	435	Subg. <i>Cancraedes</i> Edw.	443
10. <i>christophersi</i> , sp. n.	435	170 <i>a. kanarensis</i> , sp. n. . . .	444
11. <i>platystyla</i> , sp. n.	436	<i>Paraedes</i> , gen. n	446
Subg. <i>Paradixa</i> Tonm.	436	170 <i>b. barraudi</i> , sp. n	447
12. <i>kaashnurensis</i> , sp. n. . . .	437	170 <i>c argyrurus</i> , sp. n. . . .	448
Subfam. CHAOBORINÆ.		<i>Culex</i> Linn.	450
1. <i>Corethrella</i> Coq.	438	Subg. <i>Culex</i> , s. str.	451
1. <i>inepta</i> Ann.	439	236 <i>a. minuloides</i> Barr. . . .	451
2. <i>Chaoborus</i> Licht.	440	236 <i>b. fuscifurcatus</i> , sp. n. . .	452

ABBREVIATIONS USED IN TEXT.

Adult †.

- apn* Anterior pronotal lobe (prothoracic lobe).
ppn . . . Posterior pronotal lobe (propuneron).
af Anterior or upper forked cell of wing.
pf Posterior or lower forked cell of wing.

Larva

- asc* Inner (submedian dorsal) caudal hair of anal segment.
osc Outer (submedian dorsal) caudal hair of anal segment.
lh Lateral hair on saddle of anal segment.
 I-VIII .. Abdominal segments.

Distribution

An asterisk (*) denotes specimens examined by Author.

† For explanation of lettering on figures of ♂ hypopygia see p. 4;
 for numbering of wing-veins see pp. 3, 57.

ERRATA.

- Page 19. Line 8 from bottom, *for* hair 8 *read* hair 7
- „ 43. „ 2 of first footnote, *for* serrate *read* straight.
- „ 120. „ 9 from bottom, *for* *pygmeus* *read* *pygmaeus*
- „ 183. Paragraph on distribution, *for* Wuch'ang, HUPEN, *read*
Kiukiang, KIANGSI
- „ 340. Line 14, *for* p. 348 *read* p 343

Order DIPTERA.

Family CULICIDÆ.

Tribes MEGARHININI and CULICINI.

INTRODUCTION.

I. THE INDIAN AREA.

The area, as here dealt with, is shown on the map at the end of this volume. It extends from the western boundaries of Kashmir, the North-West Frontier Province, and Baluchistan, along the chain of the Himalayas, to the eastern boundaries of Assam and Burma. From the northern boundary of the Himalayas it extends southwards to include the Indian Peninsula and Ceylon, also the Andaman, Nicobar, Laccadive, and Maldive groups of islands.

Within the area many varying conditions of climate prevail—from desert conditions in Sind to areas of very heavy precipitation in the north-east and south, where the annual rainfall may be as heavy as 500 inches per annum.

II. DISTRIBUTION.

Mosquitoes are common over practically the whole area, and in many parts occur in enormous numbers. They have been collected at altitudes of 14,000 feet or more in Kashmir, and *Culex fatigans* has been found at 3,760 feet below ground-level (903 feet below sea-level) in the Kolar Gold Mines in South India.

The distribution of genera and species within the area, and beyond its limits, is given in the systematic part of this work

The mosquito fauna is mainly western Oriental in character, but there is an admixture of Palaearctic and Ethiopian species in the north and west, and of Malayan or eastern Oriental forms in the southern, eastern, and north-eastern regions.

of heavy rainfall. Several species have a world-wide distribution through the tropical and subtropical zones, such as *Aedes aegypti* and *Culex fatigans*, these two being mainly domestic. *Aedes vexans* is found in the Nearctic, Palearctic, and Oriental regions, and ranges as far eastwards as some of the Pacific islands. Some species, on the other hand, appear to be very local, such as those of *Aedes (Finlaya)*, which are known only from the Western Himalayas (possibly representing a Central Asian element in the fauna), and those of *Aedes (Aedes)*, known only from Northern Bengal or Assam.

For more general considerations regarding faunal areas, and conditions governing the distribution of mosquitoes, reference should be made to the companion volume on the Anophelini by Colonel Sir Rickard Christophers.

III. CLASSIFICATION

The arrangement of tribes, genera, and subgenera is in accordance with that adopted by Edwards (1932), the older classifications of Theobald and others having been considerably modified as a result of very intensive study of the subfamily by numerous workers during the past thirty years. It seems unlikely that any considerable alterations will be found necessary in the future.

IV CHARACTERS USED IN IDENTIFICATION AND CLASSIFICATION.

The nomenclature used for the various parts is, so far as possible, in accordance with that adopted in the companion volume on the Anophelini, and for explanation of most of the terms used reference should be made to the Introduction to that volume.

ADULT—Points of value in identification are as follows:—

Head character and coloration of scaling, and arrangement of bristles; form of bucco-pharyngeal armature of ♀ (present only in genus *Culex* and its various subgenera). **Antenna**—coloration of torus, and presence or absence of scales—the form of this part is also of importance in *Culex (Lophocerotomyia)*, length of antenna compared with that of proboscis, form of flagellar segments (e g., in *Aedomyia*, *Heizmannia*, etc.), and character of hair-whorls—scale-tufts often present in ♂♂ of *Culex (Lophocerotomyia)*. **Clypeus**: shape (as distinguishing the Megarhinini from Culicini, and genus *Harpagomyia* from others); presence or absence of scales. **Palpi and proboscis**: length, shape, ornamentation.

Thorax: shape—somewhat produced over the head in some species of *Armigeres (Leicesterna)*; scaling, ornamentation,

and chaetotaxy of mesonotum and scutellum, size of pronotal (prothoracic) lobes—usually enlarged in *Hæmagogus* and *Heizmannia*, character of scaling, if any, on posterior pronotum, arrangement of bristles on pleuræ (this is of great importance, and is shown in fig 7, p 29 The important groups of thoracic bristles are.—

- (a) *Anterior pronotal*, on pronotal (prothoracic) lobes.
- (b) *Posterior pronotal* (pro-epimeral), on the part of the side of thorax above pronotal lobe and in front of anterior spiracle.
- (c) *Propleural* (prosternal), on the part above front coxa and below anterior pronotal (prothoracic) lobe.
- (d) *Spiracular*, a row immediately in front of anterior spiracle, separated from posterior pronotal by a slight ridge forming posterior border of posterior pronotum
- (e) *Postspiracular*, on membranous part of pleuræ behind anterior spiracle (especially characteristic of the *Aedes* group)
- (f) *Sternopleural*, a more or less continuous row on posterior border of sternopleura.
- (g) *Prealar*, on a small knob immediately below and in front of wing-root
- (h) *Upper mesepimeral* (subalar), immediately below and behind wing-root, on upper part of mesepimeron, and in front of posterior spiracle.
- (k) *Lower mesepimeral*, on middle or lower part of mesepimeron
- (l) *Postnotal*, a tuft of fine hairs on postnotum.

Note position of meron (a triangular plate below mesepimeron) in relation to hind coxa.

Wings: venation or arrangement of veins. The notation of veins, as used in this volume, is shown in fig. 15, p. 57. The veins and branches are referred to by numbers, e. g., 2, 2.1, 2.2 indicate respectively the 2nd longitudinal vein (R_{2+3} *), the anterior branch (R_2), and the posterior branch (R_3) of this vein. Cross-veins are referred to by numbers also, e. g., that joining the 3rd (R_{4+5}) and 4th (M) longitudinal veins is called c.-v. 3-4 ($r-m$), indicating that it lies between veins 3 and 4. Note especially length of vein 6 and position of its termination in relation to bifurcation of vein 5; length of anterior forked-cell; size of microtrichia (small hairs on wing-membrane, minute in *Uranotania*); coloration

* As in the Comstock-Needham system of notation, used by some authors.

and shape of scales on veins, length of wing from base of costa to tip (as indicating size of a particular species), presence or absence of hairs on squama (see fig. 7, g, p. 29), and of hairs on upper and underside of wing at base (latter present only in *Theobaldia*)

Legs: ornamentation, especially presence or absence of pale rings or bands on tarsal segments, if present, whether basal or apical, or both, comparative length of tibia and of tarsal segments, presence or absence of outstanding scales, tarsal claws, whether toothed or simple (fig. 7, d-f), presence or absence of pulvilli (developed only in *Culex* and its subgenera) (Under a binocular microscope the pulvilli appear pale against a dark background)

Abdomen: ornamentation of dorsum and venter and character of scaling (tufts on terminal segments in some species of *Megarhinus*, or on venter in some species of *Aedes* (*Finlaya*))

Hypopygium structure of terminalia of ♂♂ of importance in all genera, nomenclature of parts shown in figs. 6, 35, 48, 49, 81, etc. Explanation of lettering of parts —

- 9r, 9s, ninth tergite and sternite
 - c, coxite (side-piece)
 - s, style (clasper)
- BL, basal lobe or process of coxite
- SL, or SA, subapical lobe of coxite
- EXA, external apodeme of coxite
- BA, basal apodeme of phallosome and coxite
- H, harpago (basal appendage of coxite in *Aedes*)
- HF, harpaginal fold, connecting the two harpagines
- EHF, eminence on harpaginal fold (rudimentary harpago)
- PG, proctiger or anal segment.
- PT, or VLP, paraproct (lateral or ventro-lateral chitination of anal segment)
 - OR, crown of paraproct
 - DP, dorsal plate of proctiger (tenth tergite).
 - LA, basal lateral arm of paraproct (in *Culex*)
 - PH, phallosome (mesosome or aedeagus)
 - LP, lateral plate of phallosome (when chitinations are paired)
- BF, EXP, MP, and VC, basal, external, and median processes and ventral cornu of lateral plate of phallosome (in *Culex*).
- PM, parameral plate

Terminalia of ♀♀ of special importance in *Aedes* (*Aedes*) and *Mansonia* (*Mansonioides*), nomenclature of parts shown in figs. 30, 70, 72

PUPA.—General structure shown in fig. 8 of the volume on *Anophelini*

Points of value in identification are —

Form of respiratory trumpet, arrangement of hairs on abdominal segments, shape of paddles

The pupæ of the Culicini, with the exception of one or two genera, have not been closely studied, but, where material has been available, brief descriptions have been given

LARVA —Points of value in identification. —

Character and arrangement of *frontal hairs*. The author has followed the notation proposed by Edwards and Given (1928, p. 338) for the five important pairs of hairs on the fronto-clypeus of the larvæ of the Culicini, and these are referred to respectively as *A*, *B*, *C*, *d*, and *e*, the first three are usually larger and more conspicuous than the first two, and are homologous with the three pairs of frontal hairs of Anophelini, hair *d* being homologous with the posterior clypeal and *e* with the occipital. The position of these varies greatly in different genera, as will be seen by referring to figs 31, b, 34, d, 42, etc. Characters of mouth-brush hairs, preclypeal spines, and of mentum. *Antennæ* length and thickness of shaft, position and character of shaft-hair or tuft, and of subapical bristles, presence or absence of spines or spicules on shaft. Chaetotaxy of *thorax* (see figs 3, 9, 39, 102) and abdomen. Form of *comb* and structure of teeth of comb on abdominal segment VIII. Length of *siphon-tube* compared with diameter at base. Presence or absence of an *acus* (small extension of chitin) at base of siphon-tube on each side (usually present in subgenus *Finlaya* and in genus *Culex*). Presence or absence of *pecten* on siphon, and, if present, number and form of teeth. Number and arrangement of hairs on siphon, and any modification of valves at tip of siphon. Extent of chitination of *anal segment*, characters of subdorsal (outer and inner submedian caudal), lateral, and fan-hairs, length and form of anal papillæ.

Egg —This stage of Megarhinini and Culicini has been very little studied at present, but references to available descriptions have been given in the systematic part. Eggs may be laid singly or cemented together in rafts: they differ from those of Anophelini in the absence of lateral floats.

V. BIONOMICS AND RELATION TO DISEASE.

Information on bionomics is given in the systematic part of this work, under tribes, genera, and species.

The only human diseases prevalent in India which are known to be transmitted by mosquitoes are malaria, dengue, and filariasis. The Megarhinini, being non-blood-sucking forms, are not concerned in the transmission of disease, and no species of the Culicini are known to transmit malaria. Very little work on dengue, in connection with mosquitoes, has been done in India, but *Aedes aegypti* and *Aedes albopictus* have been shown to be carriers in other parts of the world, and these two, which are both very common, and possibly other allied species,

most probably transmit the disease in India. *Culex fatigans*, one of the commonest domestic mosquitoes in most parts of the country, appears to be the chief carrier of filariasis. Other Indian species which may be concerned in the transmission of this disease are *Mansonia longipalpis*, *Aedes ægypti*, *Aedes scutellaris*, *Culex vishnui*, and certain species of *Anopheles*.

Yellow fever has not, up to now, occurred in India, but the Indian race of *Aedes ægypti* has been shown to be capable of transmitting this disease. Other common Indian species have also been shown to be potential carriers, e.g., *Aedes albopictus* and *Aedes vittatus*.

It is not possible to quote all references to this subject. Abstracts of papers will be found in the 'Tropical Diseases Bulletin,' and in the 'Review of Applied Entomology.' The following may be cited as referring to India.—James, 1900, Cruikshank & Wright, 1914, Roy & Bose, 1922, Megaw, 1923, Megaw & Gupta, 1927, Barraud, 1928 c, Korke, 1928 a, 1928 b, 1929, 1930 a, 1930 b, 1930 c, 1932, Hindle, 1929, Rao & Iyengar, 1930, 1932, Acton & Rao, 1931; Kumm, 1931 b. (For full references, see Bibliography at end of this volume.)

VI SYNONYMY AND REFERENCES

The synonymy given in the systematic part is that which seems to be agreed upon at the present time. For considerations of space only Oriental synonyms are included. Full synonymy will be found in Edwards (1932). It is most probable that some alterations will be found necessary in the future, when the life-histories of more species have been worked out. Some of the forms at present treated as varieties will perhaps be allotted specific rank, and *vice versa*.

References to original descriptions have been given in all cases. Endeavour has been made to include also all the more important recent systematic references, but it has been thought unnecessary to repeat the names of older works, as the descriptions are now, for the most part, of little value, or merely copies of the originals to which references have been given.

SYSTEMATIC.

Characters distinguishing the Tribes Megarhinini and Culicini

ADULTS.

Proboscis rigid, apical $\frac{1}{2}$ more slender and bent downwards, or hooked (fig 1), posterior margin of scutellum evenly rounded, clypeus broader than long, with front margin slightly trilobed, a V-shaped thickening in hind margin of wing between branches of vein 5 [p. 8.
MEGARHININI,

Proboscis more flexible, usually of uniform thickness, but sometimes swollen at tip (figs. 10, 12), not hooked, posterior margin of scutellum more or less trilobed, and with three distinct groups of bristles, clypeus longer than broad, rounded above and in front, no V-shaped thickening in hind margin of wing [p. 28
CULICINI,

PUPÆ.

Outer part of paddle produced beyond termination of midrib, hind border not serrated and without terminal hair, but finely fringed (fig. 5); respiratory horn not ending in a chitinous spine, nor divided nearly to the base into leaflets MEGARHININI.

Outer part of paddle not usually produced beyond termination of midrib (fig. 24, etc.), but should it have this appearance the hind margin is either serrated, or with terminal hair (fig. 22), or respiratory horn ends in a chitinous spine (fig. 31), or is divided nearly to the base into leaflets (fig 14, a). CULICINI.

LARVÆ.

Mouth-brushes each composed of 10 stout lamellæ, or flattened bristles, placed in a regular row; abdominal segment VIII without comb, but with lateral chitinous plate on each side carrying two plumose bristles MEGARHININI.

Mouth-brushes composed of numerous hairs; if these are modified into lamellæ there are about 30 or more in each brush, not arranged in a regular row; abdominal segment VIII with comb, and with or without lateral chitinous plates CULICINI.

Tribe MEGARHININI.

According to the most recent classification, this tribe includes only one genus, *Megarhinus*, which has been divided into three groups by Edwards (1932). Two of these groups are purely American, the third being confined to the Old World. The description given below applies to this last group, which corresponds to Theobald's genus *Toxorhynchites*.

[In this description the characters of adult and larva which specially distinguish *Megarhinus* from mosquitoes of the tribe Culicini are printed in italics.]

Genus MEGARHINUS Robineau-Desvoidy, 1827

Mem Soc d'Hist Nat III, p 403 Genotype, *Culex hæmorrhoidalis* Fab

Toxorhynchites Theobald, 1901, M.C 1, p 244. Genotype, *T brevipalpis* Theo

Worcesteria Banks, 1906, Phil J Sci 1, p 779 Genotype, *W grata* Banks.


Teromyia Leicester, 1908, Cul Malaya, p. 49 Genotype, *T acan-data* Leic

ADULT*.—Large or very large mosquitoes, usually with metallic blue, yellow, purple, or white flat scales, lateral tufts of coloured hairs on terminal abdominal segments in some species, *clypeus broader than long, with front margin slightly trilobed*; *proboscis curved downwards and backwards in the middle, basal $\frac{1}{2}$ rigid and usually thick, apical $\frac{1}{2}$ more slender, tapering to apex, and flexible, proboscis not adapted for piercing or sucking blood*, bucco-pharyngeal armature absent; palpi of ♂ about length of proboscis, and composed of 5 segments, segment 1 very small, 5 long and tapering to tip, 4 and 5 upturned, all segments covered with flat scales, some short spines on terminal segments, but no hair-tufts, palpi of ♀ of varying length in different species, and from about $\frac{1}{4}$ – $\frac{1}{2}$ length of proboscis, composed of 3 segments, of which the terminal one is very small and hidden by scales; antennæ of ♂ plumose, those of ♀ setose, *scutellum not lobed, posterior margin slightly and evenly rounded*, *anterior fork-cell of wing very short*; *a V-shaped thickening in wing-membrane between branches of vein 5*; squamal fringe absent,

* Edwards 1921 c, p. 284, 1932, p. 58, Howard, Dyar, & Knab, 1917, p. 928; Barraud 1929 b, p. 271; Barraud & Covell 1928, p. 676 (buccal cavity), Martini 1929, p. 114

thoracic bristles reduced, no posterior pronotal or post-spiracular, none on mesonotum except laterally, only two pairs of orbital, but a row of strong spiracular bristles present, fore and mid-tarsal claws of ♂ unequal, the larger with one tooth, those on hind legs, and on all legs of ♀, simple

Fig 1




Tribe Megarhinini, adult characters: a, wing-venation; b, clypeus, with base of antenna; c, scutellum, d, side view of ♀ (*M. splendens*), wings and legs omitted; e, head of ♂; f, central portion of wing, *M. kempi*; g, ditto, *M. splendens*.

Hypopygium of ♂* of simple structure and very similar in all Indian species. 9th sternite broadly crescentic and not specialized; coxite with a terminal style and with a hairy

* Edwards 1920 b, pp. 23-40; Christophers 1922, pp. 530-571; 1923, pp. 698-720, Macfie & Ingram 1922, p. 164, Barraud 1929 b, p. 271.

basal lobe on anal surface connected with that of the other side by the harpaginal fold; lateral plates of phallosome long and narrow, parameral plate and basal apodeme moderately

Fig 2



Larva of *Megarhinus splendens* a, whole larva, b, antenna, c, mentum; d, dorsal spine of metathorax, e, lateral plate of VIII, f, mouth-brush; g, head from above

developed, paraproct (chitimsation of anal segment or proctiger) blackened at crown and terminating in a tooth with a few minute hairs immediately below. Hypopygium of ♀ of simple type also; cerci comparatively broad and

short, postgenital plate lobe-like and not usually emarginate at apex, insula with a number of hairs arranged in a median group, three chitinated spermathecae present

Pupa.—Respiratory horn about $2\frac{1}{2}$ –3 times as long as the greatest width in side view, expanded towards tip, opening oval and oblique, in antero-posterior view the basal part is rather swollen, with a slight constriction at base of opening. Dendritic tufts on abdominal segment I well developed. The five submedian and sublateral hairs on tergites II–VII form a group towards the lateral margin, innermost always small and simple, other four varying according to the species, dorsal and lateral hairs on tergite VIII both small and usually simple. Paddles large and of characteristic shape in each species, a dark irregular mark across bases of paddles except in one species, midrib strong, no terminal hair or spine, posterior margin fringed with delicate hairs, outer part produced beyond termination of midrib (this is not the case in the Culicini, except in two or three genera, and in these the respiratory horn terminates in a strong spine, or is divided nearly to the base into leaflets).

LARVA *—Very uniform in structure throughout the genus, usually of a deep red or reddish-brown colour, those of the larger species measuring up to 16 mm in length. Head subquadrate, hairs on dorsal surface all placed very far forward, *fronto-clypeus* divided by a suture into an anterior part and a much larger posterior part, the anterior carries four hairs on each side, three fine and simple and one very minute and spine-like (the outermost), the posterior part also carries four hairs on each side, three fine and simple and one very small and branched (the innermost)†. Antenna short, shaft smooth, hair-tuft represented by two fine hairs situated some little distance below the tip, between these and the tip there is a small branched subapical hair. Mouth-parts adapted for predaceous purposes, hairs of mouth-brushes modified into ten strong lamellae with hooked tips. Thorax with some heavily chitinated plates carrying thick spinulose bristles (arrangement shown in diagrammatic form in fig. 3). Prothorax with two large plates on each side, a dorso-lateral and ventro-lateral; the former carries hairs 5–7 (5 and 7 being bristles), the latter carries hair 8, always small and branched, and the four pleural hairs (9–12) on a raised tubercle, one of these being a bristle. Mesothorax bearing three or four plates on each side, dorso-

* Barraud 1931 a, p. 1127, Green 1906, p. 160; Senior-White, 1927, p. 66; Edwards 1932, p. 58.

† [The four posterior hairs seem to correspond with hairs A, B, C, and d of Culicini, hair e being absent. The four anterior hairs presumably represent the preclypeals and anterior clypeals of Anophelini.]


lateral plate bearing hairs 2-6 (5 being a bristle), plate sometimes divided, hairs 2 and 3 being on a separate small plate, hair 7 lateral, bristle-like, arising from a tubercular plate, the ventro-lateral plate carries the pleural hairs 9-12 and hair 8, as on the prothorax. Metathorax with thirteen hairs as against fourteen on the pro- and mesothorax, hairs 1a and 14 both missing, hairs 5, 7, and 13 are bristles arising from separate plates, the last being ventral, pleural hairs and hair 8 on a ventro-lateral plate, as on the other two segments. In addition to the large plates mentioned, there are some much smaller plates, from which the finer hairs arise*. On abdominal segments I-VII there are three plates on each side, a dorso-lateral, lateral, and ventro-lateral, which carry either spinulose bristles or long plumose hairs, or both, these are very similar in all species except the dorso-lateral plate on VII, on which there may be two bristles and three hairs, or one bristle and four hairs. *Segment VIII with a large lateral plate on each side in place of a comb, with two strong bristles and some small hairs.* Siphon-tube short and wide, a single pair of strong branched hair-tufts arising near base, pecten absent. Anal segment enclosed in a strongly chitinated ring, with numerous spines along the posterior margin, both long and short irregularly alternating, lateral hair bristle-like, spinulose, both pairs of subdorsal hairs divided into a number of long branches, anal papillae quite short, anal fan well developed, the hairs split into fine branches.

* Dr F W Edwards has suggested altering the numbers of the hairs on the thorax from those given by me in 1931 in order to harmonize these with the notation used by Puri and Christophers for Anopheles (adopted here also for Culicini, figs. 9, 39, and 102). He has also pointed out that hair no. 8 (previously numbered 13 by me) is on the dorsal side of the plate carrying the pleural hairs (9-12), as in fig 3—not on the ventral side, as shown in my previous drawing. Dr Puri has very kindly re-examined material with me at Kasauli. We have found some difficulty in deciding on the probable correct numbering, for, as pointed out by Edwards, the thoracic chaetotaxy of Megarhinini seems to differ more from Anophelini and Culicini than these do from one another. The numbers in fig 3 are those which appear now to be the most satisfactory, and are in accord with those proposed by Edwards, except that we designate the median prothoracic hairs 1, 1a, 2, and 3 instead of 1, 2, 3, and 0, respectively, as we think that hair 0 (of Anophelini) is almost certainly absent in Megarhinini; the ventral prothoracic hair we designate 14 instead of 13. It appears to us that in Megarhinini hair 13 is absent from the prothorax and hair 14 from the mesothorax, another peculiarity is the association of hairs 8-12 on the same plate on each segment of the thorax (though 8 is separate on the prothorax in some species, e. g., *M. albipes*).

BIONOMICS *—The adults fly by day and visit flowers for the purpose of sucking honey. The majority of the species are found only in forest or jungle. The ♂♂ are said to congregate on leaves of bushes, forming a stationary swarm.

In India the larvæ are most frequently found in water in tree-holes or bamboos, where they prey upon and devour the larvæ of other mosquitoes living in the same situations. The larva of *M. splendens* (Wied.) is sometimes found in domestic collections of water, such as water-butts or earthenware jars in the vicinity of human dwellings, particularly on the outskirts of towns, and cities such as Calcutta. It is

Fig 3



Megarhinus splendens Wied.: chetotaxy of right side of thorax of larva. For numbering of hairs see footnote on opposite page. Hairs nos 9-12 are pleural hairs, hair no. 13 of prothorax and no 14 of mesothorax are apparently absent. Chitinous plates: DL, dorso-lateral, L, lateral, VL, ventro-lateral, and V, ventral. Mid-dorsal line at top of block, mid-ventral line at bottom.

unusual to find more than one fully-grown larva in a bamboo or tree-hole, but sometimes in a water-butt, large earthenware jar, or hollow tree containing a quantity of water several larvæ may be found living together. In such situations

* Green 1905, p. 161, Paiva 1910, p. 187; Howard, Dyar, & Knab 1912, pp. 109, 113, 118, 131; Semor-White 1920, p. 319; Edwards 1932, p. 58; Brug 1932, p. 73.

the larvæ do not appear to attack one another so long as larvæ of other species be present, but in the absence of the latter they may do so, and may even seize and suck dry pupæ of their own species.

The larva rests quietly at the surface of the water, sometimes almost horizontally, at other times in a more vertical position, and captures other larvæ with the mouth-brushes, when they approach within reach, by a rapid sideways jerk of the head and thorax

The eggs are roundish and slightly oval, they are dropped singly on water while the insect is in flight, and float buoyantly by means of air-bubbles which form amongst tubercles or spines present on the egg-shell

DISTRIBUTION —Tropics of both the Old and New Worlds, a few species being found also in the north temperate zone. About 50 species have been described, of which about 20 are American Six have, so far, been found in India, two being peculiar to that area, the other four having a distribution further east Several of the Indian species occur in the Himalayas, two being found at elevations of 7,000 feet or more

Key to Adults

- 1 Tarsi entirely dark (♂♀); small and slender species, wing-length less than 5 mm *minimus*, p. 22
Some tarsal segments with white markings, large species, wing-length usually 6 mm or more. 2
2. All three terminal tarsal segments of all legs with white markings or entirely white *albipes*, p. 16.
One or more of the three terminal tarsal segments on fore or hind legs without white markings 3
- 3 Mesonotum with conspicuous border of pale yellow scales continuous from front, along sides, over wing-roots, to scutellum *edwardsi*, p. 23
Mesonotum without a continuous pale border.. 4
- 4 Abdomen with conspicuous lateral tufts of hairs on VI–VIII, those on VII black, on VIII orange, scales on mesonotum dull bronzy- or greenish-brown except laterally *splendens*, p. 24
Abdomen without conspicuous lateral tufts of hairs, scales on mesonotum brilliant metallic green 5
- 5 Dorsum of abdomen mainly coppery purple, with narrow blue basal bands, segments 2 and 3 of mid-tarsi of ♀ mainly purple, 5th segment of hind tarsi white, segment 1 of hind tarsi of ♂ with dense bristles beneath *kempi*, p. 18
Dorsum of abdomen mainly deep blue or deep purple, with lateral pale yellow markings; segments 2 and 3 of mid-tarsi of ♀ white, segment 5 of hind tarsi dark, segment 1 of hind tarsi of ♂ without dense bristles beneath . *gravelys*, p. 20

Key to Pupæ

1. Paddle elongate, $1\frac{1}{2}$ times to twice as long as greatest width, all 5 inner hairs* near the posterior margin of tergite VII fine and short, hardly projecting beyond posterior border of that segment 2
 - Paddle wide and rounded, less than $1\frac{1}{2}$ times as long as greatest width, one of the 5 inner posterior hairs of tergite VII very long and stout, reaching beyond posterior margin of VIII 5.
2. Apical margin of paddle not emarginate nor produced into a point *gracilis*.
 - Apical margin of paddle emarginate or produced into a blunt point 3
3. One of the 5 inner posterior hairs of tergite V long and stout, the four others slender and comparatively short 4
 - Two of the 5 inner posterior hairs of tergite V long and stout, the three others quite small *albipes*.
4. One of the 5 inner posterior hairs of tergite IV long, the four others quite small, paddle $1\frac{1}{2}$ times as long as greatest width; apical margin emarginate and rounded, an irregular dark line across base *minimus*
 - Two of the 5 inner posterior hairs of tergite IV long and stout, the three others quite small, paddle twice as long as greatest width, apex produced into a blunt point, no irregular dark line across base *kemp*.
5. Larger hair on VII long and stout, usually much longer than tergite VIII, often as long as greatest width of paddle *splendens*.
 - Larger lateral hair on VII short and fine, not as long as tergite VIII *edwardsi*

Key to Larvæ (4th stage).

1. Mesothoracic dorso-lateral plate undivided 2.
 - This plate divided 4.
2. Dorso-lateral plate of VII carrying 2 bristles and 3 hairs *splendens*,
 - This plate carrying 1 bristle and 4 hairs 3.
3. Bristle 5 on prothoracic dorso-lateral plate only about $\frac{1}{2}$ the length of bristle 7, both usually single (5 occasionally bifid), mesothoracic dorso-lateral plate with a projection carrying hairs 2 and 3, hair 6 on this plate with 3 to 5 branches; chitinisations of head and body light brown or yellow *kemp*.
 - Bristles 5 and 7 on prothoracic dorso-lateral plate about the same length, 5 single, 7 bifid; mesothoracic dorso-lateral plate without a marked projection, hair 6 on this plate nearly always simple, chitinisations of head and body deep brown *edwardsi*.

* The expression "inner hairs" is used to include the five sub-median and sublateral hairs as distinct from the lateral.

MEGARHININI

Siphon distinctly shorter and smaller than chitinised part of anal segment, longer bristle on lateral plate of VIII more than $\frac{1}{2}$ length of siphon (not including valves), chitinations of head and body deep brown	<i>albipes</i>
Siphon as long as, or longer than, chitinised part of anal segment, longer bristle on lateral plate of VIII less than $\frac{1}{2}$ length of siphon, chitinations of head and body light brown or yellow	5
Small, siphon about length of chitinised part of anal segment, or slightly longer, hair 6 on mesothoracic dorso-lateral plate usually simple	<i>minimus</i>
Large, siphon longer than chitinised part of anal segment, hair 6 on mesothoracic dorso-lateral plate with 3 or 4 branches	<i>gracilis</i>

1. *Megarhinus albipes* Edwards, 1922

Ind Journ Med Res x, p 287 (♂ & ♀) TYPE-LOC Simla (Jakko), W Himalayas, 7,000 to 8,000', viii 1915 (*Christophers*)
TYPE: ♂ and allotype ♀ in Brit Mus

ADULT*—A fairly large species without conspicuous lateral tufts on terminal segments of abdomen. Leg-markings distinctive, all tarsi marked with white to a greater or less extent†. Wing 8-9.5 mm

♀—*Head-scales* dull dark purple dorsally, white laterally, usually a narrow white border to eye-margins, *proboscis* purple on stout basal part, bluish or greenish apically, *palpi* purple, reaching to base of 5th flagellar segment of antenna. *Mesonotal* and *scutellar scales* brilliant metallic green, reddish when seen in certain positions, some dark ones intermixed over wing-roots and on scutellum, scales on *apn* pale golden green, those on upper part of *ppn* purple, those on lower part and on *pleurae* silvery. *Abdominal tergites* dark blue to purple, with usually narrow white basal bands on II-V, not always complete, V-VII with pale yellow lateral patches usually visible dorsally, tergite I purple in middle, pale yellow or white at sides, segments VI and VIII with pale yellow lateral hairs, VII with black, not forming conspicuous tufts, *tergites* pale golden, with a narrow longitudinal purple stripe. *Wings* slightly brownish, membrane distinctly clouded in the region of c-vs 2-3, 3-4, and 4-5, these three‡ being fairly


* Barraud 1929 b, p 273; Senior-White 1923, p 43, footnote.

† [A closely allied species is *M. manicatus* Edw. (Formosa), which differs from *M. albipes* in having the 5th tarsal segments white.]

‡ [In *Megarhinus* (as in *Anopheles*) the base of vein 3 is short and transverse and appears like a cross-vein, it is here for convenience referred to as "c-v 2-3". In this genus, moreover, the radio-medial cross-vein (3-4) is right angled, with a more or less extended horizontal portion and a horizontal stump arising from the angle, "c-v. 3-4" here refers to the vertical portion only.]

closely approximated. *Legs* purple, with golden and white markings as follows —Femora golden at bases and beneath, fore and mid-tibiae with pale scaling posteriorly, 1st tarsal segment on all legs with basal pale ring, most distinct on mid and hind pair, apical $\frac{1}{2}$ of segment 2, and whole of

Fig 4



Larval structures of *Megarhinus*. a, *M. edwardsi* Barr., dorso-lateral plates of prothorax (above), mesothorax (middle), and abd. seg VII (below); b, *M. albipes* Edw., dorso-lateral plate of mesothorax; c, *M. kempii* Edw., dorso-lateral plates of pro- and mesothorax; d, *M. splendens* Wied., dorso-lateral plate of VII; e, *M. gravehys* Edw., siphon, chitinated part of anal segment, and (above) longer bristle of lateral plate of VIII, same scale as siphon; f, same structures in *M. minimus* Theo; g, same in *M. albipes* Edw.

3-5 on fore legs white, except part of 5 in some specimens, 2-5 on mid-legs entirely white, white scaling at apices of 2 and 3 on hind legs (in some specimens 2 is almost entirely dark), 4 entirely white, also base of 5

♂—Palpi slender and nearly as long as proboscis, purple, with yellow scaling at apex of segment 2, on sides of 3, and underside of 4, segment 5 nearly twice as long as 4, antennæ plumose, a small scale-tuft on 1st flagellar segment. Segment VII of abdomen with fairly long and dense lateral black hairs forming small tufts, lateral golden hairs on VIII

Hypopygium. 9t narrow, apical border not produced into lobes; lp with a few small teeth, very similar to that of *M. kempi*, style similar to that of *M. splendens*

PUPA † (fig 5, a)—Resembles that of *M. graveleyi* closely, but distinguished by shape of paddle, which is emarginate on the apical margin. Distinguished from pupæ of other species by characters given in key

LARVA (fig 4, b) (*vide* key).—The two parts of the mesothoracic dorso-lateral plate may occasionally be united by a narrow strip of chitin, but in all larvæ examined it is completely divided into two on one or both sides of the thorax. Approximate length (fully grown) 12-13 mm; siphon 0.73-1.0 mm; siphonal tuft with 4-7 branches.

HABITAT—Tree-holes.

DISTRIBUTION.—WESTERN HIMALAYAS Simla*, 7-8,000', viii. 1915 (*Christophers*), viii & ix 1923 (*Barraud*), viii. 1927 (*Puri*), ix., x, xi. 1930 (*Barraud*); Krol Mt.*, near Solan, 7,000', vii, viii, ix. 1923, ix. 1924, vii, viii. 1930 (*Barraud*), Dungagah*, Hazara dist, 7,000', v 1915, and Muktesar*, Almora dist, 7,000', v 1923 (*T. B. Fletcher*).

Outside India recorded from COCHIN CHINA (*Borel*)

2 *Megarhinus kempi* Edwards, 1921

Bull. Ent. Res. xii, p. 72 (*Megarhinus (Toxorhynchites) kempi*) (♂ and ♀, ♂ hypop.). TYPE-LOC. Telewadi, near Castle Rock, North Kanara dist, 3-10 x. 1916 (*S. Kemp*). TYPE. 1 ♂ and 1 co-type ♀ in Ind. Mus.; co-type ♀ in Brit. Mus.

ADULT †.—The leg-markings and coloration of abdomen serve to distinguish this species §, the ♂ differs from that of

* Throughout this volume an asterisk against a distribution record indicates that specimens from the locality have been examined by the author.

† Barraud 1931 a, pp. 1130-1131

‡ Barraud 1929 b, p. 274.

§ *M. kempi* closely resembles *M. leucesteri* Theo. from Malaya, the latter differing chiefly in having segments 2-5 of mid-tarsi of ♀ white, 9t of ♂ hypopygium broader, and lp without serrations or teeth.

other Indian species in the presence of close-set bristles on underside of 1st hind tarsal segment. Wing about 6 mm.

♀—*Head-scales* blue, greenish or purple, according to direction of light, those along eye-margins lighter, *antennae* with moderately long sparse hairs; *palpi* purple, reaching to about tip of 5th flagellar segment, *proboscis* with straight basal part very stout and purple-scaled, curved apical part bluish-green or bronzy. *Mesonotal scales* brilliant green, scales on *scutellum* and over wing-roots bright bluish-green, on *apn* and upper part of *ppn* purple, on lower part of *ppn* and on *pleurae* silvery. *Wings* without any definite clouding of membrane, c.-vs 2-3, 3-4, and 4-5 closely approximated (fig 1, f). *Legs*. femora purple dorsally, pale golden ventrally, small knee-spots of blue and white scales, tibiae purple, some light blue scales intermixed on mid-pair, tarsi purple, with white markings showing some variation, a subbasal white ring on segment 1 on all legs, usually white scaling at base of segment 4 on fore leg, segment 4 of mid-tarsi white, also larger part of 5, sometimes white scaling at base of 2 and apex of 3, segments 4 and 5 of hind tarsi mainly white, with a few dark scales at the joint. *Abdominal tergites* mainly coppery purple, with narrow basal bright blue bands on II-IV or II-VI, and basallateral silvery markings produced on to dorsum on IV-VI, tergite I bluish-green in middle, pale yellow at sides; sternites pale golden, IV purple in middle.

♂—*Palpi* about as long as *proboscis*, first four segments mainly yellowish, 5th purplish, *antennae* plumose; tarsi purple except for a pale ring near base of segment 1 on all legs, that on fore legs not always complete, segment 1 of hind tarsi with close-set bristles beneath for nearly the whole length. *Hypopygium* (fig. 6, b): 9t narrow, submedian hairy lobes not well developed; style narrow, with fairly long appendage, *lp* with a small number of minute teeth.

PUPA † (fig 5, c).—Paddle of very distinctive shape and without an irregular dark line across the base (in these characters closely resembling *M. leicesteri* Theo of Malaya).

LARVA (fig. 4, c) †.—Distinctive characters given in key; length of siphon 1.0-1.2 mm., siphonal tuft of 4-6 branches, bristle representing hair 8 on prothoracic dorso-lateral plate single (in some other species this bristle is invariably bifid).

HABITAT.—Bamboos.

DISTRIBUTION.—SOUTH INDIA: Nagargali*, Belgaum dist., viii. 1921, and Kadra and Yellapur*, N. Kanara dist., ix. & x. 1921 (*Barraud*); also type-locality* given above.

Outside India recorded from COCHIN CHINA (*Borai*); JAVA (*Brug*).

3. *Megarhinus gravelyi* Edwards, 1921

Bull Ent Res xii, p 73 (♂, ♂ hypop) TYPE-LOC Pashok, Darjeeling dist, 2,000', E. Himalayas, 26 v-14 vi 1916
(*F H Graveli*) TYPE ♂ in Ind Mus

ADULT*—A fairly large species without conspicuous lateral tufts on terminal segments of abdomen, leg-markings and coloration of abdomen distinctive †. Wing 6.5–7 mm

♀—*Head-scales* dull brownish purple, lighter along eye-margins, in certain positions the head appears mainly pale; *palpi* and basal stout part of *proboscis* purple, apical part of latter with greenish lustre, *palpi* reaching to about tip of 5th flagellar segment of *antenna*, latter with rather long sparse hairs. *Mesonotal scales* bright metallic green, those on upper parts of *apn* and *ppn* and *scutellum* coppery or purplish, those on lower parts of lobes and on *pleurae* silvery, with yellow tinge. *Wings* with slight darkening of membrane in region of c-v's 2-3, 3-4, and 4-5, these fairly closely approximated, but more separated than in *M. kempi*. *Legs* purple, with golden and pale markings, femora golden on both sides ventrally and towards base, fore tibia with a large median pale area beneath, mid-pair with wide, nearly complete pale ring in same position, hind pair dark, 1st segment of fore tarsi with a wide basal pale ring, 2 mainly white, but some dark scaling near base, variable in extent, 3, 4, and base of 5 white, basal ½ of segment 1 of mid-tarsi white, also whole of 2, 3, 4, and base of 5, segment 1 of hind tarsi with a subbasal white ring, 4 entirely white (except in a form from Haflong, Assam, in which this segment is entirely dark). Dorsum of *abdomen* deep blue or deep purple, tergite I yellowish at sides, small lateral pale yellow patches on segments IV–VII, narrow basal pale yellow bands on segments II and III, sometimes on II–VI, sternite IV deep purple, remainder pale golden, with a narrow median dark longitudinal line.

♂.—*Palpi* slightly longer than *proboscis*, segment 2 purple at base, with apical pale yellow ring, 3 mainly yellow, with basal and apical dark rings, 4 and 5 purple, with pale yellow scaling at base of 4 beneath, 5 about twice length of 4. Tarsi purple except for pale markings at base of segment 1 on all legs, that on mid-leg usually forming a complete ring. *Hypopygium* (fig. 6, c): 9t with a pair of hairy submedian lobes, more pronounced and pointed in some specimens than in others, lp without obvious teeth.


* Barraud 1929b, p 275

† *M. gravelyi* resembles *M. metallicus* Leac (Malay Penn), but in that species the legs are unbanded in the ♂, and there are no extensive white markings on the legs of the ♀

PUPA* (fig. 5, d) —Differs from that of all other Indian species in the shape of paddle, which is not emarginate on posterior border (in this respect resembling the pupa of *M. magnificus* Leic of Malaya) In abdominal chaetotaxy it agrees closely with pupa of *M. albipes*

LARVA (fig 4, e)* —Fairly easily separated from those of *M. albipes* and *M. minimus* (with which it agrees in having the

Fig 5



Pupal structures of *Megarhinus* (paddles and right halves of certain abdominal segments). a, *M. albipes* Edw.; b, *M. splendens* Wied.; c, *M. kempi* Edw.; d, *M. gravelyns* Edw.; e, *M. edwardsi* Barr.; f, *M. minimus* Theo.

mesothoracic dorso-lateral plate divided into two) by siphon being distinctly longer than chitinised part of anal segment. Length of siphon 1.2-1.5 mm., tuft 4-branched.

HABITAT.—Tree-holes and bamboos.

* Barraud 1931 *α*, p. 1131.

DISTRIBUTION —E HIMALAYAS. Tindharia*, Mungpoo*, and Kurseong*, 3-5,000', x. 1922 (*Barraud*), Darjeeling dist., ASSAM. Nongpoh*, Khasi Hills, vii. 1922 (*Barraud*), Haflong*, Cachar Hills, viii. 1922 (*Barraud*).

Also type-locality* given above.

Not recorded from elsewhere

4 *Megarhinus minimus* Theobald, 1905

Journ Bomb Nat Hist. Soc xvi, p 237 (♂, pl A, fig 1) TYPE-
LOO Yatiyantota, Ceylon, iii. 1902 (*Green*) TYPE ♂ in
Brit Mus

ADULT† —Quite distinct on account of the small size and entirely dark tibiae and tarsi ‡ Wing usually under 5 mm

♀ —*Head-scales* mainly green or bluish-green, those along eye-margins lighter, either blue with violet reflections or nearly white, *palpi* and *proboscis* purple, sometimes with brassy sheen *Mesonotal scales* metallic yellowish-green, some brilliant blue and green scales over wing-roots, *scutellar scales* green, scales of similar colour on upper borders of *ppn* and on *apn*, latter with bluish or purplish lustre when viewed in certain positions, scales on lower parts of *ppn* and on *pleurae* silvery *Legs* purple except for yellowish scaling at bases and on under-sides of femora Dorsum of first few segments of *abdomen* dark greenish-purple, remainder bright purple, indications of pale greenish basal bands on IV-VI or IV-VII, in some specimens tergites II-VII with lateral basal yellow markings, some quite small yellowish lateral tufts at apex of VI, similar tufts of mainly black hairs on VII and VIII, venter yellow, with dark scales on sternites IV, VII, and VIII, no median longitudinal dark stripe In one specimen examined tergites IV-VII are brassy-brown

♂. —Very similar to ♀, but antennae sparsely plumose, palpi a little longer than proboscis, with some yellow scaling at apex of segment 2 and on sides and under-surface of 3 and 4 *Hypopygium* ♂t strongly produced in middle into a shield-like plate bearing hairs laterally, style and appendage together about length of coxite, appendage about $\frac{1}{2}$ length of style; *bl* with one very strong hair about $\frac{1}{2}$ length of coxite and some more slender hairs, *lp* narrow, with a few minute blunt teeth near apex

PUPA § (fig. 5, f) —Fairly easily identified by characters given in key and by its comparatively small size.

† Theobald 1907, p 138, 1910, p 109, Senior-White 1919, p. 189 (♀ desc.), Barraud 1929 b, p. 278.

‡ *M. gigantulus* Dyar & Shannon (Philippine Is) is similar, but the lateral tufts on VII are orange instead of black

§ Barraud 1931 a, p 1131

LARVA (fig. 4, f) † —Comparatively small, siphon 0.7 mm long, tuft of 2 or 3 branches, or occasionally represented by a single strong hair. Diagnostic points given in key.

HABITAT ‡ —Bamboos.

DISTRIBUTION —S INDIA · Yellapur* and Kadra*, N. Kanara dist, vii 1907 (*T. R. Bell*) and ix & x 1921 (*Barraud*), Anamali Hills*, Madras Pres., 3,500' (*Shaffi*). CEYLON: Colombo*, 1913 (*James*), Suduganga, iii 1919 (*Senior-White*), and type-locality mentioned above

Recorded from SUMATRA (*Brug*).

5 *Megarhinus edwardsi* Barraud, 1924

Ind. Journ. Med. Res. xi, p 900 (♂ & ♀). TYPE-LOC Solan, Krol Mt., W Himalayas, 7,000', viii 1923 (*Barraud*) TYPE: ♂ & ♀ in Brit Mus

ADULT † —Very distinct on account of the broad pale yellow border to the mesonotum. Wing about 7 mm.

♀.—*Head-scales* pale green on vertex, pale yellow along eye-margins, stout basal part of *proboscis* dark blue, apical slender part bronzy, *palpi* from $\frac{1}{2}$ – $\frac{1}{3}$ length of proboscis, dark bluish; *mesonotum* with broad pale yellow border from front margin, along sides, over wing-roots to scutellum, disc covered with dark green metallic scales; scutellar scales greenish-blue, scales on *apn* and *ppn* and on *pleura* pale yellow, dorsum of *abdomen* dark metallic blue, tergites I, III, V, and VI with lateral pale yellow patches, III and V with incomplete medial pale bands, VI with pale yellow lateral tufts, VII with similar tufts of mainly golden hairs, VIII with small orange tufts, sternites II–V and VI pale, with a median longitudinal dark stripe, IV and VII nearly all dark; *wings* without definite clouding of membrane, c.-v. 2–3 nearer apex of wing than 3–4 by about 3 times its length. *Legs*. femora dark bluish-green, broadly yellowish beneath and at knee-joints, fore tibia dark green, with yellow scales beneath near apex, mid-tibia yellow, with an apical dark ring, a wide sub-apical yellow ring on hind tibia, segment 1 of fore tarsi dark above at base, yellow beneath, remainder and whole of segment 2 white, 3, 4, and 5 dark bluish, but a few pale scales at base of 3; basal $\frac{1}{2}$ of segment 1 of mid-tarsi yellow, the tip white, 3, 4, and the larger part of 5 white; hind tarsi dark bluish-green, segment 1 with subbasal yellow ring, 2 with wide basal white ring.

♂.—*Palpi* a little longer than proboscis, with pale yellow rings at base of segment 2 and near apex of 3; scutellar

† Barraud 1929 b, p. 276.

‡ Senior-White 1920, p. 319.

scales dark blue; hair-tufts on abdominal segment VII black instead of golden, fore tibia entirely dark, mid-tibia yellow, with basal and apical dark rings, hind tibia with ill-defined subapical yellow ring, fore tarsi with apical white ring to segment 1, 2 all white, basal white ring on 3, 4, and 5 dark blue. *Hypopygium* not examined (the type is the only ♂ known).

PUPA (fig 5, e) †.—Very similar to *M. splendens*, differing only in the length of the larger lateral hair on segment VII

LARVA (fig. 4, a) †—Resembles *M. splendens*, but distinct on characters given in key. Siphon 1.0–1.2 mm. long, tuft usually of 5 branches

HABITAT.—Tree-holes.

DISTRIBUTION.—W. HIMALAYAS, Krol Mt., 7,000', near Solan*, viii. 1923 and vii 1930 (*Barraud*), E HIMALAYAS, Surel*, Darjeeling dist., 5,300', x 1922 (*Barraud*).

Not known from elsewhere

6. *Megarhinus splendens* (Wiedemann), 1819

Culex splendens Wiedemann, Zool. Mag 1, iii, p 2 (♂) TYPE-LOC Java TYPE ♀ in Vienna Mus

Culex regius Thwaites, in Tennent, 1859, 'Ceylon,' p 268 TYPE-LOC Ceylon TYPE non-existent.

Megarhina (species from Sikkim) Giles, 1900, Handbook, 1st ed. p 131 TYPE-LOC. Himalayas, 1,800', vi 1896 (*G C Dudgeon*) TYPE: ♀ in Brit Mus

Megarhina unimisericors Giles (nec Walker), Journ Bomb Nat. Hist Soc xii, 1901, p 604 (May)

Megarhina sikkimensis id., ib. TYPE-LOC Sikkim TYPE: ♀ in Brit Mus. (*G C Dudgeon*) (vide above)

Megarhinus unimisericors Theobald (in part), M C 1, p 225, pl. vii. fig. 28

Megarhinus gilesi Theobald, ib. p 227 (Nov.), pl ix, fig 33 TYPE-LOC. Upper Burma (*Watson*) and Sikkim (*Dudgeon*). TYPE: 3 ♀♀ and 1 ♂ from Burma, 1 ♀ from Sikkim (latter Giles's type of *M. sikkimensis*) in Brit Mus

Worcesteria grata Banks, 1906, Phil. Journ. Sci. 1, pp 780–782 (Sept.), pl 1, figs 1–10 (in part, type-series included specimens of both *M. splendens* Wied. and *M. amboinensis* Doi, latter not an Indian species) TYPE-LOC.: Negros Occidental and Cebu, Philippine Is (*Banks, McGregor*). TYPE number of specimens, both sexes, in Bureau of Science, Manila

Tocorhynchites argenteotarsus Ludlow, 1906, Can Ent. xxxviii, p 367 (type ♀) TYPE-LOC.: Mindanão, Philippine Is. (*Dr. H Newton Kierulff*) TYPE ♀ in US Nat Mus

ADULT (fig 1) †.—This is the largest Indian species, distinguished from *M. edwardsi* by the absence of a wide pale

† Barraud 1931 a, p. 1132

‡ More important recent systematic references Edwards 1913 b, p 222 (syn under *T. unimisericors*), 1917, p 202 (syn under *T. regius*), 1922 a, p 469, and 1929 a, p 4 (syn under *M. splendens*), Dyar & Shannon 1925, p 66 (syn under *M. amboinensis*), Barraud 1929 b, p. 277.

yellow margin to the mesonotum, and from other species by presence of well-developed lateral tufts of yellow, black, and orange hairs on terminal abdominal segments, and by leg-markings and general coloration of body, the dull bronzy scales* on dorsum of mesonotum being characteristic †
Wing 8.5-9 mm


♀—*Head-scales* green and blue with purple reflections, a narrow pale border to eyes, stout basal part of *proboscis* purple or dark blue, *palpi* purple, about as long as first six flagellar segments of *antenna*, latter with rather short, fine sparse hairs. *Mesonotal scales* dull bronzy or greenish-brown, lighter bluish or greenish scales at sides, scutellar scales and those over wing-roots bright bronzy or bluish-green, scales covering *apn* brassy with bluish reflections, or deep blue or green, on *ppn* bluish above, silvery below, on *pleura* silvery. Membrane of *wings* brownish, especially anteriorly, scales at base of wing deep blue or purple, c-vs 3-4 and 4-5 much nearer the base of the wing than 2-3. *Legs*: fore and mid-femora purple, pale golden ventrally and posteriorly, hind pair mainly pale golden, purple dorsally on distal $\frac{1}{2}$, fore tibia purple, with usually some pale golden scaling posteriorly, mid-pair mainly pale golden, purplish at base and apex, hind pair purple, with greenish reflections, tarsal markings variable, segments 3-5 on fore and hind tarsi dark, segment 1 of fore tarsi white, with basal dark ring and sometimes dark tip, a basal white ring on segment 2 variable in width; segments 1-5 on mid-tarsi sometimes entirely white except for a dorsal dark mark at base of 1 and a dark tip to 5, in some specimens the white markings reduced to a subbasal white ring on 1 and a basal white ring on 2 (numerous intermediate variations occur), segment 1 of hind tarsi dark or with yellow or white scaling, forming a more or less complete basal or subbasal ring, 2 often mainly white, but sometimes with a subbasal pale ring only. Specimens

* [In some specimens, especially females, and more so in those from Java, the mesonotal scales appear much less dull, having a rather pronounced greenish-metallic gloss.]

† The most nearly allied species are *M. quasiferon* Loew of Malaya, which differs in having tergite 1 entirely green, instead of dark blue or dark green in middle with pale lateral patches, and *M. rubriflor* Dol. (*immisericors* Walk.), of Dutch East Indies, which differs only in markings on underside of abdomen. The last may only be a variety of *M. splendens* Wied (vide Edwards, Bull. Ent. Res. xiv, p. 356, 1924). There has been a good deal of confusion regarding the synonymy of this species, but that given above appears now to be established, most of the types having been re-examined in recent years. It should be noted that *M. splendens* Wied. as described by Theobald is now regarded as a synonym of *M. aurifluus* Edw., 1921, from Formosa.

from the eastern parts of India usually show more extensive white markings than those from the west. Dorsum of abdomen dark blue or dark green, tergites I-III and V with lateral pale yellow patches visible dorsally, IV and VI with

Fig 6



♂ hypopygial structures of *Megarhinus* a, *M. splendens* Wied.
b, *M. kempi* Edw c, *M. graveleyi* Edw (with variation in shape
of style) Lettering as on p. 4

smaller similar markings; lateral outstanding tufts of hairs on VI-VIII, those on VI yellow and black, on VII black, on VIII orange; sternites II, III, V, and VI pale yellow with median purple line, IV, VII, and VIII mainly dark purple.

♂—*Palpi* about length of proboscis, purple, with yellow or white scaling at apex of segment 2 and near apex of 3, variable in extent, *antennæ* densely plumose, 1st flagellar segment with tuft of scales. *Tarsi*. segments 3-5 on all legs dark, fore tarsi entirely dark or with pale scaling at bases of segments 1 and 2 beneath, mid-tarsi with pale scaling near base of I sometimes forming a wide ring, II with basal white ring very variable in width, segment I of hind tarsi dark or with pale scaling at base, 2 with wide basal or sub-basal white ring. *Hypopygium* (fig 6, a) 9t wide and only very slightly emarginate on apical border, *lp* without marked serrations.

PUPA (fig 5, b)*—Diagnostic points given in key, of the five submedian and sublateral hairs one is long and black on tergites 2, 6, and 7, and two on tergites 3, 4, and 5, larger lateral hair on segments 2-7 long.

LARVA (figs 2, 3, 4, d) † (*vide key*)—Length when fully grown from 15-16 mm, colour crimson or deep reddish-brown, length of siphon-tube 1.1-1.5 mm, tuft with 4-8 branches, ventro-lateral plate on I sometimes divided into two, head and chitinous parts of body in larval skin deep rich brown.

HABITAT ‡.—Tree-holes, bamboos, and sometimes in domestic collections of water such as water-butts, jars, etc.

EGG §.—Green states that the eggs are scattered singly and separately on the surface of water and do not tend to run together in strings. The egg is of a regular oval form, 0.55 mm. long by 0.37 mm. broad, of a creamy-white colour; surface closely studded with spinous granules, some of which are larger than the rest and disposed at more or less regular intervals. Each of these larger granules has a prominent apical point. This granular formation doubtless accounts for the buoyant manner in which it floats, the whole contour of the egg being visible above the surface-film. When the larva hatches, the egg divides transversely across the equator, the empty halves float on the surface, with the convexity upwards.

DISTRIBUTION.—Common along the HIMALAYAN foothills from Dehra Dun (U.P.) to the north-east boundaries of Assam, through ASSAM and BURMA, to the ANDAMANS, in BENGAL, BIHAR, Orissa down to the Madras boundary; on the west coast from Savantvadi State to TRAVANCORE and Ceylon.

* Barraud 1931 a, p. 1132; Edwards 1926 a, p. 115.

† Barraud 1931 a, p. 1131; Theobald 1903 a, p. 118, Senior-White 1927, p. 66.

‡ Green 1905, p. 160; Paiva 1910, p. 187; Senior-White 1920, p. 319.

§ Green 1905, p. 160.

It has not, so far, been found in Central India nor west of a line drawn from Simla to Ratnagiri (on the coast south of Bombay)

Outside India its range extends through the MALAY PENINSULA and islands, SIAM, COCHIN CHINA, PHILIPPINES, and HONG KONG.

SEASONAL PREVALENCE.—Throughout the greater part of the year in South India and Ceylon, and as far north as Calcutta, further north and in the north-east it is most abundant during the monsoon.

Tribe CULICINI.


This tribe is represented in India by fifteen genera, several of which are divided into subgenera, according to the classification adopted in this work. The number of species at present known is 239, and rather more than half these are included in the larger genera *Aedes* and *Culex*. The characters now used for the separation of genera, and applicable to both sexes of the adult, are often rather obscure, and are to be found chiefly in the pleural chaetotaxy and in the venation of the wing. Other characters, such as scaling and length of palpi, have also been taken into consideration, but these are used much less than formerly, as it has been found that a good deal of variation occurs sometimes in these respects in closely allied groups of species. The presence or absence of pulvilli and of hairs on the squama of the wing are also of importance in distinguishing certain genera. The structure of larvæ and pupæ, which often shows more definite generic characters than the adults, has also been applied.

Subgeneric divisions are more often based on characters present only in the ♂ sex (structure of hypopygium, length of palpi, etc.). In the larger genera *Aedes* and *Culex* the larvæ do not show well-marked subgeneric characters, and such characters are sometimes not very definite in the adult ♀♀.

For general remarks on classification and grouping of genera of the world the reader is referred to Edwards (1932), pp 63-65.

CHARACTERS ADULT.—Mosquitoes variable in size, but not as large as the majority of Megarhine species, and including some very small forms. Usually less highly ornamented with coloured scales than the Megarhinini, though silvery metallic, blue, green, or violet scales are present in some of the smaller species. Clypeus longer than broad, rounded above and in front. Proboscis flexible and nearly always uniformly slender and not hooked, but sometimes swollen at tip. Palpi of ♀♀ never as long as proboscis, those of ♂♂ of very variable length. Scutellum more or less trilobed and with three

Fig. 7.



Tribe Culicini: generic characters of adults.

a, side view of thorax, showing chaetotaxy, genus *Theobaldia*.

h, ditto, genus *Aedes*.

apn, anterior pronotal lobe and bristles; ppn, posterior pronotal lobe and bristles; pp, propleural bristles; sp, spiracular bristles; psp, postspiracular bristles; stp, sternopleural bristles; pa, prealar bristles; u.m.e., upper mesepimeral; l.m.e., lower mesepimeral.

b and c, tip of abdomen of ♀ *Culex* and *Aedes* respectively: 6, 7, 8, abdominal tergites VI, VII, and VIII; c, cerci.

d, e, and f, tip of last tarsal segment of *Theobaldia*, *Aedes*, and *Culex* respectively: e, empodium; p, pulvilli; c, claw.

g, base of wing from beneath, genus *Theobaldia*: s, squama, with fringe of hairs; a, alula (also fringed); sc, base of subcostal vein bearing hairs.

2687

591.954

separate groups of bristles. Hind margin of wing without a V-shaped thickening. Abdomen with tergites and sternites covered with scales, which usually lie flat on the surface. Hypopygium of ♂ usually with well-developed paraprocts (chitinisations of anal segment), phallosome without leaflets, but often with chitinised lateral plates bearing teeth.

PUPA —Paddle usually with one terminal hair (two in *Culex*, but none in *Tripteroides*, *Topomyia*, *Harpagomyia*, most *Ficalbia* and *Mansonia*). Outer part of paddle not usually produced beyond termination of midrib, but, should it have this appearance, the hind margin is either serrated, or with terminal hair, or the respiratory trumpet ends in a chitinised spine, or is divided nearly to the base into leaflets.

LARVA —Mouth-brushes composed of hairs, of which the median series have, in many species, serrations on one side apically, not usually prehensile, except in *Aedes* (*Mucidus*) and *Culex* (*Lutzia*), in which the hairs are modified into somewhat hooked lamellæ, 30 or more in number in each brush (in *Megarhinus* the brush consists of ten lamellæ arranged in a line). Clypeus with six pairs of hairs, in addition to a pair of preclypeal spines. The character and relative positions of hairs *A*, *B*, *C*, *d*, and *e* are of importance. The sixth pair is nearly always very small and lies on the front of the clypeus slightly external and dorsal to the preclypeal spines. Chatotaxy of thorax and abdomen variable*. In some forms one or more pairs of strong spines are present on the thorax. Segment VIII of abdomen with comb of teeth on each side (absent in *Megarhinini*), and a siphon-tube arising from dorsum, having respiratory openings and movable valves at its tip, and one or more pairs of hairs or hair-tufts along its length (occasionally these are very small). Siphon with or without pecten. Anal segment with either a chitinised saddle or completely enclosed in a ring of chitinisation, two pairs of subdorsal hairs, one pair of lateral hairs, two pairs of papillæ surrounding the anal opening, and a ventral fan of hairs (occasionally reduced to one pair).

* [The thoracic and abdominal chatotaxy has not yet been closely studied in the various genera of Culicini, it is practically certain that in all genera of this tribe the same set of hairs exists as in *Megarhinini* and also in *Anophelemi*, but the development of the individual hairs varies very greatly in different genera and species. The thoracic chatotaxy of *Tripteroides*, *Aedes*, and *Culex* is illustrated in figs 9, 39, and 102 respectively. Among points of distinction between Culicini and *Megarhinini* it may be noted that hair 8 is apparently never associated with the pleural hairs (9-12) in Culicini, as it is in *Megarhinini*, hairs 5-7 of prothorax and 4-6 of mesothorax are not included in one chitinous plate, hair 1 of mesothorax is never duplicated, and hair 14 is present.]

Egg—The eggs of Indian *Culex* have been very little studied. They are longer than wide, with hard shell, without lateral floats, and more or less fusiform in shape, not roundly oval or spiny, as in the *Megarhinini*. Those of many species are able to resist desiccation over long periods. They may be laid singly, or cemented together in the form of rafts

KEYS TO GENERA

Adults

1. Margin of squama fringed (fig. 7, g) (fringe usually complete, but interrupted in some small species), vein 6 ending well beyond level of fork of vein 5 (figs 21, 32, 34) 2
- Margin of squama quite bare (fig. 13, b), vein 6 short, usually ending at about level of fork of vein 5 (figs 13, 15) 15.
2. Pulvilli present (fig. 7, f); bucco-pharyngeal armature present in ♀ (fig. 80, a-g) (spiracular and postspiracular bristles absent) *CULEX*, p. 332.
- Pulvilli absent or rudimentary (fig. 7, d, e) bucco-pharyngeal armature absent in ♀ (fig. 80, h-k) 3.
3. Postspiracular bristles absent (fig. 7, a) 4
- Postspiracular bristles present (fig. 7, h) 13
4. Spiracular bristles present (from 3-10) (fig. 7, a) 5
- Spiracular bristles absent (fig. 7, h) 6.
5. Fairly numerous hairs on underside of wing at base of vein sc (fig. 7, g), large mosquitoes, palpi of ♂ long, the tips spatulate *THEOBALDIA*, p. 86.
- Base of vein sc without such hairs; small mosquitoes; palpi of ♂ quite short, like those of ♀ *TRIPTEROIDES*, p. 34.
6. *apn* lobes enlarged and approximated behind head, mesonotum without dorso-central or prescutellar bristles 7.
- apn* lobes of normal size and well separated; dorso-central and prescutellar bristles well developed 8
7. Postnotum bare (♂♂ of Indian species unknown) *HEMAGOGUS*, p. 300.
- Postnotum with group of small hairs, antenna of ♂ with last two segments very long *HEIZMANNIA*, p. 299.
8. Postspiracular area with scales; tarsal claws of ♀ usually toothed (palpi of ♀ 4, or more, length of proboscis) [TERRIA), p. 320.
- Postspiracular area bare; claws of ♀ simple *ARMIGERUS* (LEUCES- 9.
9. All segments of antennae of ♀, and last two of antennae of ♂, short and thick (fig. 32); femur of middle leg with tuft of scales at tip *ANDOMYIA*, p. 181.
- Antennae normal, slender; femur of middle leg without scale-tuft 10.

- 10 First segment of fore tarsi longer than last four together, 4th very short in both sexes, shorter than 5th (fig 21, b), wings spotted as in many species of *Anopheles* (fig 21, a) [p 94.
ORTHOPODOMYIA,
- First segment of fore tarsi not longer than last four together, 4th not shortened in ♀, wings not spotted, but may be speckled 11
- 11 Fork-cells of wing short, anterior being only about $\frac{1}{2}$ length of stem, palpi of ♂ clubbed at apex [(MIMOMYIA), p. 105
FIOALBIA
- Fork-cells of wing not unusually short, anterior being about length of stem, palpi of ♂ not clubbed at apex 12
- 12 Small mosquitoes with comparatively large wing-scales, proboscis of ♂ greatly widened towards apex (fig 27, h), that of ♀ slightly enlarged [p 105.
FIOALBIA (part),
- Larger yellow or yellowish-brown mosquitoes, wing-scales normal, proboscis not enlarged [LETTIDIA), p 119
MANSONIA (COQUIL-
13. Wing-scales unusually broad, and many asymmetrical, tergite VIII of ♀ with a row or patch of short tooth-like spines (fig 30), apical segment of palpi of ♂ minute, claws of ♀ simple [OIDES), p 123.
MANSONIA (MANSONI-
- Wing-scales not unusually broad and not asymmetrical, tergite VIII of ♀ without spines, apical segment of palpi of ♂ usually of moderate length, claws of ♀ usually toothed 14
14. Proboscis rather stout, somewhat laterally compressed, and turned downwards towards tip (fairly large dark species, with flat scales on vertex and scutellum) Proboscis fairly slender and straight, ornamentation and scaling very various. AÆDES, p. 134.
15. Wing-membrane with very small microtrichia requiring a magnification of more than 80 for their detection, anterior fork-cell usually very short (fig 15) URANOTÆNIA, p 56
- Wing-membrane with microtrichia, as usual, visible under a magnification of 50 or less, anterior fork-cell not usually very short 16.
- 16 Proboscis very hairy and much enlarged at tip (fig 12, b) HARPAGOMYIA, p. 47.
- Proboscis not hairy and not enlarged 17.
- 17 Wing-scales emarginate at tips (fig 13, c). HODGESIA, p. 52.
- Wing-scales normal TOPOMYIA, p. 45.

Larvæ (4th stage)*

- 1 Ventral fan or brush of anal segment represented by a single pair of hairs (fig. 12) 2.

* Larvæ of Indian species of *Hæmagogus* and *Topomyia* are unknown.

- Ventral fan of anal segment of more than 2 separate hairs 3.
- 2 Metathorax with a long spine on each side (fig 9) *TRITETEROIDES*.
HARPAGOMYIA,
[*TOPOMYIA*.]
- Metathorax without long spines
3. Antenna very large, somewhat flattened; siphon hairy all over, and with a pair of very long subposterior branched hairs, the branches longer than siphon (fig 33) *ABDOMYIA*
- Antenna not very large and not flattened, siphon not usually hairy all over, siphonal hair-tufts with branches of only moderate length or quite small 4.
- 4 Siphon with several subposterior hair-tufts usually arranged in pairs *CULEX*.
- Siphon with not more than one pair of subposterior hair-tufts 5
- 5 Siphonal valves modified for piercing tissues of aquatic plants (figs 31, a) *MANSONIA*.
- Siphonal valves not modified for piercing (but compare *Ficalbia hybrida*, p. 112) 7.
7. Hair-tufts on siphon very near base of tube (figs 14, 20) 8
- Hair-tufts on siphon near middle of tube or towards apex 9.
- 8 Antennal tuft not much beyond middle; size large *THEOBALDIA*.
- Antennal tuft at $\frac{1}{2}$ from base, size small *HODGESIA*.
9. Siphon without pecten 10.
- Siphon with pecten 12.
- 10 Abdominal segment VIII with lateral chitinated plates more or less developed (sometimes also on VII); comb of alternating large and small teeth in a row *ORTHOPODOMYIA*.
- Abdominal segments VII and VIII without lateral chitinated plates; comb-teeth all about same size 11.
11. Antenna comparatively short, with smooth shaft and with a small hair representing antennal tuft (fig. 75) *ARMIGERES*
- Antennae fairly long, with small spicules on shaft and a large branched hair-tuft (figs. 25-27) *FICALBIA* (part).
12. Segment VIII with lateral chitinated plates (*vide* figs. 16, 54) 13.
- Segment VIII without lateral chitinated plates 14.
13. Pecten-teeth usually with fringe, or lateral denticles on both sides from base to apex (fig 16). *URANOCEPHALA*.
- Pecten-teeth usually with or without a few lateral denticles near base on one side only (fig. 54) [*STEGOMYIA*].
14. Pecten-teeth few (1-4) and simple (without lateral denticles); both *osc* and *psc* with a number of branches *AÆDES* (some species of *FICALBIA* (part)).
- Pecten-teeth usually fairly numerous, and majority with lateral denticles; *osc* usually single and long *AÆDES* (most species);
HYMOMANIA;
(?) *HEMAGOGUS*.

Genus **TRIPTEROIDES** Giles, 1904

Journ. Trop. Med vii, p 368. Genotype, *Runchomyia phalippinensis* Giles.

Rachnotomyia Theobald, 1905, Journ. Bomb. Nat. Hist. Soc. xvi, p 248. Genotype, *R. ceylonensis* Theo.

Colonomysia Leicester, 1908, Cul. Malaya, p 233. Genotype, *C. caeruleocephala* Leic.

Skeuromysia id., ib. p 248. Genotype, *S. fusca* Leic.

Squamomyia Theobald, 1910, Rec. Ind. Mus. iv, p 28. Genotype, *S. inornata* Theo.

Tricholeptomysia Dyar & Shannon, 1926, Insec. Insc. Mons. xiii, p 72. Genotype, *Wyeomyia nepenthicola* Banks.

Until very recently this genus has been known as *Rachnotomyia*, but Edwards (1932) has adopted the present name, which was suggested, but not adopted, by Giles in 1904, and has given an up-to-date definition of the genus. It is now divided into four subgenera, of which only one, *Tripteroides* (s. str.), is represented in India. The characters of the adults, larvae, and pupae, common to the Indian species, are given below. The eggs of these are at present unknown.


ADULT* (fig. 8).—Small mosquitoes with slender legs, proboscis usually long and slender, sometimes as long as whole body, palpi quite short in both sexes, usually about $\frac{1}{2}$ length of proboscis or less (*T. edwardsi* about $\frac{1}{4}$); antennae of ♂ moderately plumose, with last two segments elongate, those of ♀ slender, with segments of flagellum of about equal length, hairs fairly long; eyes touching for some distance above antennae; head mainly flat-scaled, some upright scales on nape, one pair of strong occipital bristles separated from small lower orbital bristles; thoracic bristles reduced, 1 posterior pronotal, 3–6 spiracular, no postspiracular, no lower mesepimeral, usually none along upper border of sternopleura, some large patches of flat scales on pleura; *apn* fairly large but well separated, flat-scaled, mesonotal scales variable in form in different species, either narrow and almost hair-like or distinctly broad, bristles few, dorso-central series sometimes absent, giving the thorax a smooth appearance; scutellar scales flat, postnotum usually with a pair of slight furrows and without setae (except in *T. edwardsi*). Fork-cells of wing moderately long, vein 6 terminating nearer tip of wing than commencement of 2 and fork of 5, squama with some hairs. Legs slender, hind tibia shorter than others, segment 1 of hind tarsi usually longer than tibia, fore tarsal claws of ♂ unequal, the larger with a blunt tooth (except in *T. powelli* var. *indica*), claws of ♀ simple. Abdomen of ♀ with hairs and bristles at tip, but few hairs on other segments; hypopygium of ♂ (fig. 10) rather prominent, coxite comparatively small, with hairy lobe on anal surface and long terminal style.

* Edwards 1921 a, p 283; 1924, p. 359; 1932, p 73; Barraud 1929 a, p. 1058; Barraud & Covell 1928, p. 676 (buccal cavity).

carrying a small, stout appendage at its tip, 9t produced into a pair of well-developed submedian lobes carrying strong bristles, which may be flattened, resembling leaflets in some species, paraproct with a few strong teeth at crown and a few minute hairs, phallosome resembling an incomplete tube, feebly chitinated, sometimes with a few minute teeth at apex

PUPA*—Paddles small and bluntly pointed, without terminal hair or fringe, but some very small hairs on posterior border near termination of midrib, conspicuous lateral branched hairs on outer posterior corners of VII and VIII, those on VIII extending beyond tip of paddle, long single hairs on IV-VI or IV-VII.

Fig 8



Adult characters of *Tripteroides*: a, *T. similis*, ♀, side view, legs and wings omitted; b, *T. aranoioides*, ♂ head; c, *T. similis*, mid-femur, front view; d, diagram showing posterior frontal and spiracular bristles


LARVA: 4th stage† (fig. 9)—Those of the Indian species, so far as they are known, have the following characters:—Head small, antenna quite short and comparatively stout; a double dorso-lateral black spine on each side of metathorax [representing hair 7], a pair of smaller, single spines on mesothorax [also representing hair 7]; hair 5 also thickened in some species; propleural group of hairs (9-12) latero-ventrally placed, so that the tubercle is almost or quite visible from

* de Meijere 1910, p. 280; Brug 1931 p. 20; Edwards 1932, p. 74.

† Edwards 1921 c, p. 283; 1926 b, p. 393; 1932 (see also under *aranoioides*).

above, almost as well-developed as mesopleural and metapleural groups, hair 12 of metapleural group small but distinct, hair 6 of mesothorax on plate with 7, on pro- and metathorax separate, hair 8 of mesothorax not very large], numerous stellate hairs on thorax and abdomen [those on dorsum of thorax representing hairs 0 and 1 of prothorax, 1 of mesothorax, and 1 and 3 of metathorax], a comb of rather

Fig 9



Larval characters of *Tripteroides*: a, *T. vicina* Edw. (a Malayan species), whole larva showing general appearance; b-g, *T. arancoides*, details b, mentum, c, comb; d, preclypeal spines, e, antenna, f, spine of metathorax, g, chaetotaxy of dorsum of thorax [numbers as employed by Puri in *Anopheles*, the small plate bearing hairs 1-3 of prothorax is shown too large, hairs 9-12 are the pleural hairs of pro-, meso-, and metathorax borne on strong tubercles, hair 7 on meso- and metathorax is transformed into a spine; hair 13 (not shown in figure) is a large stellate hair, ventral in position and far removed from the pleural tubercle.]

large teeth on either side of segment VIII set in a single but sometimes irregular row, siphon-tube of moderate length, tapering towards tip, some fairly long hairs along posterior border of tube from base to apex arranged more or less in a double row, either 2-branched or single, the pair nearest the base usually divided into several branches, similar single or double hairs irregularly arranged over the lateral and anterior surfaces, pecten consisting of a row of fine bristle-like teeth commencing some distance from base and continued nearly to apex, a chitinous saddle on anal segment enclosing dorsum and sides, some strong spines on posterior border, *osc* single, *asc* branched, lateral hair very long and either single or 2-branched, anal fan absent, being represented by one pair of branched hairs, fully-grown larva about 7 mm long

BIONOMICS* —Little is known of the habits of these species in India. The adults are most probably diurnal, but never appear to be abundant or troublesome. Senior-White states that ♀♀ of *T. aranoides* do not bite, that he has seen adults at flowers of *Mikania scandens* at dusk in February on the Suduganga Estate in Ceylon, and by day on the windows of a bungalow. Most of the species are found in forest or jungle during the rains, but *T. aranoides*, which is by far the commonest, may breed near human habitations where suitable conditions exist. They are mainly bamboo-breeders, but the larvæ of *T. aranoides* have been found in water in a hollow *Erythrina*-tree (Senior-White), and also very commonly in pitcher-plants in Malaya; those of *T. affinis* occurred in a tree-hole at Belgaum (Barraud). The larvæ of *T. aranoides*, and probably those of other species, can be fairly easily separated from those of other bamboo-breeders by the pale creamy colour of the body, against which the numerous black stellate hairs can be clearly seen.

DISTRIBUTION —The subgenus *Tripterooides* occurs in the south-western, southern, and eastern parts of the Indian area, including Ceylon. Further east it extends through Malaya, Cochin China, Japan, Australian region, Philippines, and Fiji. Five species have been found in India up to the present.

Key to Adults.

- | | | |
|--|----|--|
| 1. Femora spotted | 2 | |
| Femora unspotted | 3. | |
| 2. Anterior pronotal lobes covered with flat silvery scales | | <i>similia</i> , p. 38. |
| Anterior pronotal lobes with a row or two of small flat brown scales | | [p. 39. <i>powells</i> var. <i>indicus</i> , |

* de Meijere 1910, p. 928, Senior-White 1920, p. 320, Buxton & Hopkins 1927, p. 76; Dover 1928, p. 28.

3. Mesonotum with a continuous white-scaled margin from front, along sides, over wing-roots to scutellum, postnotum with well-marked tuft of setæ *edwardsi*, p. 41.
 Mesonotum without a white margin, postnotum bare 4.
 4 White scales on *ppn* and pleuræ, a narrow blue border to eyes 5
 Brown scales on *ppn*, silvery scales on pleuræ, a wide deep blue border to eyes *affinis*, p. 44.
 5 Light scales forming a lateral straight border to tergites for length of abdomen. *aranoides*, p. 42.
 Light scales forming lateral triangular patches, or a serrated border, to tergites for length of abdomen [p. 43.
aranoides var. *serrata*,

Key to known Larvæ.

- Comb of about 20 teeth, subequal in size, and set in a close row *affinis*.
 Comb of 10 or 12 teeth, of which about 5 are larger than remainder, a few towards dorsum widely spaced, ventral hairs of anal segment branched *aranoides*.
 Comb of 5 large teeth; ventral hairs of anal segment single *dofleini*, p. 45.

Key to Pupæ.

- Sublateral hairs on posterior border of tergite VII all comparatively small and extending only slightly beyond posterior margin of that segment, ends of branches of larger lateral hair on VII not extending to outer posterior corner of tergite VIII *aranoides*.
 One or two sublateral hairs on posterior margin of tergite VII comparatively long and reaching as far as posterior margin of VIII, ends of branches of larger lateral hair on VII extending to, or beyond, outer posterior corner of VIII *affinis*.

7 *Tripteroides similis* (Leicester), 1908

Odonemyia similis Leicester, Cul Malaya, p. 235 (♂ & ♀). TYPE-LOC: Bukit Kutu, Malay Penin. (Leicester). TYPE: ♂ and co-type ♀♀ in Brit Mus

ADULT*—Distinguished by silvery scales on *apn*, each femur marked with two silvery spots, no dorso-central bristles on mesonotum, the scales on this part narrow, metallic scales on scutellum, and wing-scales fairly broad and rather dense†. Wing 3.3–3.6 mm.

* Barraud 1929 a, p. 1080 (*Rachnonotomyia*).

† [*T. aenea* Edw. (Malay) and *T. plumosa* Brug. (Sumatra) are similar, but have brown scales on *apn*, the latter has a tuft of scales at end of ♂ style.]

♀.—*Head* a patch of deep blue scales on vertex, wider in front than behind, remainder of head clothed with brownish-black scales except for a patch of white scales laterally, some dark upright scales on nape, antennæ with pale pubescence on shaft, hairs dark brown, torus brown, clypeus, palpi, and proboscis dark brown, palpi about $\frac{1}{2}$ length of proboscis, latter very long, as long as whole body. *Thorax* mesonotal scales narrow and many hair-like, greenish-brown or yellowish-brown, no dorso-central bristles, flat bronzy-green scales on scutellum, flat silvery scales on *apm*, a large black area in middle of pleuræ surrounded by brown, mainly covered with patches of silvery scales. *Wings* fairly densely clothed with rather broad scales. *Legs* tibiae and tarsi dark brown, unbanded, all femora marked with two silvery spots on anterior surface, one near apex and one near middle, the medial one on hind femur not always very distinct, being sometimes confluent with a silvery marking running from base. *Abdomen* dark purplish or bronzy, with large lateral silvery patches extending on to dorsum to form narrow apical bands on II-V or II-VI.

♂.—Very similar to ♀, antennæ moderately plumose, palpi only about $\frac{1}{2}$ length of proboscis. *Hypopygium* (fig 10, c) lobes of 9t wide and armed with a number of flattened spines or leaflets, style enlarged on apical $\frac{1}{2}$ and comparatively short.

PUPA, LARVA, and EGG—Unknown

HABITAT.—Bamboos (*Barraud*)

DISTRIBUTION.—In India recorded only from DARJEELING DIST., Surel* and Mungpoo*, c 5,000', x 1922 (*Barraud*), Sukna* and Marianbarrie Estate*, 500', viii 1928 (*Sobha Ram*)

Known also from the MALAY PENINSULA and ARCHIPELAGO.

8 *Tripteroides powelli* (Ludlow), 1909, var. *indicus* (*Barraud*), 1929.

Uranotenia powelli Ludlow, Can. Ent. xii, p. 235 (♀). TYPE-LOC.: Camp Wilhelm, Yababas, P.I., 3. 1. 1909 (*W. A. Powell*). TYPE: ♀ in U.S. Nat. Mus.

Rachionotomyia powelli (Ludl.) var. *indica* Barraud, Ind. Journ. Med. Res. xvi, p. 1061 (♂ & ♀, ♂ hypop., figs.). TYPE-LOC.: Sukna, Darjeeling dist., 500' (*Sobha Ram*), viii. 1928. TYPE: ♂ & ♀ in M.S.I. coll., Kasauli.


ADULT †.—The following characters taken together distinguish this species from its nearest allies in India and

† Theobald 1910 b, p. 519 (*Uranotenia*); Dyar & Shannon 1925, p. 72 (*Rachionotomyia*).

Malaysia*.—Coloration of mesonotal integument, presence of dark brown scales on *apn*, all femora marked with two silvery spots, larger tarsal claw of fore leg of ♂ simple, presence of several pairs of dorso-central bristles on mesonotum, wing-scales not very dense and fairly narrow. Wing about 2.5 mm.

♀.—*Head* a broad area of flat metallic deep blue scales on anterior $\frac{1}{2}$ of dorsal surface, extending to sides, the straight posterior margin of this area bordered with deep violet or brownish-black scales, a patch of silvery scales low down at each side, clypeus, palpi, proboscis, and antennal hairs dark brown, proboscis slender, about length of whole

Fig 10



Hypopygial details of *Tripteroides* a, *T. powelli* var *indica*, coxite and style from inside, half of tergite; b, *T. araneoides*, whole hypopygium, side view, c, *T. similis*, half of tergite, style

body, palpi about $\frac{1}{2}$ length of proboscis. *Thorax*: integument of prescutum and scutellum light brown, of scutum and postnotum dark brown, several pairs of dorso-central bristles, mesonotal scales narrow and yellowish, with a few darker ones.

* [Allied Oriental species are: (1) *monetifera* Dyar (Philippines), *apn* with narrow golden scales, (2) *hybrida* Leic. (Malaya) and (3) *nin-dwenter* Giles (Philippines), *apn* with flat blue scales, (4) *bambusa* Yam. (Japan) and (5) *vicina* Edw (Malaysia), thoracic integument blackish, only narrowly pale on front margins, *bambusa* having a slight notch, *vicina* a distinct tooth on larger front claw of ♂ (not *vice versa*, as stated in Edwards's key, 1922 *l.*, p. 444), (6) *caeruleocephala* Leic and (7) *proxima* Edw (Malaya), thoracic integument all yellowish, these two also differing in front claws of ♂]

intermixed, scutellar scales rather small, flat, and dark brown, integument of pleuræ of mesothorax black, an oval dark area around anterior spiracle, pleuræ of prothorax and metathorax, and coxæ pale yellowish, a large area of brilliant silvery scales covering larger part of sternopleura and mesepimeron, a row or two of flat dark brown scales on *apn*, a few dark hair-like scales on *ppn*. *Wing-scales* not very dense, and mainly rather narrow. *Legs* markings as in *T. similis*. *Abdomen* tergites brownish-black, with yellowish or bronzy sheen, large lateral silvery patches not usually forming complete bands on dorsum, venter pale yellowish.

♂—Antennæ moderately plumose, wing about 2.3 mm., larger claw of fore tarsi simple. *Hypopygium* (fig 10, a) lobes of 9t longer and narrower than in *T. similis*, each armed with about six flattened bristles or leaflets.

LARVA, PUPA, and EGG—Unknown.

HABITAT—Bamboos (*Barraud*).

DISTRIBUTION of var *indica*.—ASSAM Golaghat*, Subsagar dist. (*Christophers*), Nongpoh*, Khasi Hills, vii. 1922 (*Barraud*). DARJEELING DIST, Sukna*, 500', x. 1922 (*Barraud*) and viii. 1928 (*Sobha Ram*). UPPER BURMA. Maymyo*, viii. 1930 (*R. C. Wats*).

The type-form is known only from the Philippine Islands †.

9 *Tripteroides edwardsi* (*Barraud*), 1929

Heizmannia edwardsi Barraud, Ind. Journ. Mod. Res. xvii, p. 264 (♀). TYPE-LOC. Yellapur, N. Kanara dist., x. 1921 (*Barraud*).

TYPE: one co-type ♀ in Brit. Mus., one co-type ♀ in M. S. I. coll., Kasauli.

ADULT—Distinct on account of the border of white scales to mesonotum and presence of setæ on postnotum. Wing-length 3.8–4 mm.

♀—*Head* a wide area of flat grey scales on vertex, a narrow pale border to eyes, and a large pale patch at each side of head, a few dark upright scales on nape, torn with some pale scales on inner sides, flagellar segments and hairs dark, clypeus pale grey, palpi and proboscis black with dark bluish sheen, palpi about $\frac{1}{4}$ length of proboscis, latter of only moderate length. *Thorax*: mesonotal scales broad, very dark brownish-grey on disc, a border of white scales along sides, over wing-roots to scutellum, scales over wing-roots large, no dorso-central bristles, scutellar scales broad and flat, white on lateral

† No specimens of the type-form are available for comparison, but according to the description it differs from the variety in having the mesonotal scales dark brown. It should be noted that several other closely related species occur in Malaysia.

lobes, mostly dark on mid-lobe, a well-marked tuft of setæ on postnotum, dull white scales on *apn*, flat white scales (not silvery) almost entirely covering pleuræ, including *ppn*, postspiracular area, sternopleura, and mesepimeron. *Legs* tibiae and tarsi brownish-black, unbanded, fore and mid-femora dark on anterior surface, mostly pale posteriorly, hind femur white on both sides, except along dorsal edge, knee dark. *Abdomen* deep brownish-black with large irregular lateral white patches, widest towards apex of each segment, not forming bands over dorsum.

♂ and early stages unknown.

HABITAT —Bamboos (*Barraud*)

DISTRIBUTION —Known only from type-locality.

10 *Tripteroides aranoides* (Theobald), 1901

Wyeomyia aranoides Theobald, M C n, p 274 (♀) TYPE-LOC. Taiping, Perak, Malay Penin (*Wray*), 22 xi and 21 xii. 1899
TYPE: ♀ in Brit Mus.

Rachionotomomyia ceylonensis Theobald, 1905, Journ Bomb Nat Hist. Soc. xvi, p 248 (♀) TYPE-LOC: Peradeniya, Ceylon (*Green*), x 1901 TYPE ♀ in Brit. Mus.

Skewromyia fusca Leicester, 1908, Cul Malaya, p 248 (♂ & ♀). TYPE-LOC Malay Penin (*Leicester*) TYPE ♂ & ♀ in Brit. Mus. (from larva in bamboos).

Squamomyia mornata Theobald, 1910, Rec. Ind Mus iv, p 28 (♂). TYPE-LOC west slopes of Dawna Hills, Lower Burma, 2-3,000' (*Annandale*), 2 or 3 in 1908 TYPE ♂ in Ind Mus.

Ficalbia tenax de Meijere, 1910, Ann Jard Bot Butenzorg (2), Supp iii, p 828 (♂ & ♀, adult, larva, and pupa) TYPE-LOC Tjibodas, Java TYPE Amsterdam Mus.

var *serrata* Barraud (of *R. aranoides*), 1929, Ind Journ Med Res xvi, p. 1059 TYPE-LOC Nagargali, Belgaum dist, viii. 1921 (*Barraud*). TYPE ♂ & ♀ in M S I coll.

ADULT* —The only other Indian species with which this could be confused is *T. affinis*, but the two are distinct on characters given in key† Wing 3 2-3 4 mm

♀ —*Head* mainly covered with large dark brown flat scales, a narrow border of blue scales along eye-margins, palpi and proboscis dark brown, with sometimes a bronzy sheen, palpi about $\frac{1}{2}$ length of proboscis, latter long and slender, longer than abdomen, clypeus partly covered with white scales. *Mesonotum* covered with broad dark greyish-brown scales, no dorso-central bristles, scutellar scales flat and dark greyish-brown, white scales on *apn* and *ppn*, similar white scales (not silvery) almost covering pleuræ, integument brown or yellowish. *Legs* dark brown, unbanded, undersides of femora

* Theobald 1907, p. 518, and 1910 b, p 529 (*Squamomyia*); Edwards 1913 b, p 241, and 1922 d, p. 462 (syn.); Barraud 1929 a, p. 1059.

† [Another similar Oriental species is *nepenthes* Edw. (Malaya), with shorter proboscis, only one claw on hind tarsi, and clypeus bare.]

lighter Dorsum of *abdomen* brownish-black with dark bluish sheen, white scales on venter appearing bluish in some positions, whitish scales on sides of tergites*

♂—Resembles ♀ except for more plumose antennae *Hypopygium* (fig 10, b): 3-4 strong spines on lobes of 9t, as in *T. affinis*

PUPA †—Points of difference from *T. affinis* given in key

LARVA ‡—Peclypeal spines strong, curved, and pointed, arising from slight prominences of preclypeus *Frontal hairs* all fine and single (except 4, which is usually 2-branched), 3 pairs placed well forward, a fourth pair, longer than others, further back, posterior to level of bases of *antennae*, latter short and stout, shaft smooth, a single hair at about $\frac{1}{2}$ length from base, 2 small preapical spines very near tip, 2 small spines at apex, and a papilla *Mouth-brushes* moderately developed, median hairs slightly hooked, and with minute serrations apically. *Mentum* narrow from before backwards, with large central tooth and 8-10 smaller on each side. Numerous stellate hairs on *thorax* and *abdomen*, each hair with about 12 branches, the tips apparently blunt, but actually ending in several minute points, some long lateral hairs also on *thorax* and *abdomen*, a strong single, slightly barbed spine on mesothorax dorso-laterally, a longer double strong spine arising from a large chitinated tubercle on metathorax, one arm longer than the other, actual length of spine, and of shorter arm compared with longer, variable, but longer arm usually twice or more length of shorter, both arms sharp-pointed *Comb* of 10-12 pointed teeth, of which about five are larger than others, a few towards dorsum widely spaced; indications, in most specimens, of a feebly chitinated lateral plate from which the comb arises. *Siphon* usually darker in middle than towards base and apex, a narrow dark ring at base, no acus, length of tube about 0.5-0.7 mm., $2\frac{1}{2}$ -3 times length of width at base; a double row of 2-branched hairs along posterior margin, the pair nearest base usually 4-branched, scattered single and 2-branched hairs on lateral and anterior surfaces; *pecten* of 5-8 fine teeth, pointed and minutely fringed, commencing some distance from base of tube and continued nearly to apex. *Anal segment* enclosed dorsally and laterally with chitinated saddle, posterior margin of which carries 9-12 rather long sharp spines, osc single, smooth, and long, *asc* with 5 or 6 fairly long branches, all

* In the type-form the line separating the dark and light parts of the *abdomen* is serrate, in the var. *serrata* it is jagged, the white area on each tergite being broader towards the base.

† de Meijere 1910, p. 939.

‡ Edwards 1926 a, p. 116; Senior-White 1927, p. 66; Edwards & Given 1928, p. 387; Brug 1931 a, p. 15;

about same length, *lh* about same length as *isc* and either 2- or 3-branched or single, pair of ventral hairs representing fan or brush each with 5-6 shorter branches, both pairs of anal papillæ nearly as long as lateral hair, tips slightly rounded

HABITAT—Bamboos, occasionally tree-holes In Malaysia found very commonly in pitcher-plants (*de Meijere*, *Given*, *Dover*) †, as well as bamboos (*Leicester*)

DISTRIBUTION—S INDIA Nagargah*, Belgaum dist., vii 1921 (*Barraud*), Yellapur* and Kadra*, N Kanara dist., x and xi 1921 (*Barraud*), Nilgiri Hills*, x 1915 (*Khazan Chand*) DARJEELING DIST.: Sureil* and Mungpoo*, c 5,000', x 1922, and Sukna*, 500', x 1922 (*Barraud*) ASSAM Shilong*, vi & viii 1922, 5,000', Haflong*, Cachar Hills, viii. 1922, and Golaghat*, Sibsagar dist., v 1925 (*Barraud*), Khumtai* Sibsagar dist., v 1925 (*Christophers*) BURMA Rangoon*, i 1920 (*Christophers*), and 1930 (*Feegrade*), west slopes of Dawna Hills*, iii 1908 (*Annandale*) ANDAMAN Is *, vii 1926 (*Sobha Ram*) CEYLON Peradeniya, x 1901 (*Green*), Colombo*, 1913 (*James*), Matale*, Suduganga Estate, xi. 1919 (*Senior-White*)

Also known from MALAYA, COCHIN CHINA, BORNEO, and JAVA

11 *Tripteroides affinis* (Edwards), 1913

Rachnotomyia affinis Edwards, Bull. Ent. Res. iv, p. 241 (nom. nov.)

Phonomyia caeruleocephala Theobald, 1910, M.C. v, p. 577 (♂ & ♀, wing fig.). TYPE-LOO, Hakgala, Ceylon, iii 1907 (*Green*)

TYPE ♂ & ♀ in Brit. Mus. (nec *Colonymyia caeruleocephala* Leicester, 1908, Cul. Malays, p. 233)

ADULT—Very similar both in markings and in structure of ♂ hypopygium to *T. aranoides*, but may be distinguished by the characters given in the key†. It should be noted that the scales on the pleuræ are distinctly silvery, a very unusual character among those species of the subgenus *Tripteroides* which lack spots on the femora.

PUPA.—Distinguished from *T. aranoides* by characters given in key

LARVA—Resembles *T. aranoides*, except as follows:—Comb of more numerous teeth, about 20 in a close-set row, spine on mesothorax, and both arms of spine on metathorax, ending in several minute points

HABITAT—Tree-holes (*Barraud*)

† [Larvæ from pitcher-plants differ in some respects (chiefly in the smaller number of branches of various hairs) from those described above from bamboos, and it may be that two species or races are involved (*Edwards* 1926, p. 117).]

‡ *Vide Barraud* 1929 a, p. 1060

DISTRIBUTION.—S INDIA Santi Kappa*, Coorg, v 1914 (*T. B. Fletcher*), Nilgiri Hills*, x 1915 (*Khazan Chand*), Belgaum*, Bombay Deccan, viii 1921 (*Barraud*) Ceylon Hakgala, c 6,500', iii. 1907 (*Green*); Peradeniya, v 1912 (*J. C. F. Fryer*), Rakwara, v 1929 (*Henry*)
Not known from elsewhere

[12 *Tripteroides doffeini* (Gunther), 1913.

Zeitschr. Wiss. Ins.-biol. ix, pp 204-7 and 259-269 (larva and pupa only) TYPE-LOC. Udugama, Ceylon (*Gunther*) TYPE Freiburg-i.-Br (?)

ADULT.—Unknown

LARVA.—Very fully described and figured by Gunther, in most respects closely resembles those of *T. aranoides* and *T. affinis*, but seems to differ from both in having only 5-6 branches in the abdominal stellate tufts. As, however, the number of branches in these tufts is liable to increase with the age of the larva, and the stage of Gunther's larvæ is not clearly stated, it is possible that they may prove to be second- or third-stage larvæ of *T. aranoides*. The following features noted in the original description may be of specific importance.—Median hairs of mouth-brush strongly combed. Mentum with about 13 approximately equal teeth. Metathoracic spine with a short basal branch, barely $\frac{1}{2}$ as long as the main spine, tip of latter slightly rounded, not split into points. Comb of 5 large, widely-spaced teeth (smaller ventral teeth not mentioned). Ventral hairs of anal segment single, *isc* with 2 branches only; *osc* finely barbed; *lh* single

HABITAT.—Pitchers of *Nepenthes destillatoria*

DISTRIBUTION.—Recorded only from type-locality.]

Genus **TOPOMYIA** Leicester, 1908

Cul Malaya, p 238. Genotype, *T. minor* Leic.

Pseudograhamia Theobald, 1910, Rec. Ind. Mus. iv, p 26. Genotype, *P. aureovenier* Theo.

ADULT †.—Small mosquitoes with slender legs, thorax usually ornamented with a median silvery stripe, and with silvery scales on *apn*, scutellum, and pleuræ. Microtrichia on wing-membrane of normal size, squama without hairs; wing-scales normal and not emarginate at tips; clypeus unusually small; antennæ alike in the two sexes, not strongly plumose, segments of flagellum subequal in length; palpi quite short in both sexes, proboscis not hairy, about as long

† Theobald 1910 b, p. 551; Edwards 1922 d, p. 437; 1932, p. 90; Brunetti 1914, p. 72; Barraud 1929 b, p. 270.

as abdomen, apparently bent under the body when insect is at rest, scutellum unusually small

LARVA†—[Some Malayan species are very similar to *Harpagomyia*, but *T. argenteoventralis*, which is closely allied to the Indian species, has a very peculiar larva, with the maxilla modified into a strong horn, siphon with long ventral spines, and anal papillæ very long and slender]

BIONOMICS‡—Adults are usually found near jungle-streams (*Leicester*), none has yet been found sucking human blood. larvæ breed in water in bamboos and leaf-bases and flowers of certain plants

DISTRIBUTION—The genus is purely Oriental; about 13 species are known, and nearly all these occur in Malaya, one of these has also been found in South India.

13 *Topomyia aureoventer* (Theobald), 1910.

Pseudograhamia aureoventer Theobald, Rec Ind Mus iv, p 27 (♀)
TYPE-LOC Pallode, 20 miles north-east of Trivandrum,
Travancore (*N Annandale*) TYPE ♀ in Ind Mus

ADULT§.—May be identified on generic characters and on markings, as given below ||. Wing 3.5 mm

♀—*Head*: flat-scaled, a large triangular patch of silvery scales in middle of vertex, remainder covered with brownish-black scales, palpi quite small, clothed with pale scales; proboscis black, slightly swollen at tip *Thorax* mesonotum deep brown, with a median silvery line of flat scales, lateral areas of ochreous narrow scales towards the front extending to bases of wings, flat silvery scales on *apn*, pleuræ, and mid-lobe of scutellum, flat golden scales on *ppn*. *Legs*. brownish-black, femora paler beneath *Wings* with rather scanty dark scales, forks rather long, with broadish scales, vein 6 scarcely beyond level of base of fork of 5 *Abdomen*. laterally compressed, large lateral apical silvery patches on tergites I, II, IV, V, and VI (none on others) not forming complete bands on dorsum, sternites dull golden

♂ and early stages unknown (see note under genus).

DISTRIBUTION—In India known only from type-locality* given above, and from Virajpet*, Coorg, 20–24 x. 1915 (*T. B. Fletcher*)

Also occurs in MALAY PENINSULA

† Brug 1931 b, p. 8; Edwards 1932, p. 90

‡ Leicester 1908, p. 239; Brug 1913 b, p. 8, Edwards 1932, p. 90.


§ Edwards 1913 b, p. 240, and Barraud 1929 b, p. 270 (as *T. argenteoventris*), Theobald 1910 b, p. 551 (*Pseudograhamia*)

|| It differs from *T. argenteoventralis*, with which it was at one time confused, by the absence of a large patch of silvery scales on the dorsum of the 2nd abdominal tergite and shorter 5th vein. For distinctions between the various other Malayan species, see Edwards 1932 d, p. 438.

Genus **HARPAGOMYIA** de Meijere, 1909.Tijd v Ent. iii, p 165 Genotype, *H. splendens* de Meij*Malaya* Leicester, 1908, Cul Malaya, p 258 (nec *Malaya* Hellor).Genotype, *M. genurostris* Loew

This genus includes only two Indian species*. They are small mosquitoes ornamented with brilliant silvery, blue or violet scales, the hairy proboscis is peculiar in having a flexible joint and a very much enlarged tip carrying four long hairs (fig. 12, b). This character alone distinguishes these mosquitoes from all other Culicines

Fig 11



Harpagomyia taking food from mouth of an ant (*Cremastogaster*).

(After Jacobson, reproduced by permission of Messrs. F. Warne and Co. from 'Standard Natural History.' The raised hind tibia of the mosquito is shown too long in the figure.)

PUPA †.—Distinguishing characters are given below under *H. genurostris*.

LARVA †.—In 4th stage this resembles those of *Topomyia* and *Trypteroidea* in the absence of an anal fan or brush, but

* See also Theobald 1910 b, p. 547; Edwards 1922 a, pp. 496-503, 1930 b, p. 548, and 1932, p. 91; Barraud 1926 p. 348; Barraud & Covell 1928, p. 676 (buccal cavity).

† de Meijere 1911, p. 162.

‡ de Meijere 1911, p. 162; Ingram & de Meillon 1927, p. 76; Evans 1929, p. 467.

differs from the latter in the absence of large spines on the thorax. There do not appear to be any constant characteristics by which the larvæ may be distinguished from those of *Topomyia*. Important points of structure are given in the descriptions below.

BIONOMICS* —The following details are given by Edwards (1932) —Larvæ have been found in tree-holes, in old water-filled nests of tree-dwelling ants, in leaf-bases of pineapple and other plants. The habits of the adults are of exceptional interest, and were first observed in detail by Jacobson in Java, having been confirmed subsequently by James in Ceylon and Farquharson and others in tropical Africa. The flies haunt tree-trunks where ants of the genus *Cremastogaster* are found, and obtain their food from the ants; the proboscis of the mosquito is evidently highly specialized for this purpose, and they probably do not feed in any other way. The *Harpagomyia* places itself directly in front of an advancing ant, sometimes even nipping the ant between its front legs (fig. 11), and not releasing it until it stops and opens its jaws, when the mosquito thrusts the swollen tip of its proboscis into the ant's mouth and rapidly absorbs the food offered. While the interchange is taking place the ant strokes the tip of the mosquito's proboscis with its palpi. When not in use, the proboscis of the mosquito is folded backwards under the body —a most unusual position in this family. Whilst feeding, the wings are vibrated and the hind legs held high so that the tarsi curve forwards over the head, as in *Wyeomyia* and related genera.

DISTRIBUTION —In India the genus appears to be confined to the warmer southern and eastern regions of heavy rainfall. Two other species are known from the Oriental region, and four from Tropical Africa.

14. *Harpagomyia genurostris* (Leicester), 1908

Malaya genurostris, Cul. Malaya, p. 258 (♂) TYPE-LOC Malay Penin., caught in bungalow (Dr. Daniels) TYPE ♂ in Brit. Mus.

ADULT† —Differs from *H. jacobsoni* in having clypeus yellowish-white, a line of silvery scales between eyes, a larger silvery patch laterally on tergite IV, and scales of *apn* and on front of vertex silvery white, with only a faint bluish tinge. Wing 2·2–2·4 mm.

* Jacobson 1909 a, p. 246; 1909 b, pp. 158–164, 1911, pp. 158–161; de Meijere 1909, p. 169, 1911, pp. 162–167; James 1914, p. 233; Farquharson 1919, pp. xxxix–xxxix, Edwards 1932, p. 92.

† See also Edwards 1913 b, p. 240, and 1930 b, p. 543; Barraud 1926, p. 349.

♀—*Head* a subtriangular patch of silvery flat scales in middle of vertex in front, with a slight bluish or violet tinge in some specimens, the scales continued as a narrow line downwards between eyes, a patch of similar scales at each side of head, remainder covered with brownish-black flat scales, a wide space separating the two vertical bristles from the lower orbital bristles, antennæ dark brown, a little longer than proboscis, hairs fairly long, clypeus fully twice as long as width at base, rather pointed and beak-like, yellowish, with silvery prunescence, palpi a little longer than clypeus, pale yellow or creamy, proboscis comparatively short, about 1 mm, mainly yellow, except the swollen part, which is brown. *Thorax* integument of mesonotum dark brown, clothed with brown narrow scales, a sharply-defined median line of flat bluish-violet scales with metallic lustre from front margin to about middle of mesonotum, postnotum and pleura dark brown, latter, as well as coxæ, almost entirely covered with flat silvery scales, those covering *pm* with a yellowish tinge, on *apm* with a bluish or purplish tinge. *Wings*. dark scaled, termination of vein 6 usually a little nearer tip of wing than level of fork of 5 and base of 2, but in some specimens it ends at about level of those points, *af* longer than *pf*. *Legs* dark brown, undersides of femora paler, all legs with yellowish sheen when seen in certain positions, segment 1 of hind tarsus a little longer than tibia. *Abdomen* tergites brownish-black, with large lateral silvery patches on I and II and IV-VI, that on IV being larger than others, those on V and VI apical, VII nearly all silvery, venter silvery with yellowish sheen.


♂.—Integument of mesonotum usually rather lighter than in ♀, and the median silvery purple line not always so well pronounced, only tip of swollen apex of proboscis definitely dark, vein 6 usually ending beneath fork of 5 and base of 2 or slightly nearer base of wing than this. Antennæ similar to those of ♀. *Hypopygium*. *9t* with a pair of submedian hairy lobes; coxite short and broad, a basal lobe on inner side bearing 4 strong bristles in a row and 2 similar bristles above, style (fig. 12, a) moderately long with short terminal appendage; *lp* with small teeth at apices; paraprocts pointed, without teeth or hairs.

PUPA.—Paddles feebly chitinised, semi-transparent, somewhat triangular and pointed, without terminal fringe or hair, part inside midrib rather narrow; lateral tufted hair on outer posterior corners of VII and VIII well developed; one of the hairs on posterior borders of V and VI long and single, corresponding hair on IV also single and moderately long; dendritic tufts on tergite I only moderately developed;

respiratory trumpet short, about 3 times length of width, of normal shape

LARVA (fig 12, g-j) —*Head* towards front of clypeus there are three pairs of fine hairs in a somewhat convex row, the outer (A) and inner each 3-branched, the mid single or 2-branched, posterior to these there is a pair of single or

Fig 12



Harpagomyia genurostris: style of ♂ hypopygium (a) and proboscis (b); also details of larval structure of *H. jacobsoni* (e-f) and *H. genurostris* (g-j). e, i, comb-scales, d, h, pecten-teeth

2-branched fine hairs internal and slightly posterior to A and at about level of bases of antennæ; posterior to these four pairs there is a fine hair on each side at about level of upper margin of eye, preclypeal spines moderately long and stout, tips pointed. Antenna short, slightly curved, shaft smooth, a single hair some little distance from tip, several apical bristles. A number of median hairs of mouth-brush

hooked, and with minute serrations on one side apically. Mentum narrow from before backwards, a fairly large median tooth and 12 smaller on each side. *Thorax* lateral tufted hairs well developed, two fan-shaped tufts, one larger than the other, on prothorax, arising from tubercles. *Abdomen* lateral tufted hairs on segments I and II fairly long, those on following segments with fewer branches, comb composed of about 40 teeth in a large subtriangular patch. *Siphon* about 0.7 mm long, slightly curved and tapering, about 4 or 5 times length of basal width, scattered hairs over surface, one pair with 4 or 5 fairly long branches towards posterior margin at about middle, a few 2- or 3-branched hairs on anterior and lateral surfaces, and a row of single and some smaller branched hairs between median tuft and apex on posterior surface. Pecten variable, in some specimens an irregular group of small transparent fringed teeth towards base, and a widely spaced row of 4 or 5 larger teeth, also minutely fringed, on rather more than basal $\frac{1}{2}$ of tube, in other specimens the smaller teeth towards base appear to be absent. *Anal segment* enclosed in a chitinous ring, *asc* of 6 or 7 branches, *osc* single, *lh* with one very long branch and one much finer and shorter, a patch of spines on margin of ring on each side, a pair of fine single ventral hairs in place of fan or brush, one or both occasionally split into two. Both pairs of anal papillae very long, about twice length of anal segment and nearly as long as siphon-tube, moderately wide, tips rounded.

HABITAT.—Water in leaf-bases of a large *Arum* (*Barraud*).

DISTRIBUTION.—BIHAR: Pusa*, 2 xii 1926 (*Mathur*), CHITTAGONG HILL TRACTS: Rangamatti*, ix 1922 (*Barraud*); New Forest*, Dehra Dun, U.P., 10. iv 1930 (*Mathur*). BURMA: Rangoon*, 13 xi. 1930 (*Feegrade*). CALCUTTA*: North-West Soap Co's Factory, Matlabruz, near west end of King George's Dock, 25 v. 1931 (*Senior-White*). ASSAM: Golaghat*, Sibsagar dist., 14. x. 1924, 1 & xi. 1925 (*Barraud*). CEYLON: Colombo*, x. 1913 (*James*).

Only known with certainty elsewhere from MALAY PENINSULA.

15. *Harpagomyia jacobsoni* Edwards, 1930

Bull. Ent. Res. xxi, p. 543 (♀). TYPE-LOC.: Fort de Kock, Sumatra, 1926 (*E. Jacobson*). TYPE: ♀ in Brit Mus.

ADULT.—Resembles *H. genurostris*, but differs as follows:—Larger and darker, wing 3–3.5 mm.; clypeus dark, both this and tori with pale pruinescence, no scaled line between eyes, scales on front of vertex and on *apn* definitely blue or bluish-violet, not silvery-white with bluish sheen,

integument of mesonotum and of pleuræ black, swollen apical part of proboscis entirely dark in both sexes, lateral silvery patch on abdominal tergite IV smaller

PUPA and Egg —Unknown

LARVA —Differs from *H. genurostris* as follows —The five pairs of frontal hairs have much the same arrangement, but are all single and somewhat thicker; comb of about 16–20 teeth arranged more or less in two rows, anal papillæ more pointed, ventral hair of anal segment with 3 fairly long branches. In other details the resemblance to larva of *H. genurostris* is very close

HABITAT —Water in leaf-bases of a large species of *Arum*, in association with *H. genurostris* (Barraud)

DISTRIBUTION —CHITTAGONG HILL TRACTS Rangamatti*, ix 1922 (♀) (Barraud). DARJEELING DIST, Marianbarrie Tea Estate*, near Sukna, viii 1928 (♀) (Sobha Ram), Rangamatti*, Jalpaiguri dist., 10 xi 1929 (♂) (Sinton)

Originally recorded from SUMATRA, not known from elsewhere

Genus **HODGESIA** Theobald, 1904

Journ Trop Med vii, p 17 Genotype, *H. sanguina* Theo.

ADULT †.—Very small mosquitoes, distinguished from all other Culicini by the form of the outstanding scales on the outer $\frac{1}{2}$ of the wing, these scales (fig 13, c) are narrow and emarginate at their tips, and appear as though the sides were prolonged into minute spines beyond the end of the scale. As in *Harpagomyia* and *Uranotaenia*, vein 6 is short and ends at about level of fork of 5 and base of 2 (fig 13, a), a line drawn through these points forming a right-angle with the costa, *af* about length of its stem, squama quite bare (fig. 13, b). Proboscis normal, not swollen at tip, and about length of abdomen; palpi very short in both sexes, antennæ alike in both sexes, not plumose, all flagellar segments subequal in length. Pleural bristles much reduced, no spiracular, postspiracular, or prealar, one strong lower mesepimeral bristle and one or two upper, two strong posterior pronotal, one sternopleural, or this may be absent. Postnotum without setæ. Tarsal claws simple. Coxite of ♂ hypopygium short and without lobes, style without distinct appendage, phallosome simple, paraprocta strongly developed, with 5 or 6 strong teeth at crown, cerci of ♀ small, usually hidden, only one chitinated spermatheca.


† Theobald 1907, p 679, Leicester 1908, p 229, Barraud 1929 a, p 1081; Edwards 1930 a, p 299, 1932, p 93.

PUPA*.—Those of Indian species are unknown, the pupæ of two African species have been collected and show the following characters —Respiratory trumpets (fig 14, a) cleft almost to base into two leaflets of unequal widths, the narrower one notched at tip, abdomen with conspicuous dark pattern, VII and VIII without lateral tufts, paddles notched at tip, serrate round apical margin, about equally divided by strong midrib, at tip of which is one hair

LARVA* (fig 14) —The chief points of structure are given below in the description of that stage of *H. malaya*

BIONOMICS —Nothing is known regarding the habits of the two species occurring in the Indian area. Larvæ have been found in small pools in swampy ground and in pools in jungle*.

Fig 13.



Hodgesia malaya: a, wing, showing venation, b, base of wing, showing bare squama, c, scales from lower fork

DISTRIBUTION.—Five species are known from the Oriental and Australasian regions, of which one occurs in S. India and one in Ceylon. Four other species are known from Tropical Africa.

Key to Adults.

Integument of mesonotum and of pleura uniformly reddish-brown or ferruginous	<i>malaya</i> , p. 54.
Integument of mesonotum pale yellowish, with a large brownish-black oval area over, and in front of, each wing-base; sternopleura and lower part of mesepimeron black, pleura otherwise yellowish	<i>bailyi</i> , p. 55.

* Wigglesworth 1929, pp. 60-62; Edwards 1932, p. 93.

16 *Hodgesia malaya* Leicester, 1908


Cul Malaya, p 231 (♂ & ♀) TYPE-LOC Kuala Lumpur, Malay Penn (Leicester) TYPE ♂ & ♀ in Brit Mus

Hodgesia ampya Dyar, 1920, Insec Inse Mens viii, p 176 (♀)
TYPE-LOC . Philippine Is TYPE two co-type ♀♀ in US Nat Mus

ADULT* — Small species with unmarked abdomen† Wing 2-2.5 mm (fig 13)

♀ — *Head* flat-scaled, scales along eye-margins and on vertex white, on nape dark brown, torus and flagellum light

Fig 14



odgesia, pupal and larval characters a, respiratory horn of pupa, *H. nigerica*, b-f, *H. malaya*, larval details, c, pecten-teeth, f, comb-teeth (NB — The row of fine ventral hairs on the anal segment is accidentally omitted in fig b)

brown, hairs dark, clypeus reddish-brown, palpi and proboscis dark brown, palpi very short *Thorax*. integument of mesonotum reddish-brown clad with bronzy-brown narrow scales and dark brown bristles, scutellar scales brown, integument of pleurae reddish-brown, with several patches of silvery-white scales, similar scales on *apn* *Wings*: venation and

* Dyar & Shannon 1925, p 70 (syn), Edwards 1930 a, p 299

† [The other Oriental species, *quasisanguinea* Leic, has silvery markings on abdomen]

scales typical *Legs*: dark brown, femora conspicuously lighter *Abdomen* dark brown, without bands or spots.

♂—Resembles ♀ very closely

PUPA and EGG—Unknown

LARVA * (fig 14)—Preclypeal spines moderately long and stout, the five pairs of frontal hairs all placed rather far back, *A* with about 6 short branches, *B* single and long, *C* placed almost directly behind *B*, with about 8 short branches, *d* internal and slightly anterior to *B*, with 3 fairly strong branches, *e* external and slightly anterior to *C*, with 2 or 3 fine branches Mouth-brushes relatively large *Antenna* of moderate length, slightly curved, shaft with small spines and 4 bristles very near tip (3 long and 1 short), the longer bristles with numerous short spines, tuft at $\frac{5}{8}$ of length from base *Mentum* rectangular, with 4 teeth on either side of large median tooth. *Thorax* with relatively large tufts of slightly plumose hairs *Comb* of about 7 teeth arranged in a row, teeth pointed, with delicate lateral fringes *Siphon* short and cylindrical, about same length as anal segment, valves comparatively large, pecten of about 6 stout spines, without lateral denticles, along about basal $\frac{1}{2}$ of tube, tuft arises very near base of tube, divided into about 8 rather stout branches, of equal length with siphon Anal segment completely enclosed in chitinous ring, both *isc* and *osc* divided into a number of branches, *th* of 2 or 3 stout branches, one at least of which is spinous, a row of several fine hairs along ventral surface of segment (omitted in figure), as well as a fairly well-developed fan or brush, latter of about 6 fairly long-branched hairs, no fan-plate

HABITAT.—Junglo-pools (*Leicester*)

DISTRIBUTION.—CEYLON. Colombo, 1913 (*James*)

Not known in India, but occurs in MALAY PENINSULA and PHILIPPINE ISLANDS

17 *Hodgesia baillyi* Barraud, 1929

Ind Journ Med. Res. xvi, p 1062 (♀). TYPE-LOC. Virajpet, Coorg, South India, vi. 1927 (*J. D. Bailly*) TYPE ♀ in Brit. Mus

ADULT —Wing 2.5 mm

♀—*Head*: flat-scaled, a broad band of silvery scales along eye-margins, widest in middle, where it occupies about $\frac{1}{2}$ the dorsal surface, behind the straight posterior margin of this band there are black scales continued to nape. Tori brown, flagellar segments and hairs of antenna dark brown; clypeus dark brown; palpi and proboscis brownish-black.

* Wigglesworth 1929, p 63.

the former minute *Thorax* integument of mesonotum area pale yellowish on anterior $\frac{1}{2}$, a large oval brownish-black over and in front of each wing-base, scutellum and a small space on mesonotum in front of it yellowish, a sparse covering of black hair-like scales and dark bristles on mesonotum, postnotum dark brown, *ppn*, postspiracular area, upper part of mesepimeron, and coxæ yellowish, sternopleuræ and lower part of mesepimeron black, upper part of dark area overlaid with patches of flat silvery scales, similar scales on *apn*, two black *ppn* bristles. *Wings* scales and venation typical. *Legs* tibiae and tarsi dark brown, with bronzy metallic sheen, fore femora light on about basal $\frac{1}{2}$, otherwise dark brown, mid-pair dark brown anteriorly, pale posteriorly and ventrally for whole length, hind pair nearly all white, except for a short dorsal brown streak to the knee. *Abdomen*, entirely bluish-black, with metallic lustre, blunt at extremity owing to retraction of terminal segments, cerci very small and completely hidden, one chitinated spermatheca.

♂ and early stages unknown

HABITAT —Small pool in swampy area (*J D Barry*)

DISTRIBUTION —Known only from type-locality

Genus **URANOTÆNIA** Lynch Arribalzaga, 1891.


Rev Mus La Plata, 1, p. 375. Genotype, *U pulcherrima* L. Arrib.
Anisacheleomyia Theobald, 1905, Entomologist, xxxviii, p. 52.
 Genotype, *A. nigripes* Theo.

ADULT* —Small mosquitoes, distinguished from all other Culicinae by form of microtrichia on wing-membrane; these are very much smaller and denser than in any other genus, requiring a magnification of more than 80 for their detection, at this magnification the membrane appears clear, vein 6 short (fig. 15) and curved down to wing-margin at tip, this point being usually nearer base of wing than level of commencement of 2 and fork of 5; *af* usually very short, always much shorter than its stem; squama without hairs. *Palpi* in both sexes quite short. *Proboscis* often slightly enlarged at tip, especially in ♂, and about length of abdomen, except where otherwise mentioned in descriptions. *Antennae* usually a little longer than proboscis in ♀, segments subequal in length and setose, about length of proboscis in ♂, plumose, last two segments elongate. *Eyes* touching above antennae and below

* Theobald 1901, p. 241, Leicester 1908, p. 203, Edwards 1912 a, p. 37; 1921 c, p. 282, 1932, p. 96, Howard, Dyar, & Knab 1917, p. 898, Kirkpatrick 1925, p. 67; Barraud 1926, p. 331, Barraud & Covell 1928, p. 676 (buccal cavity), Martini 1930, p. 196.

proboscis Orbital bristles not numerous, but the row not conspicuously interrupted. Vertex and nape covered with broad, flat, and upright scales, the latter variable in number, in some species few and confined to nape, in others more numerous and extending forwards nearly to eye-margins Dorsum of thorax clothed with narrow scales and bristles, in some species there are flat scales along lateral margin in front of wing-root, or pale narrow scales in this position, which may or may not form a continuous border round the front margin, the presence or absence of such scales is of importance in identification. *apn* widely separated, with usually only 3 bristles, usually only 1 *ppn*, 1 spiracular (absent in some species, as mentioned in descriptions*), no postspiracular, about 2 prealar, several upper sternopleural, 1 or 2 upper and 1 lower mesepimeral Upper margin of meron well above base of

Fig 15



Wing-venation of *Uranotania*, showing notation used in this volume
(Note shortness of vein 6 in this genus)


hind coxa. Postnotum bare Scutellum, in all Indian species, covered with flat dark brown scales, at least on the mid-lobe. All tarsal claws of both sexes simple, those of fore and hind legs in both sexes (except claws of fore leg of ♂ of *U unguiculata*) short and equal, but one sometimes broader than the other, claws of mid-leg of ♀ equal, of ♂ unequal (in *U edwardsi* ♂ both are very long and equal) No pulvilli. In several species the tarsi of one or more pairs of legs are specialized in the ♂ (fig 17), whilst in others the comparative length of tibia and of segment 1 of tarsi of fore and hind legs form easily seen distinctions *Hypopygium* of ♂† (fig. 18) with coxite short and broad, with lobe on inner side bearing several long thick bristles; style of varying length in different

* This character may be variable, only a few specimens of each species have been available for examination.

† Macfie & Ingram 1922, p. 184; Christophers 1922, p. 566; Kirkpatrick 1926, p. 69, Barraud 1926, p. 381, Martini 1930, p. 196, Edwards 1932, p. 196.

species, with terminal appendage usually quite small, phallosome divided into lateral plates bearing a few teeth, anal segment membranous, without distinct paraprocts. Only one chitinated spermatheca in ♀, as in *Hodgesia* and *Ficalbra*.

Fig 16



Larval structures of *Uranotenia* (head, mentum, comb-plate, and pecten-teeth) a-d, *atra*, e-h, *unguiculata*, i, *annandalei*; j-l, *recondita*, m-p, *maxima*

PUPA* —Among Indian species only that of *U. unguiculata* is known, a short description is given under that species. In this, and in the majority of pupæ described from other regions, the respiratory trumpet is rather short, with small

* Edwards 1932, p. 96. See also under *unguiculata* and other species.

opening, paddle without distinct fringe, though apical margin may be serrated, part of paddle inside midrib wider than that outside, one small terminal hair

LARVA, 4th stage* (fig 16)—The larvæ of only eight or nine Indian species are known, these show fairly well-marked generic characteristics—head small, antenna short, preclypeal spines usually stout and short, arising from projections of preclypeus. In many species one or more pairs of frontal hairs are stout, in some cases resembling flattened leaf-like bristles. Lateral hairs on thorax and on segments I and II of abdomen fairly well developed. Abdominal segment VIII bears a comb on each side, the teeth usually arising in a row along the posterior margin of a chitinated plate. Siphon with one pair of hair-tufts at about middle, aous well developed, pecten present, the teeth of characteristic form, usually scale-like, wide at tip, with lateral and terminal delicate fringes. Anal segment enclosed in a chitinous ring, the hind margin spinose; both pairs of subdorsal hairs of 2 or more branches, anal fan small.

EGG—The eggs are laid in boat-shaped masses. Those of Indian species are unknown.

BIONOMICS*—Little is known of the habits of the adults; they do not appear to attack man to any extent. The larvæ of some species live in stream-pools and swamps, whilst those of several others have been taken from tree-holes, bamboos, and pitcher-plants. From their habit of resting almost horizontally in the water just below the surface they may usually be distinguished from other Culicine larvæ, but may be mistaken for those of *Anopheles*.

DISTRIBUTION—Members of the genus are found in most parts of the Indian area, but are most abundant in the regions of heavy rainfall. Numerous species are known from the tropical and subtropical regions of both the New and Old Worlds.

Key to Adults.

- | | |
|---|-------------------------|
| 1. Hind tarsi with white markings, or pale at tip | 2. |
| Hind tarsi entirely dark | 6 |
| 2. No white scales on wings, hind tarsi pale at tip only (segments 4-5 and distal part of 3) | <i>testacea</i> , p. 74 |
| Some white scales on wings, segments 2-4 of hind tarsi with pale markings (5 or 4-5 all pale) | 3 |
| 3. Costa of wing with pale areas | 4 |
| Costa entirely dark; proboscis dark; segments 1-3 of hind tarsi with basal and apical white rings | 5 |

* Senior-White 1920, p. 319; Edwards 1932, p. 97

4. Proboscis mainly white above, segments 1-3 of hind tarsi with apical white rings only (4-5 white) *alboannulata*, p. 62.
 Proboscis dark segment 1 of hind tarsi all dark, 2-4 with basal white marks only *edwardsi*, p. 65
5. Hind tibiae with 3 white patches, femora without white knee-spots *rutherfordi*, p. 63
 Hind tibiae with an apical white ring only, a white knee-spot on each femur *christophersi*, p. 64.
6. Some white, blue, or creamy flat broad scales along lateral margin of mesonotum in front of wing-root 7
 No pale or blue broad flat scales along margin of mesonotum (though there may be narrow pale scales in this position) 13
7. A border of flat creamy scales along lateral margin of mesonotum from wing-root, continued round front, wings mainly pale scaled *orientalis* p. 66.
 A line of white or blue flat scales not forming a complete border round front of thorax, wings mainly or entirely dark scaled 8
8. Flat scales along lateral margin of mesonotum not continued anteriorly beyond *ppn*; mid-femur without a pale longitudinal line 9
 Flat scales continued nearly to neck, a pale line along basal $\frac{1}{2}$ of anterior surface of mid-femur *unguiculata*, p. 67
9. Proboscis longer than whole body, abdomen black, with large lateral apical white patches (which may unite on dorsum) on segment IV only *longirostris*, p. 71
 Proboscis of moderate length, abdomen with either pale lateral patches, or median pale markings dorsally, on more than 3 segments 10
10. Lateral apical markings on abdominal tergites, no median pale markings on dorsum, foretarsi of ♂ peculiar (fig. 17, a) *ata*, p. 72
 Median pale markings on I-IV, V with complete apical white band, foretarsi of ♂ normal 11
11. Dorsum of mesonotum covered with a mixture of pale brown, ochreous, and dark brown scales *macfarlanei*, p. 70.
 Dorsum of mesonotum covered with deep brown scales only 12.
12. Tergites II-IV with lateral pale spots separate from the median pale apical markings [p. 69.
campestris var. *zelena*,
 Tergites II-IV without separate lateral pale spots [p. 68.
campestris (type-form),
13. A line of bluish-white scales across pleurae, a patch of rather broad greyish-brown scales over wing-root *annandalei*, p. 75
 No line of bluish-white scales across pleurae, no patch of greyish-brown scales over wing-root 14

- 14 Abdomen banded 15
 Abdomen unbanded 18
15. Bands well defined on all segments, no border of pale narrow scales round margin of mesonotum 16
 Bands faintly indicated, but usually definite on one or more terminal segments, a border of pale narrow scales from wing-roots round front of mesonotum *maculima*, p 77
luteola, p 80
- 16 Pleurae uniformly pale 17
 Pleurae with dark patches
- 17 Pleurae pale yellow, with conspicuous brownish-black markings, mesonotum dark brown, with scales of same colour, except at sides in front, where they are lighter *stischlandi*, p 80.
 Pleurae dark brown, with slightly paler and darker areas, mesonotum brown, covered with pale ochreous scales *viridis*, p 83
- 18 Some narrow pale scales on front or lateral margins of mesonotum 18
 No pale scales on front or lateral margins of mesonotum 22
- 19 A large velvety black spot in front of each wing-root *bimaculata*, p 78.
 No large black spot in front of wing-root 20
20. Pale scales on front and lateral margins of mesonotum forming an almost continuous border from wing-root round front margin, mesonotal bristles rather long, curved, and conspicuous 21
 Pale lanceolate scales on margin of mesonotum for short distance in front of wing-root only, none on front margin, mesonotal bristles rather short and inconspicuous *hebes*, p 81
nervipectus, p 76
- 21 Pleurae uniformly pale *maxima*, p 77.
 Pleurae with a dark band across the middle, lower $\frac{1}{2}$ of mesopneuron dark
- 22 Pleurae with conspicuous dark markings on a pale ground *maculipleura*, p 82
 Pleurae uniform in colour 23
23. Mesonotal bristles long, curved, and dense, pleurae pale brown, dorsum of abdomen brown, venter paler *recondita*, p 83
 Mesonotal bristles neither long nor dense, pleurae rather dark brown, dorsum and venter of abdomen dark brown *novobscura*, p 84.

Key to known Larvæ (4th stage).

- 1 Comb-teeth with fringed rounded ends *recondita*
 Comb-teeth ending in a single sharp point 2
2. Numerous strong stellate hairs on thorax and abdomen, the tips of branches stout and ending in several minute sharp points. *maxima*.
 Hairs on thorax and abdomen not so 3
3. Pecten of 17-20 teeth 4.
 Pecten of 9-14 teeth 5.

- 4 Antennal hair single, at about middle of shaft
Comb of 8-9 teeth, of which lateral are smaller
than median Siphonal tuft of about 8
branches; *U* 4-branched *unguiculata*
Antennal hair with several fine branches, at
about $\frac{3}{4}$ length of shaft from base Comb
of 7 fairly large subequal teeth Siphonal
tuft 3-branched, *U* 2-branched *novoboscua*
- 5 Frontal hairs *B* and *C* very stout, resembling
flattened leaf-like bristles 6
Frontal hairs *B* and *C* less stout, not flattened,
resembling finely barbed bristles *atra*
- 6 Antenna with three large, leaf-like appendages
Plates of VIII connected dorsally *annulata*
Antenna normal Plates of VIII separate 7
- 7 Pecten of about 14 teeth Frontal hair *B* pos-
terior to *A* *macfarlanei**
Pecten of about 11 teeth Frontal hair *B* in
transverse line with *A* [*campestris**,
(*longirostris* ?),

18 *Uranotaenia alboannulata* (Theobald), 1905.

Anisopheleomyia alboannulata Theo, Entom xxxviii, p 51 (♂, 6g
ungues fore leg) TYPE-LOC India (James) (probably Travancore)
core) TYPE ♂ in Brit Mus.

ADULT † —A rather small blackish species, highly ornamented with white or bluish-white scales on proboscis, head, body, and legs, quite distinct from any other in India. Wing about 2.2 mm †

♀ —*Head*: a narrow border of bluish-white scales to eyes, widest at sides, some very long white scales projecting between eyes in front, a small patch of pale scales in middle of vertex towards nape, brownish-black scales over remainder of vertex, some dark upright scales towards nape, palpi a little longer than clypeus, upper side of proboscis white scaled, except for a dark interruption near middle and a dark tip, underside mainly dark. *Thorax*: integument of mesonotum black, clothed with dark brown narrow scales, with a narrow, sharply-defined border of bluish-white flat scales from wing-roots all round margin, a line of bluish flat scales across pleurae, continuous from head to base of abdomen, including *apn*,

* The key is based on scanty material, and in some cases it has been necessary to rely on descriptions and drawings by other writers. In Senior-White's figure of the larva of *campestris* and in Brug's figure of the larva of *macfarlanei* no lateral plate on VIII is shown, but these may have been omitted from the drawings. Of *longirostris* only a third-stage larva has been examined.

† Theobald 1907, p 573, Edwards 1913 b, p 238, and 1922 b, p 91 (syn.), Barraud 1926, p 334.

‡ A Malayan species (*U. trilineata* Leic.) is very similar, but differs in having a white ring in middle of segment 1 of hind tarsus.

sternopleura, and mesepimeron, another line of similar scales immediately above coxæ, integument of pleuræ almost black. *Wings* a rather large patch of white scales anteriorly in middle of wing from costa to cross-veins, another patch at tip of wing, including costa, fork-veins, and apical third of 3, some white scales at base of wing on 1, 4, 5, and 6. *Legs* femora dark brown with a line of white scales from base extending nearly whole length of each, a white knee-spot and a dark interruption between this and end of white line, on fore and mid-femora the line is along dorsal edge on anterior surface, and on fore femur there is a white spot between end of line and knee-spot, a white line on posterior surface of mid-femur, a short white line on apical half of posterior surface of hind femur, tibiae of fore and mid-legs dark, lighter at tips of tibiae, three wide white markings on hind tibia not forming complete rings beneath, one at apex, one in middle, and one near base, the actual base being brown, as are also the spaces between the white markings and underside, tarsi of fore and mid-legs brown, appearing very pale in certain positions, segments 1 and 2 of hind tarsi with rather wide apical white rings, the tip of 3 and whole of 4 and 5 white. *Abdomen* bluish-white scales covering tergite I and base of II, pale scales continued as a median longitudinal band on II, III, and IV, widening out into an apical band on IV, a pale apical band on VI, very small pale lateral patches on V and VII, remainder brownish-black, venter dark brown.

♂—Very similar to ♀, wing about 1.8 mm, antennæ plumose and about length of proboscis, palpi as in ♀. *Hypopygium* style short and wide (fig. 18, a).

LARVA and PUPA.—Unknown.

DISTRIBUTION.—S INDIA: Malabar Coast*, x 1915 (*Khazan Chand*), Kerwar, N Kanara dist, 1901–5 (*Cogill*), Mudigere, Mysore*, 1931 (*Anantaswami Rao*), Travancore (?) (*James*).

Not recorded from elsewhere.

19. *Uranotænia rutherfordi* Edwards, 1922.

Ind. Journ. Med. Res. x, p. 288 (♀). TYPE-LOC: Peradeniya, Ceylon, 30. iv 1914 (*A. Rutherford*). TYPE. ♀ in Brit Mus.

ADULT †.—Resembles *U. alboannulata*, but differs as follows.—Proboscis entirely dark, femora without white knee-spots, the mid as well as the fore pair with a white spot between the end of the longitudinal line and knee, the first of the three white markings on hind tibia situated at base, and not slightly removed from it, segments 1 to 3 of hind

† See also Barraud 1926, p. 335.

tarsi narrowly white at both base and tip, costa of wing entirely dark

♂ and early stages unknown

DISTRIBUTION —Known only from the type-locality

20 *Uranotænia christophersi* Barraud, 1926

Ind Journ Med Res xiv, p 336 (♂ & ♀, ♂ hypop fig) TYPE-
LOC Andaman Is, ix 1911 (*Christophersi*) TYPE ♂ & ♀ in
Brit Mus

ADULT —Diagnostic points are given in key The ♂ may be recognized by modifications of tarsi Wing about 2 mm

♀ —*Head* a line of bluish-white flat scales on sides from near *apn*, continued forward to eye-margins in middle, some long white scales projecting forward between eyes, brownish-black flat scales and a few black upright scales covering remainder of upper surface, a patch of brownish-black flat scales low down at each side, palpi scarcely exceeding clypeus in length, proboscis and antennæ dark brown
Thorax mesonotum dark brown, covered with dark brown scales, a moderate number of dark bristles, a thin line of broad white flat scales along margin from wing-root to *ppn*, integument of pleuræ dark, broad white scales on *apn* continued as a line across pleuræ to mesepimeron, small patches of similar scales on coxæ *Wings*. white scales along basal $\frac{1}{2}$ of stem of vein 5 and at extreme base of 4, cross-veins light in colour, otherwise wing is dark *Legs* brown, with pale markings, pale knee-spots on femora, most pronounced on hind legs, a short line of pale scaling on anterior surface of mid-femur at base, a thin white line dorsally on anterior surface of hind pair, pale markings at tips of tibiae of fore and mid-legs, a pale apical ring on hind tibia continued over the joint between this and segment 1 of hind tarsi, similar apical and basal rings over the joints between segments of hind tarsi, segment 5 being entirely pale, tarsi of fore and mid-legs brown *Abdomen*. lateral apical white patches on tergites, median apical white patches on I-III, a wide apical white band on V, no apical white bands or median patches on IV, VI, or VII

♂ —Markings as in ♀, segment 1 of hind tarsi (fig 17, d) unusually short, with a double row of stiff curved hairs beneath, segment 1 of fore tarsi $\frac{1}{2}$ length of tibia *Hypopygium* style (fig. 18, e) very similar to that of *U. alboannulata*

LARVA and PUPA. —Unknown

DISTRIBUTION —Known only from type-locality.

21 *Uranotænia edwardsi* Barraud, 1926*.

Ind Journ Med Res xiv, p 337 (♀) TYPE-LOC Golaghat, Sibsagar dist, Assam, 23. ix 1924, caught in bungalow (♀) (Barraud); Gauhati, Kamrup dist, Assam, 31 x 1926 (♂) (Barraud) TYPE: ♀ and allotype ♂ in Brit. Mus

ADULT.—A small, dark mosquito, with mainly white-scaled wings, lines of bluish-white scales in front of wing-root and across pleuræ; fairly easily recognized by diagnostic points given in key The ♂ has remarkable proboscis and mid-tarsi (fig. 17, e, f) Wing about 2 mm

♀.—*Head*. vertex covered with flat bluish-white scales, some long white scales projecting forward between eyes, scales towards nape brownish, very few upright scales, a patch of dark brown scales low down at each side, palpi projecting very slightly in front of clypeus, covered with dark scales and hairs, proboscis dark scaled, a little longer than abdomen, slightly enlarged at tip, approximate length 1.5 mm *Thorax*: mesonotum dark brown, covered with dark brown narrow scales and fairly numerous long black bristles, a line of bluish-white flat scales from wing-root to *ppn*, postnotum brownish-black, pleuræ almost black on upper part, lighter towards coxæ, a line of bluish-white flat scales from *apn* across pleuræ to mesepimeron, no spiracular bristle. *Wings*. spotted, white scales at tip of wing including costa, fork-veins, and apical part of 3, a narrower white area in middle of wing, including costa and vein 1, spreading posteriorly on to cross-veins, base of 5 1, and whole of 5.2, further white scaling at base of wing on costa, a small area on 1 removed from base, on base of 4, on more than basal $\frac{1}{2}$ of 5, and less than basal $\frac{1}{2}$ of 6, remaining wing-scales brownish-black. *Legs*. fore and mid-pairs dark brown, undersides of femora paler, an indistinct pale marking at tip of hind tibia on underside, segment 1 of hind tarsi dark brown, indistinct pale markings at bases of 2, 3, and 4, chiefly on underside, 5 entirely pale; segment 1 on fore and hind legs shorter than tibia. *Abdomen*. dorsum dark brown, venter paler, indistinct apical pale markings on tergites III and IV.

♂.—Markings as in ♀, proboscis longer, about 1.9 mm., apical $\frac{1}{2}$ with a number of long hairs projecting from upper

* [This is probably only a variety of the Malayan *U. micans* Leis., which is similarly adorned and has the same structure of proboscis and middle tarsi in ♂, differing chiefly in having broad and complete white rings at bases of segments 2-4 of hind tarsi. Other Oriental (Malayan) species which are somewhat similar are. (1) *navipes* Theo., with a broad band of flat white scales all round margin of mesonotum, hind tarsi broadly white-tipped; (2) *unimaculata* Leis., proboscis unusually long, tarsi dark; (3) *bimaculata* Leis., scales at base of wing dark.]

side and slightly curved downwards (fig 17, f); tarsi of mid-leg specialized, segment 1 about $\frac{1}{3}$ length of tibia and slightly shorter than 2, 4 rather shorter than 5, with several strong spines at its tip, 5 with a pair of very large simple claws apparently articulated at about middle of segment and not at tip (fig 17, e) *Hypopygium*. style comparatively short and very wide, widest at about $\frac{3}{4}$ length from base and tapering sharply from this point to tip.

Early stages unknown

DISTRIBUTION.—Known only from type-localities.

22. *Uranotænia orientalis* Barraud, 1926

Ind Journ Med Res xiv, p 339 (♂, hypop fig) TYPE-LOC : Golaghat, Sibsagar dist, Assam, 1 1925, caught in jungle (Barraud) TYPE. ♂ in Brit Mus

ADULT—Distinct on account of the general pale colour, mainly pale wings, and unbanded legs* Wing 1.5 mm

♂—*Head* creamy or pale brown flat scales on vertex, a few upright scales towards nape, a small patch of dark brown flat scales low down at each side, palpi scarcely longer than clypeus, proboscis dark brown, about length of abdomen, approximate length 1.4 mm.; antennæ slightly shorter than proboscis, plumes brown. *Thorax* dark brown narrow scales covering mesonotum, except for a band of flat creamy scales from wing-roots round front margin forming a continuous and rather broad pale border, bristles on dorsum black, not unusually long nor numerous, scutellar scales brownish-black, postnotum dark brown, integument of pleuræ very pale brownish, some creamy flat scales on *apn* and *ppn*; other details of pleural scaling cannot be given, as the only known specimen is pinned through this area. *Wings* mainly pale scaled, costa dark brown, except at tip of wing, subcosta dark scaled, except at base, some dark scaling along middle part of veins 1 and 2, otherwise wing-scaling is almost entirely pale. *Legs* brown, without paler markings, uniformly yellowish when seen in certain positions, segments 1 and 2 of fore tarsi together equal in length to tibia, segment 1 of hind tarsi shorter than tibia. *Abdomen* creamy scales covering tergite I, pale scaling on II, except for a small dark brown patch on each side at base, dark brown beneath, III with apical pale band, IV nearly all pale, with a few dark scales at base dorsally, V-VII dark brown, except for an apical median pale patch dorsally. *Hypopygium* style (fig. 18, f) short and widened near tip, resembling that of *U atra*

♀ and early stages unknown

DISTRIBUTION —Known only from type-locality.

* [All the other Oriental species with pale wing-tip have the pleuræ at least partly dark and abdomen more extensively dark.]

23 *Uranotænia unguiculata* Edwards, 1913

Journ. & Proc. Asiatic Soc. of Bengal (n.s.), ix, p. 51 (♂) TYPE-
 Loc. Tiberias, Palestine, x 1912 (N. Annandale) TYPE ♂
 in Ind. Mus.

ADULT* —A fairly large and robust species of general dark colour and distinctive ornamentation of bluish-white scales. Wing 2.5–3 mm.

♀.—*Head* vertex and nape covered with brownish-black flat scales and numerous dark upright scales, usually a small number of pale scales in middle of vertex towards nape, a thin line of bluish-white scales to eye-margins widening out slightly at sides, conspicuous white scaling on inner sides of tori, antennæ, clypeus, palpi, and proboscis dark brown, the last paler beneath, palpi exceeding clypeus by about its length. *Thorax* integument of mesonotum dark brown or black, covered with dark brown narrow scales and dark bristles, a line of bluish-white flat scales along lateral margin from wing-root nearly to neck, not forming a complete border round front margin, integument of pleuræ brownish-black, lighter towards coxæ, some black lanceolate scales on *ppn*, flat white scales on *apn* continued as an irregular line across pleuræ on to mesepimeron. *Wings* dark scaled, except for a short line of white scales at base of vein 1, fork-cells rather longer than usual. *Legs* very small pale knee-spots to femora, fore pair dark brown anteriorly, paler posteriorly, a pale longitudinal line along basal $\frac{1}{2}$ of anterior surface of mid-pair, posterior surface pale, except along dorsal edge, hind femur pale along both surfaces for nearly whole length, except along dorsal edge, tibiæ of fore and mid-legs brown, faintly pale at tips, hind tibia brown, with conspicuous white mark at apex and with a short white line dorso-anteriorly at about middle, tarsi of all legs brown. *Abdomen*. tergites dark brown, with basal dull white or creamy lateral patches, sometimes indications of narrow basal pale ochreous bands on dorsum, venter pale brown.

♂ —Claws of fore leg unequal, segment 1 of hind tarsi shorter than tibia, palpi only very slightly longer than clypeus, antennæ plumose, lateral pale markings on tergites are not so well developed as in ♀, and may be absent, VIII pale scaled dorsally (after rotation) *Hypopygium* style (fig. 18, d) moderately long and comparatively wide for whole length, slightly curved.

* Waterston 1918, p. 11; Edwards 1921 c, p. 283; Kirkpatrick 1925, p. 68; Barraud 1926, p. 338, Martini 1930, p. 196

PUPA†.—Respiratory trumpet about 4 times length of width, with transverse folds near base, abd. seg. I with dendritic tuft, lateral hair on IV–VI fairly long, with 5–6 branches, on VII quite small, on VIII long, with about 8 branches, two or three of the sublateral and submedian hairs on IV–VII well developed, with 3–9 branches, paddle with part inside midrib wider than that outside, outer and apical margins serrate, terminal hair single and short

LARVA† (fig. 16, e–h).—Head and antenna dark brown or black, antenna short, with a few minute spicules, a single hair at about middle, preclypeal spines fairly long, median hairs of mouth-brush hooked and minutely serrate apically, frontal hairs as in fig 16, e, *B* and *C* fairly strong. Hairs on thorax fairly well developed, lateral series branched, lateral hairs on abdominal segments I and II fairly stout and long, those on remaining segments rather small, many with a number of branches, comb-plates (fig 18, g) with 8 or 9 teeth along posterior margin, upper and lower teeth smaller than median. Siphon from 0.7–0.85 mm long, 3 or 4 times length of width at base, pecten of about 19 teeth, of usual form, tuft of about 8 branches, its base opposite the most distal pecten-tooth, *osc* 2-branched, *asc* 3- or 4-branched, *lh* of 4 rather long branches, both pairs of anal papillae short, tips pointed, fan of about 10 fairly long branched hairs arising from rudimentary fan-plate.

HABITAT.—Reedy and weedy pools in swampy ground, borrow-pits, less frequently in disused wells, rice-fields, drains between rice-fields, sides of canals. Larvae cannibalistic, at least in captivity.

DISTRIBUTION.—KASHMIR: locality unrecorded, 1922* (*C. A. Gill*); Jhelum River*, near Wular Lake, 13. x. 1923 (*Sinton*); Srinagar* and Gunderbal*, x. 1923 (*Sinton*); Srinagar*, ix. 1929 (*Barraud*).

Known also from TURKESTAN and westwards to ITALY and ALGERIA.

Specimens from Kashmir are usually larger than those from Egypt and Palestine.

24. *Uranotaenia campestris* Leicester, 1908.

Cul. Malaya, p. 213 (♂ & ♀) TYPE-LOC. Malay Peninsula (*Leicester*) TYPE co-types ♂ & ♀ in Brit Mus.

var. *n. zelena* (see p. 69) TYPE-LOC. Pir Pao, Bombay, xii. 1919 (*Christophers*) TYPE: ♂ & ♀ in Brit Mus

ADULT†.—Distinguished by the apically banded abdominal tergites. Wing about 1.8 mm.

† Joyeux 1918, p. 534, Séguy 1924, p. 64; Theodor 1924, pp. 341–5; Kirkpatrick 1925, p. 70, Martin 1930, p. 197; Montschadsky 1930, p. 580

† Edwards 1922 b, p. 91, Barraud 1926, p. 339

♀—*Head*: vertex and nape covered with brownish-black or purplish flat scales, a thin line of bluish-white scales to eye-margins, turning inwards at sides towards *apn*; very few upright scales on nape. Antennæ, clypeus, palpi, and proboscis dark brown, the last indefinitely paler beneath; palpi scarcely longer than clypeus, proboscis about 1.2 mm. long. *Thorax*: integument of mesonotum deep brown, covered with bronzy dark brown scales and dark bristles; a line of broad bluish-white scales from wing-root along lateral margin to *pym*. Postnotum and integument of pleuræ dark brown, usually a lighter area in middle of mesepimeron. Flat bluish-white scales on *apn*, continued as a line across pleuræ to middle of anterior margin of mesepimeron, small spots of similar scales on coxæ. *Wings* dark scaled, except for a line of pale scales along vein 1 from base to nearly level of base of 2. *Legs*. unmodified, dark brown, undersides of femora paler. *Abdomen*. dorsum dark brown, with apical pale median patches on I-IV, which appear dull white, greyish, ochreous, or brownish, according to angle of light; V with complete white apical band continued to sides, no separate lateral pale patches on abdomen, tergites VI and VII completely dark brown; sternites mainly pale-scaled.

♂—Resembles ♀, except for plumose antennæ. *Hypopygium* (fig 18, e): style short and wide for the whole length.

var. n. *zelena*.—Differs from type-form as follows.—Abdomen with separate lateral pale patches on II-IV, usually a few pale scales at apex of VI in middle. Few, if any, pale scales at base of wing in ♀. (Larva of this form at present unknown.)

PUPA and EGG.—Unknown.

LARVA of type-form (Ceylon) †.—The following points can be made out from figures given by Senior-White (1927).—Frontal hairs *B* and *C* stout leaf-like barbed bristles, *A* with about 4 fine branched, *d* with 3, and *e* with 2; about 9 pointed comb-teeth (no lateral plate shown), about 11 pecten-teeth, of usual form, siphon-tuft with about 8 branches, its base close to the most distal pecten-tooth; *lh* with 5 branches, anal papillæ fairly long, narrow, and pointed.

HABITAT.—Streams and rock-springs (*Senior-White*).

DISTRIBUTION.—Type-form:—INDIA. Pusa*, Bihar, 20. ix. 1912 (*Sharma*), and 17. x. 1922 (*Shaffi*), Delhi*, 24. viii. 1927 (*Senior-White*); Saharanpur*, U.P., 25. ix. 1927 (*Sinton*). ASSAM: Golaghat*, Sibsagar dist., xii. 1924, ii., xi., & xii. 1925 (*Barrand*). CEYLON: Suduganga*, Matale dist., 26. ii. 1920 (*Senior-White*).

† Senior-White 1920, p. 319, and 1927, p. 66.

var *zelena* —BOMBAY. Pir Pao *, as above. COORG, Somwarpet *, vi 1927 (*J D Baily*) N. KANARA DIST. . Karwar, 1901-5 (*Cogill*).

Known also from MALAY PENINSULA

25 *Uranotænia macfarlanei* Edwards, 1914.

Bull Ent Res v, p 127 (♀) TYPE-LOC Hongkong (*Macfarlane*) TYPE. ♀ in Brit Mus

ADULT †—Distinguished by abdominal markings and absence of blue tint in thoracic scales ‡ Wing 2-2.5 mm

♀ —*Head* : a narrow border of white scales to eyes, widening out at sides, remainder of vertex and nape covered with brownish-black, or purplish-black, flat scales, very few upright scales on nape. Tori brown, darker on inner sides, flagellum, clypeus, palpi, and proboscis dark brown, underside of last indistinctly pale, length of proboscis about 1.6 mm, palpi projecting slightly in front of clypeus. *Thorax* : integument of mesonotum deep brown, fairly densely covered with a mixture of dark and pale brown and ochreous narrow scales, a line of lanceolate white scales along lateral margin from wing-root, not continued beyond level of anterior spiracle, scales anterior to this ochreous, a moderate number of fairly long dark bristles, postnotum dark brown, integument of pleuræ black, an indistinct greyish area on mesepimeron; a line of brownish-white scales on *apn*, continued across pleuræ to middle of anterior margin of mesepimeron, small white spots on coxæ. *Wings* : dark scaled, except for a line of pale scales along about basal $\frac{1}{4}$ of vein 1. *Legs* : unmodified, dark brown, undersides of femora paler, tips of femora and tibiae indistinctly pale. *Abdomen* : dorsum dark brown, with well-defined apical white, or creamy, bands on tergites I and V, continued to sides; median apical markings of same colour and small separate lateral white patches on II-IV, VI with a small median pale patch and small lateral pale markings, VII dark brown, venter hidden.

♂ —Markings as in ♀, except that, in some specimens, there is a small area of pale scales on vertex. *Hypopygium* : style short, about 4 times as long as wide, of fairly even width throughout.

LARVA —Has not been isolated in India, but has been described by Brug (from Java) § as follows. —“Head rather broad, with the usual two pairs of thick spiny hairs. Mental plate a little but broader than long, bearing 13 teeth. Antenna

† Edwards 1922 *d.*, p. 480

‡ The Malayan *U. argyrotarsis* Leic is rather similar, but has the hind tarsi broadly white-tipped

§ Brug 1932, p 74

with a simple short hair near its base, terminal bristles shorter than half the antenna. Comb on the eighth segment consisting of about 10 smooth, pointed teeth. Siphon cylindrical, index 4, pecten-teeth about 14, densely fringed, siphon-tuft beyond the middle with about 9 branches. Inner dorsal hair of anal segment three-branched, outer one two-branched, lateral hair moderately long, six-branched. Gills shorter than anal segment. Described from three larval skins and two larvae, found in a small pool with dirty water (rotten leaves) near Dago (Preanger Regentschappen, Java) at about 900 m. above sea-level."

DISTRIBUTION.—ASSAM Golaghat*, Silsagar dist., XII. 1924, 1 & 11 1925, caught in jungle (*Barraud*).

Known also from MALAY PENINSULA, SUMATRA, JAVA, and HONG KONG.

26 *Uranotaenia longirostris* Leicester, 1908

Cul Malaya, p. 217 (♂) TYPE-LOC. Malay Penin (*Leicester*).
TYPE ♂ in Brit Mus

ADULT.—A small blackish species recognized by the unusually long proboscis, and by the single white mark on the abdomen. Wing about 1.5 mm.

♀—*Head* mainly covered with flat bluish-white scales, some towards nape darker, appearing purplish-bronze, very few upright scales, antennæ, clypeus, palpi, and proboscis brownish-black, palpi scarcely longer than clypeus, proboscis longer than whole body, approximate length 2 mm. *Thorax* mesonotum brownish-black, scales narrow and dark brown, a line of broad bluish-white scales from wing-root, along lateral margin, to *pym*, bristles dark brown, fairly numerous and long, brownish-black flat scales on scutellar lobes, post-notum almost black, integument of pleuræ brownish-black, a line of bluish-white flat scales on *apn* continued across pleuræ, no spiracular bristle. *Wings* dark scaled. *Legs*: rather long, unmodified, dark brown, undersides of femora conspicuously white. *Abdomen*: black, tergite IV with large lateral white patches, which may meet on dorsum to form a narrow apical band.

♂—Markings as in ♀, antennæ moderately plumose, hairs very fine, tarsal claws of fore leg rather broad. *Hypopygium*: style short and wide, with a moderate appendage.

PUPA and Egg.—Unknown.

LARVA (3rd stage).—Preclypeal spines short, stout, bluntly pointed; frontal hairs *B* and *C* very stout leaf-like bristles, finely barbed; hairs *A* and *d* small, each of 2 fine branches, hairs *A*, *B*, and *d* in transverse line; *C* posterior and slightly internal to *B*; *e* posterior and external to *C*, very fine.

and split into 2. Lateral hairs on thorax and on segments I and II of abdomen fairly strong, majority branched, a few single, hairs on remaining segments very small and fine, no definite lateral plates on VIII, comb of 5 or 6 pointed teeth, one in middle larger than others. Siphon about 0.32 mm. long (not including valves, which are comparatively very large), only slightly longer than anal segment, and of about even width throughout, pecten of about 7 comparatively large teeth occupying basal $\frac{1}{2}$ of siphon, tuft at about middle of tube, slightly nearer base than most basal pecten-tooth, of about 8 fine branches. Both *asc* and *osc* of 2 branches, anal fan very small, a few hairs arising close together, no definite fan-plate.

HABITAT.—Ponds (*Leicester*), clear pools at side of stream (*Chowdhury*).

DISTRIBUTION.—ASSAM. Hathikuli Tea Estate*, Sibsagar dist., vii 1930 (*K. L. Chowdhury*).

Originally described from the MALAY PENINSULA.

27. *Uranotaenia atra* Theobald, 1905.

Ann. Mus. Nat. Hung. iii, p. 114 (♀). TYPE-LOC. Muna, New Guinea, 1900 (*Biró*). TYPE. ♀ in Nat. Mus. Hung., Budapest.

Uranotaenia lateralis Ludlow, 1905, Can. Ent. xxxvii, p. 385 (♀, *U. ceruleocephala* var. *lateralis*). TYPE-LOC. Mindanão, Philippine Is., 25 vi 1905 (*Vedder*). TYPE. 4 co-type ♀♀ in US Nat. Mus.

Uranotaenia cancer Leicester, 1908, Cul. Malaya, p. 215 (♂ & ♀). TYPE-LOC. Port Swettenham and Klang, Malay Penin. (*Leicester*). TYPE. ♂ & ♀ in Brit. Mus.

Uranotaenia ceylonica Theobald, 1910, M. C. v, p. 503 (♀, figs. wing). TYPE-LOC. Galle, Ceylon, 10. iv. 1907 (*T. B. Fletcher*). TYPE. ♀ in Brit. Mus.

ADULT †.—A small dark brown species, with bright blue scales on head and pleurae, a line of flat scales in front of wing, and lateral pale spots on abdomen; ♂ with modified fore and mid-tarsi ‡. Wing about 1.6 mm.


♀—**Head**.—a border of bright blue or bluish-white flat scales to eye-margins, widest in middle, behind this there are dark bluish or bronzy flat scales and a few upright scales; tori, clypeus, and palpi light brown, proboscis and flagellum of antenna darker brown, palpi only a very little longer than clypeus. **Thorax**.—integument of mesonotum brown, thickly covered with narrow dark brown scales, a number of rather

† Theobald 1907, p. 563, Edwards 1913 b, p. 238; 1922 d, p. 460 (syn.), 1924, p. 357, Barraud 1926, p. 340.

‡ [*U. subnormalis* Mart (*rogersi* Edw.), of Borneo and Singapore, is rather similar, but hind (not front) tarsi of ♂ modified.]

long dark bristles, a short line of bright blue flat scales in front of wing-root, scutellar scales flat and dark brown, postnotum dark brown, integument of pleuræ dark brown on upper part, lighter near coxæ and on lower part of mesepimeron, some flat blue scales on *apn* and a large patch of similar scales in middle of sternopleura, not forming a line across pleuræ. *Wings*: dark scaled, anterior fork-cell very short. *Legs*: dark brown, segment 1 of fore tarsi $\frac{2}{3}$ length of tibia, that of hind leg a little shorter than tibia. *Abdomen*: dorsum brownish-black, with small apical lateral white patches, venter mainly pale scaled.

Fig. 17



Secondary sexual characters of ♂ *Uranotenias*. a, b, c, *atra*, base of front tarsus, tip of middle tarsus, tip of hind tibia; d, *christophersi*, base of hind tarsus; e, f, *edwardsi*, tip of middle tarsus, proboscis.

♂—Segment 1 of fore tarsi only $\frac{1}{2}$ length of 2 and a little more than $\frac{1}{4}$ length of tibia, thickened at base and tip where there are long hairs (fig. 17, a), tibia of mid-leg longer than all the tarsal segments together, segments 4 and 5 of tarsi modified (fig. 17, b), hind tibia with a few peculiar scales near tip (fig. 17, c); markings as in ♀, antennæ plumose. *Hypopygium*: style (fig. 18, g) rather short and widened on apical $\frac{1}{4}$.

PUPA and Egg.—Unknown.

LARVA (fig. 16, a-d) †.—Preclypeal spines short and stout, tips pointed, frontal hairs as figured, *B* and *C* moderately stout. Hairs on thorax and segments I and II of abdomen moderately developed, those on III-VII fine and small, comb-plates large, with 6-9 pointed teeth, finely fringed laterally, on posterior border in a row. Siphon about 0.44 mm. long, index rather less than 3, pecten of 9-12 teeth, occupying less than basal $\frac{1}{2}$ of siphon, teeth transparent, enlarged towards apex, densely fringed, hair-tuft of about 12 fine branches, base slightly beyond most distal pecten-tooth. Both *asc* and *osc* of 2 branches, *lh* of 6-8 fine short branches.

HABITAT.—Crab-holes, stagnant pools, or swamps (*Leicester*), brackish water on coral islet (*Brug*)

DISTRIBUTION.—BOMBAY. Pir Pao*, xu 1919 (*Christophers*). ANDAMAN IS*, ix. 1911, caught from crab holes (*Christophers*). Ceylon Colombo*, 1913 (*James*), Galle, 10 iv 1907 (*T B Fletcher*)

Range extends through MALAY PENINSULA and ARCHIPELAGO and SIAM to NEW GUINEA, AUSTRALIA, and the PHILIPPINES

28 *Uranotaenia testacea* Theobald, 1905.

Ann. Mus. Nat. Hung. iii, p 113 (♀, pls u and m, figs. wing and wing-scales) TYPE-LOC Singapore, 1902 (*Biro*) TYPE. ♀ in Nat Mus Hung, Buda Pest

Uranotaenia falcipes Banks, 1906, Phil. Journ. Sci. i, p 1004 (♂ & ♀) TYPE-LOC. Rizal, P.I., 22 ii 1906 (*Banks*). TYPE: ♂ & ♀ in Bureau of Sci, Manila

Uranotaenia unilineata Leicester, 1908, Cul. Malaya, p. 220 (♂ & ♀). TYPE-LOC Malay Penin (*Leicester*) TYPE. ♂ & ♀ in Brit. Mus.

ADULT ‡.—A medium-sized dark species with blue scales on head and pleuræ, but none above wing-root. Wing 2.4-2.7 mm.

♀.—*Head*: upper surface covered with bright blue or bluish-silvery flat scales, a darker median patch on vertex, a few small dark upright scales on vertex and nape; tori and clypeus pale brown, former darker on inner sides, flagellum, palpi, and proboscis dark brown, palpi only slightly longer than clypeus, proboscis about 1.7 mm. long. *Thorax*: integument of mesonotum bright brown, with a thin covering of brown narrow scales, a moderate number of dark bristles, no flat scales along lateral margin, postnotum light brown, a line of flat bluish-silvery scales from head across pleuræ to anterior border of mesepimeron, including *apn*, integument

† Brug 1924 a, p. 441

‡ Theobald 1907, p 560, Edwards 1913 b, p. 239 (syn.); Barraud 1926, p 342

dark above and below this line, also upper part of mesepimeron, remainder pale brown, no spiracular bristle. *Wings*: dark scaled. *Legs* dark brown, femora paler ventrally, apical $\frac{2}{3}$ of segment 3 and whole of 4 and 5 of hind tarsi creamy white, sometimes a pale area beneath tibio-tarsal joint, tarsi of fore and mid-legs appear very pale in some positions, but are not definitely pale-scaled, segment 1 of hind tarsi a little longer than tibia. *Abdomen* dark brown dorsally, paler ventrally.

♂—Markings as in ♀, antennæ plumose [Legs unmodified, except that the middle femora have a few long, fine, erect hairs about the middle, such hairs not occurring in related species] *Hypopygium*: style (fig. 18, h) of moderate length and width, appendage unusually long and stout.

LARVA, PUPA, and EGG.—Unknown

DISTRIBUTION.—LOWER BURMA: base of Dawna Hills*, 4. iii. 1908 (Annandale). ASSAM: Gauhati*, Kamrup dist., x 1920 (Senior-White), Nongpoh*, Khasi Hills dist., 14 xi 1921 (Christophers), Golaghat*, Sibsagar dist., 1. & 11 1925 (Barraud), S INDIA: Mudigere*, Mysore, 1931 (Anantasaiah Rao)

Also known from the MALAY PENINSULA and PHILIPPINES

29 *Uranotænia annandalei* Barraud, 1926.

Ind Journ Med Res xiv, p 343 (♂ & ♀, ♂ hypop fig) TYPE-LOC.: Golaghat, Sibsagar dist., Assam, xii 1924 and 1. 1925, caught in jungle (Barraud). TYPE: ♂ & ♀ in Brit Mus.

ADULT ♀.—Very similar to *U. testacea*, differing as follows.—*Head*: a border of bluish-silvery flat scales to eyes, brownish or dark bluish flat scales over remainder of upper surface; antennæ, clypeus, palpi, and proboscis dark brown. *Thorax*: mesonotum darker, covered with dark brown narrow scales, a small patch of greyish-brown rather broad scales over each wing-base, not forming a line along lateral margin. *Legs* tarsi entirely brown, segment 1 of hind tarsi same length as tibia.

♂.—Segment 1 of hind tarsi shorter than tibia. [Middle femora without long hairs in middle.] *Hypopygium*: style (fig. 18, l) very short, wide on basal $\frac{1}{2}$, much narrower towards tip.

PUPA and EGG.—Unknown

[LARVA (4th stage) †.—Has not been isolated in India, the following description being based on specimens from

† Edwards 1932, p. 559; ascribed doubtfully to *U. testacea*, but fresh material received from Dr. R. B. Jackson has established its identity with *annandalei*.

Hong Kong Head as in fig 16, i. Antennæ differing from those of all other species of which the larva is known in bearing three large leaf-like plates, representing modified apical bristles, one being retracted nearly to middle of antenna and borne on a thumb-like projection, shaft-hair subapical in position. Preclypeal spines long and not very stout, with the usual emargination between them. Frontal hair *A* slender and single, *B* and *C* both strong, black, and much flattened, *d* and *e*, both short and single. Abdomen practically bare dorsally, with well-developed lateral hairs only on the first three segments. Comb-plates connected by a dorsal saddle which is not much narrower than the plates; comb-teeth sharp-pointed, about 4 in number. Siphonal index about 2.5, large tuft near middle, with 8-10 somewhat flattened, narrowly lanceolate branches. Pecten-teeth about 10 in number, widened at tip and fringed on distal margin only. Anal segment ringed, lateral hair small and simple, about 6 bifid or trifid hairs in fan.

HABITAT.—Shady pool in bed of a partly dried stream.]

DISTRIBUTION.—ASSAM Golaghat*, Sibsagar dist., XII. 1924 and i 1925 (*Barraud*). LOWER BURMA: base of Dawna Hills*, 4. III. 1908 (*Annandale*).

Also found in HONG KONG.

30. *Uranotænia nivipleura* Leicester, 1908.

Cul. Malaya, p. 219 (♂ & ♀). TYPE-LOC.: The Gap, Selangor, Malay Penn. (*Leicester*) (♂), and Singapore, bred from larva found in pitcher-plant (*Dr. Finlayson*) (♂). TYPE. ♂ & ♀ in Brit. Mus.

ADULT †.—A fairly large brown species, with a continuous narrow pale border to mesonotum, pleuræ conspicuously lighter than dorsum of thorax and abdomen. Wing about 3 mm.

♀.—*Head* a line of pale brown scales to eye-margins, dark brown flat scales and numerous upright fork-scales on vertex, latter appearing pale brown in side view; antennæ, clypeus, palpi, and proboscis dark brown, palpi about twice length of clypeus, proboscis a little longer than abdomen, approximate length 1.8 mm. *Thorax*: integument of mesonotum dark brown, covered with brown scales and fairly numerous long curved bristles, a thin line of narrow white scales along margin of mesonotum from wing-roots continued round front, more or less interrupted by a pair of submedian bare lines running back from front margin, postnotum dark brown; pleuræ uniformly pale yellowish-brown, a few

† *Barraud* 1926, p. 343

pale flat scales on sternopleura and *ppn*. *Wings* dark scaled. *Legs* rather long but unmodified, dark brown, femora paler beneath, segment 1 of hind tarsi a little shorter than tibia. *Abdomen* dorsum dark brown, venter pale brown.

♂—Resembles ♀ except for plumose antennæ. *Hypopygium*: style tapering from base to apex, resembling that of *U. recondita*.

LARVA, PUPA, and EGG.—Unknown

HABITAT —Pitcher-plants (Singapore, *Dr Finlayson*); tree-holes (Muktesar, *Fletcher*).

DISTRIBUTION —W HIMALAYAS. Muktesar*, Kumaon, 7,500', ix 1922 (*T B Fletcher*); Bhowali, Naini Tal dist., 4,600', vi–vii 1925 (*S Sundar Rao*).

Known also from MALAYA

31. *Uranotænia maxima* Leicester, 1908

Cul Malaya, p. 221 (♀). TYPE-LOC. The Gap, Selangor, Malay Penin (*Leicester*) TYPE: ♀ in Brit Mus

ADULT †—A large brown species, distinguished from *U. nivicpleura* by presence of dark areas on pleuræ and narrow bands ‡ on some abdominal segments. Wing 3.2–3.7 mm.

♀—*Head* a narrow border of dull white broad scales to eyes, vertex and nape mainly dark scaled and with numerous black upright scales extending forwards nearly to eye-margins, a few scales on nape in some specimens, antennæ, clypeus, palpi, and proboscis dark brown, palpi exceeding clypeus by about its length, proboscis about 2 mm. long. *Thorax* integument, scales, and bristles of mesonotum dark brown, the last moderately numerous, long, and stout, a continuous border of dull white narrow and lanceolate scales from wing-roots along lateral and front margins, more or less interrupted by a pair of submedian bare lines running back from the front, postnotum dark bronzy brown; pleuræ dark brown on upper part bordering mesonotum, including *ppn*, a light transverse band below this, followed by another dark band, remainder light brown, except for indefinite dark patches between coxæ, two patches of creamy flat scales on sternopleura, dull white narrow scales on *apn*. *Wings*: dark scaled, forks longer than usual. *Legs*: dark brown, femora conspicuously pale ventrally, segment 1 of hind tarsi a little shorter than tibia. *Abdomen*: scales on dorsum brownish-black, indistinct pale basal bands, not always pronounced, but usually definite on one or more of the terminal segments.

† Barraud 1926, p. 347, Edwards 1923, p. 2.

‡ As these bands are not always pronounced, and may be overlooked, this species is brought down in two places in the key (p. 61).

♂.—Very similar to ♀, antennæ moderately plumose *Hypopygium*: style (fig 18, j) large, swollen on apical $\frac{1}{2}$ and tapering sharply to a point, fairly numerous hairs towards apex, a moderately long terminal appendage

PUPA and EGG.—Unknown

LARVA (fig 16, m-p).—Distinct from that of any other known Indian species by presence of numerous stellate hairs on thorax and abdomen, the branches of which are stout, with apparently blunt tips, but actually terminating in several minute sharp points. Frontal hairs as figured, *C* short and very stout, resembling a leaflet, *B* single and fairly long. Comb of about 11 fairly large sharp teeth, minutely fringed along sides, comb-plates large, but not connected dorsally. Siphon about 1 mm long, not including valves, index about 3.3, pecten of about 16 teeth of usual form, *osc* of 2 long branches, *isc* divided into 3, *lh* 3-branched, anal papillæ long and slender, with pointed tips, longest fan-hairs about length of papillæ, no distinct bars to fan-plate

HABITAT.—Rock-pools at edge of a stream (Malaya, *Hacker*).

DISTRIBUTION.—ASSAM: Shillong *, Khasi Hills, 4,500', ix. 1917 and 20. ix. 1918 (*T. B. Fletcher*). DARJEELING DIST.: Sureil *, 5,000', x. 1922 (*Barraud*)

Originally recorded from the MALAY PENINSULA.

32 *Uranotænia bimaculata* Leicester, 1908.

Cul. Malaya, p. 226 (♀) TYPE-LOC. The Gap, Selangor, Malay Penn. (*Leicester*) TYPE. ♀ in Brit Mus

ADULT †.—A brown species of moderate size, distinguished from all others in India by the presence of an oval velvety black spot in front of each wing-base. Wing 2.6–3 mm.

♀.—*Head*: some flat creamy scales at each side continued along eye-margins to middle point as a very narrow border, dark brown scales and numerous light-coloured upright scales over remainder; antennæ, clypeus, palpi, and proboscis dark brown, palpi rather more than twice length of clypeus, proboscis about 1.6 mm long. *Thorax*: integument of mesonotum dark brown, fairly thickly covered with brown narrow scales, a moderate number of fairly long dark bristles, a large oval velvety black spot immediately in front of each wing-base, some inconspicuous dull white narrow scales on front margin, integument of pleuræ dark greyish-brown, coxæ and *apm* lighter, a patch of dull creamy scales on sternopleura. *Wings* dark scaled, in some specimens a few scales at base of vein 1 appear pale, *af* only a little shorter than *pf*. *Legs*: unmodified, dark brown, undersides of

† Edwards 1921 c, p. 283; Barraud 1926, p. 346, Martini 1930, p. 108.


femora paler. *Abdomen*: brownish-black dorsally, paler ventrally.

♂—Resembles ♀, but antennae densely plumose, upright scales on head smaller. *Hypopygium* (figs 18, a): style moderately long, wide on basal $\frac{1}{2}$, apical part recurved and tapering slightly to tip.

PUPA, LARVA, and EGG.—Unknown.

HABITAT—Tree-holes and bamboos (*Barraud*).

Fig 18



a, ♂ hypopygium of *Uranotaenia bimaculata*; b-m, ♂ style of other species of *Uranotaenia*. b, *alboannulata*; c, *christophersi*; d, *unguiculata*; e, *campestris*; f, *orientalis*; g, *atra*; h, *testacea*; i, *annandalei*; j, *maxima*; k, *stricklandi*; l, *recondita*; m, *luteola*.

DISTRIBUTION.—DARJEELING DIST.: Kurseong*, Sureil*, and Mungpoo*, 5-6,000', ix. & x. 1922 (*Barraud*); Marian barrie Tea Estate*, near Sukna, c. 500', viii. 1928 (*Sobha Ram*).

Also known from the MALAY PENINSULA and JAPAN.

33. *Uranotænia luteola*, sp. n. †.

TYPE-LOC Malabar Coast, x 1915 (*Khazan Chand*) TYPE: ♂ & ♀ in Brit. Mus

ADULT.—Recognized by the uniformly pale pleuræ and banded abdomen and absence of flat or pale scales on lateral margin of mesonotum. Wing about 2.7 mm

♀.—*Head*. mainly covered with dark brown flat scales and fairly numerous upright scales, latter extending forwards nearly to eye-margins, scales along eye-margins appear lighter, tori and clypeus brown, palpi and proboscis dark brown, palpi about twice length of clypeus, proboscis about 1.6 mm long. *Thorax* integument of mesonotum chestnut-brown, scales brown and scanty, bristles black, no flat or pale scales along lateral margin, postnotum deep brown, contrasting with uniform very light brown colour of pleuræ. *Wings* dark scaled. *Legs*: dark brown, undersides of femora not conspicuously pale, segment 1 of hind tarsi longer than tibia, segment 4 more than 3 times length of 5. *Abdomen*. dorsum dark brown, with well-defined narrow yellowish basal bands on tergites II–VII, all of about equal width, sternites covered with yellowish-brown scales

♂.—Very similar to ♀, but antennæ plumose *Hypopygium* style (fig 18, m) unusually long and slender

LARVA, PUPA, and EGG.—Unknown.

DISTRIBUTION.—ANDAMAN ISLANDS *, ix 1911 (*Christophers*) MALABAR COAST *, x. 1915 (*Khazan Chand*)

34. *Uranotænia stricklandi* Barraud, 1926.

Ind. Journ. Med. Res. xiv, p. 345 (♂ & ♀, ♂ hypop. fig.). TYPE-LOC Balasum River, near Kurseong, Darjeeling dist., vi. 1923 (♂) (*O. Strickland*), Nilgiri Hills, S India, x. 1915 (allotype ♀) (*Khazan Chand*) TYPE ♂ & ♀ in Brit Mus

ADULT.—A brown species of moderate size with well-marked yellowish bands on abdomen and conspicuous dark markings

† [This was described by the author (1926, p. 344) as *U. lutescens* Leio., but, owing to certain discrepancies from Leicester's description, he came to doubt the identity of the Indian form and asked me to compare a pair of specimens from the Malabar Coast with Leicester's types. I have done so, and find a number of distinctions which certainly indicate that two species are involved. *U. lutescens* (of Malaya and Borneo) differs from *U. luteola* as follows.—Close-lying scales of head pale (Leicester said light bronze or fawn-brown), upright forked scales confined to one row of at most a dozen on the nape. Antennæ of ♂ rather more densely plumose. Mesonotum (including postnotum) pale yellowish-brown, no darker than pleuræ. Whitish basal bands on abd. seg V and VI more distinct than on other segments, that on VI in ♂ occupying $\frac{1}{2}$ the tergite or more, those on II–V in ♀ often interrupted in middle, style of ♂ not quite so slender. Leicester's species was bred from bamboos.—F. W. EDWARDS.]

on pleuræ; differs from *U. bicolor* Leic in pleural markings and in character of mesonotal scaling. Wing 2.2-2.3 mm

♀—*Head* vertex covered with dark brown flat scales, a narrow border of greyish-white or pale ochreous broad scales along eye-margins and a few similar scales in middle of vertex, very numerous upright pale scales scattered over vertex and nape, continued anteriorly nearly to eye-margins, torus of antenna pale brown, flagellum dark brown, clypeus, palpi, and proboscis dark brown, palpi about twice length of clypeus, proboscis about 1.5 mm long. *Thorax*. integument of mesonotum pale at sides, dark brown in middle, darkest over wing-roots, a scanty covering of brown or golden scales and a moderate number of dark bristles, no line of flat scales in front of wing-root, postnotum brownish-black, pleuræ yellowish-white, marked with conspicuous brownish-black patches, *apn* and *ppn* dark, a dark area below anterior spiracle extending on to upper part of sternopleura, another dark area in middle of this sclerite, larger part of mesepimeron dark, some broad pale scales between the two dark areas on sternopleura. *Wings*: dark scaled. *Legs* unmodified, dark brown, femora paler beneath. *Abdomen*. dorsum dark brown, with well-marked basal ochreous basal bands, venter pale

♂—Markings as in ♀, antennæ plumose, segment 1 of hind tarsi a little longer than tibia. *Hypopygium*: style (fig 18, k) long and moderately slender, but shorter than that of *U. luteola*

LARVA, PUPA, and EGG—Unknown

DISTRIBUTION—Type-localities mentioned above, also Mercara *, Coorg, S. India, iv. 1928 (*Abdul Majid*).

35. *Uranotænia hebes* Barraud, 1931.

Ind. Journ. Med Res xix, p. 609 (♀) TYPE-LOC.: Nongpoh, Khasi Hills dist., Assam, iii. 1927, caught in jungle (*Barraud*)

TYPE: one ec-type ♀ in Brit Mus, one in M S I coll., Kasauli.

ADULT.—A brown species of moderate size, resembling *U. maculipleura* in general appearance, but differing in having only one small dark area on pleuræ, some pale scales in front of wing-root, shorter palpi, and much less expanded upright scales on head. Wing about 3 mm

♀.—*Head*: covered with broad flat dark brown scales, those along eye-margins appear pale when viewed in certain positions, a moderate number of upright scales scattered over dorsal surface extending forwards near to eye-margins, the ends of these scales very slightly expanded, torus pale brown, darker on inner side, clypeus brown, palpi

DIPT.—VOL. V.

G

quite short, projecting only very slightly in front of clypeus, proboscis dark brown *Thorax*: integument of mesonotum rather dark brown, covered with narrow brown scales and dark bristles, some lanceolate pale brown scales over wing-root and a small collection of white or creamy lanceolate scales along margin in front of wing-root, some of which are fairly broad, but no line of flat scales in this position, postnotum dark brown, integument of pleuræ pale brown, a small dark area behind anterior spiracle, white lanceolate scales with silvery sheen covering *apn*, some broader pale or silvery scales on *ppn*, sternopleura, and mesepimeron, no spiracular bristle. *Wings* scales dark brown, forks of about equal length *Legs* dark brown, femora paler beneath, segment 1 of fore tarsi a little longer than last four together, segment 1 of hind tarsi very little shorter than tibia *Abdomen*: dorsum very dark brown, without markings, venter pale brown.

♂ and early stages unknown

DISTRIBUTION.—Known only from type-locality

36 *Uranotaenia maculipleura* Leicester, 1908

Cul. Malaya, p. 223 (♀) TYPE-LOC 6 miles from Kuala Lumpur Malay Penin. (*Leicester*) TYPE. ♀ in Brit. Mus.

ADULT.—A medium-sized species of fairly robust build and of general dark brown colour, numerous upright scales on the head with markedly expanded ends. Wing about 2.3 mm.

♀—*Head*: vertex and nape covered with dark brown flat scales and numerous upright scales extending forwards nearly to eye-margins and with markedly expanded ends, scales along eye-margins appear pale; antennæ, clypeus, palpi, and proboscis dark brown, palpi twice length of clypeus or slightly more *Thorax*: integument of mesonotum dark purplish-brown, scales narrow and brown, bristles dark brown, no flat or pale scales along margin, postnotum deep brown; pleuræ pale brown, with dark brown patches, *ppn* dark, a small dark area below anterior spiracle, two small dark patches on sternopleura, mesepimeron entirely dark, a few pale broad scales on sternopleura and on *apn*. *Wings*: dark scaled. *Legs*: dark brown, femora not conspicuously paler, segment 1 of hind tarsi very slightly longer than tibia. *Abdomen*: dorsum brownish-black, venter pale brown

♂ and early stages unknown.

DISTRIBUTION.—DARJEELING DIST.: Marianbarrie Tea Estate*, near Sukna, c. 500', viii. 1928, caught in jungle (*Sobha Ram*)

Originally recorded from MALAY PENINSULA

37. *Uranotænia bicolor* Leicester, 1908

Cul Malaya, p 225 (♂ & ♀) TYPE-LOC. Kuala Lumpur, Malay Penin (*Leicester*) TYPE. ♂ & ♀ in Brit Mus

Uranotænia fusca Leicester, 1908 (nec Theobald), Cul Malaya, p 227 (♂ & ♀). TYPE-LOC. Sungai Limbing, Malay Penin (*Daniels*)

Uranotænia leicesters Edwards, 1913, Bull Ent Res iv, p 239. Nom nov for *U. fusca* Leic.

ADULT*.—A rather large brown species, distinguished from others with banded abdomen by characters given in key Wing about 3 mm

♀—*Head* vertex and nape covered with dark brown flat scales, and numerous brown upright scales extending well forwards, a narrow border of pale scales to eyes, torus yellowish-brown, flagellum of antenna dark brown, clypeus and palpi brown, proboscis darker, palpi about twice length of clypeus. *Thorax*. integument of mesonotum yellowish-brown, with darker areas, one in front of scutellum and one over each wing-root, in some specimens, however, the mesonotal integument is uniformly rather dark brown, a moderate covering of pale ochreous narrow scales, no line of flat scales or narrow white scales on lateral margins, bristles dark brown; pleuræ brown, lower part of mesepimeron distinctly darker, meron and coxæ pale, some pale flat scales on sternopleura. *Wings*. dark scaled. *Legs*. dark brown, femora lighter beneath towards base. *Abdomen* dark brown, with well-defined pale ochreous or creamy basal bands on tergites I-VI or I-VII

♂.—Resembles ♀, but abdominal pale bands usually broader and antennæ plumose. *Hypopygium*. not examined.

PUPA, LARVA, and EGG.—Unknown

HABITAT.—Marshy edges of jungle-stream (*Leicester*)

DISTRIBUTION.—S INDIA. Mudigere, Mysore, 1931 (*Ananta-swamy Rao*).

Known also from MALAYA

38. *Uranotænia recondita* Edwards, 1922.

Bull. Ent. Res. xii, p. 91 (♂ & ♀). TYPE-LOC. Karwar, N. Kanara Dist., viii. & x. 1902 (*Coghill*). TYPE. ♂ and paratype ♀♀ in Brit Mus

ADULT†.—A rather small obscure brown species, distinguished by the very bristly mesonotum, absence of ornamentation, uniformly pale pleuræ, and presence of fairly numerous longish upright scales on head. Wing 2.3-2.6 mm.

♀.—*Head*: vertex and nape covered with dark brown flat scales, those along eye-margins somewhat lighter, fairly numerous and rather long brown upright scales scattered

* Edwards 1922 d, p. 486

† Barraud 1926, p. 346
G 2

over vertex and nape, tori and palpi brown, flagellum, clypeus, and proboscis dark brown, palpi from 2 to nearly 3 times length of clypeus, slightly variable in specimens from different localities, proboscis about 1.7-1.8 mm long. *Thorax*. integument of mesonotum light brown and often translucent, covered with unusually long and numerous curved dark bristles, scales brown, no flat or narrow white scales on lateral margins, postnotum dark brown, pleuræ uniformly pale brown. *Wings* dark scaled. *Legs* unmodified, dark brown, femora paler beneath. *Abdomen* dorsum deep brown, venter paler.

♂—Resembles ♀, but antennæ plumose. *Hypopygium* style (fig 18, 1) long and moderately wide, tapering slightly to tip.

PUPA and EGG—Unknown.

LARVA (fig 16, j-l)—Peclypeal spines leaf-like, widening out from base, outer side curved, inner straighter, tips bluntly pointed, frontal hairs missing in two damaged skins available, mentum triangular, with 8 or 9 rather blunt teeth on either side of central one. Lateral thoracic hairs well developed, with numerous long branches, some smaller many branched hairs also present, lateral hairs of abd seg I and II stout and long, either single or 2-branched, finely barbed, hairs on following segments fine, some long, comb-plates small, comb of 12-14 teeth, with delicate lateral fringes, tips rounded and also fringed (in side view teeth may appear to end in a single point). Siphon slightly more than 1 mm. long, pecten of 22-26 teeth, with broad fringed tips, tuft of about 8 fairly strong branches, the base slightly beyond the most distal pecten-tooth. Both *isc* and *osc* of 2 branches, *lh* of 3 short branches; papillæ damaged, both pairs apparently of moderate length, fan-hairs missing, fan-plate distinctly barred.

HABITAT—Tree-holes, during the monsoon.

DISTRIBUTION—N. KANARA DIST Karwar, viii. & x 1902 (Coggill), ix. 1921* (Barraud), Kadra*, ix 1921 (Barraud). CENT. PROV Buldana, 15. ii. 1923 (S Sundar Rao). N. MADRAS COAST: Vizagapatam*, xi 1927 (J. D. Barry), Muniguda*, Vizagapatam Agency, 12. ii. 1933 (Senior-White). ORISSA Ranchi*, 10 xi. 1930 (Senior-White).

Not recorded from elsewhere.

39 *Uranotaenia novobscura*, sp. n.

TYPE-LOC Sukma, Darjeeling dist, c 500', x. 1922 (Barraud).

TYPE ♂, no 1276/1881, allotype ♀, and other specimens, from same place, viii 1928 (Sobha Ram).

ADULT.—A small dark brown species without ornamentation, apparently indistinguishable from *U. obscura* Edwards.

in the adult stage, but differing in the larval stage. Wing 2.2-2.6 mm

♀—*Head* vertex and nape covered with dark brown flat scales and a moderate number of upright scales, some scales at sides appear lighter when seen in certain positions, antennæ, clypeus, palpi, and proboscis dark brown; palpi scarcely twice length of clypeus *Thorax* integument of mesonotum and scales dark brown, bristles less dense than in *recondita*, pleuræ pale brown *Wings* dark scaled. *Legs* dark brown, undersides of femora, pale brown, segment 1 of hind tarsi distinctly longer than tibia *Abdomen* both dorsum and venter dark brown, without markings

♂—Resembles ♀, but antennæ plumose *Hypopygium*: style fairly long, 8-9 times length of width in middle

LARVA, 4th stage (described from one damaged skin, from which type ♀ resulted)—*Head* dark brown, lighter at sides *Antenna* very short and rather stout, about 0.14 mm long, shaft dark brown, bare, except for a hair with several fine branches at about $\frac{2}{3}$ from base *Frontal* hair *A* with 4 branches, *d* also with several fine branches, *B* and *C* missing, the latter apparently attached some distance posteriorly to other three pairs *Mentum* longer than width at base, 6-7 regular teeth on either side of median tooth *Thorax* with some chitinated plates or tubercles at bases of tufts, the larger tubercle on metathorax with small thorn-like spines *Abdomen* with long 2-branched hairs on I and II, arising from small plates *Longer* hairs on following segments 3-branched, a long single hair on either side of VII *Comb-plates* large but not united on dorsum to form a saddle (as is the case in *obscura*). *Comb* of 7 fairly large pointed teeth in a row, each tooth fringed at base. *Subsiphonal* tuft of 5 fairly long subplumose branches. *Anal* segment completely ringed by plate, hind margin finely spinose. *Subdorsal* and *fan-hairs* missing; *lateral* hair of 2 strong branches, *fan-plate* with distinct bars; *papillæ* damaged. *Siphon* moderately dark, slightly tapering, length about 0.5 mm., not including valves; width at base, when compressed, about 0.2-0.3 mm. *Acus* fairly large. *Pecten* 17-20 teeth, broad and fringed as usual (not simple as in *obscura*), in a close-set rank, extending beyond middle of tube; *siphonal* tuft 3-branched, arising near most distal pecten-tooth.

HABITAT.—Bamboos (*Barraud*).

DISTRIBUTION.—Known only from type-locality.

Genus **THEOBALDIA** Neveu-Lemaire, 1902.

C R. Soc. Biol. liv, p 1331 Genotype, *Culex annulatus* Schrank.
Pseudotheobaldia Theobald, 1907, MC iv, p 271 Genotype,
P. inverticollata Theo
Allotheobaldia Brolemann, 1919, Ann Soc Ent France, lxxxviii,
 p 91 Genotype, *Culex longiareolatus* Macquart

ADULT*—Mosquitoes resembling large species of *Culex* in general appearance, but distinguished by presence of 5-10 spiracular bristles arranged in a row along anterior margin of anterior spiracle, by presence of hairs on remigium (base of vein 1), especially on underside of wing (fig. 7) †, and by absence of pulvilli. Vertex clothed with curved and upright scales, both narrow, and usually also hair-like false scales, scutellar scales narrow or lanceolate, antennæ in both sexes shorter than proboscis, those of ♂ plumose, palpi of ♀ from $\frac{1}{3}$ — $\frac{1}{2}$ length of proboscis, latter of moderate length, palpi of ♂ widened and spatulate at apices, either slightly longer or a little shorter than proboscis, a few posterior pronotal bristles, a row of spiracular, no post-spiracular. Wings fairly broad, subcosta ending slightly nearer tip of wing than level of base of vein 3, vein 6 somewhat sinuous. Tarsal claws of ♀ all simple, those of fore and mid-legs of ♂ toothed.

The genus is divided into several subgenera, two of which occur in the Indian region, *Theobaldia*, s.str., and *Allotheobaldia*, the latter includes only one species, *T. longiareolata* (Macq.)

♂ *hypopygium* Subgenus *Theobaldia* (fig 19, b, c): coxite fairly long, with small subapical and basal lobes, style of moderate length, with terminal appendage, phallosome divided into lateral plates, paraprocts with teeth at crown; tergite VIII with a few strong spines in middle on apical border. Subgenus *Allotheobaldia* (fig 19, a): coxite without subapical lobe, style with double terminal appendage; paraprocts pointed at crown, no spines on tergite VIII.


* Edwards 1921 c, p. 285; 1924, p. 363, 1932, p. 101; Bami Prashad 1915, p. 501; Christophers 1922, pp. 530-572 (♂); 1923, pp. 698-720 (♂); Barraud 1924 l, p. 139, Kirkpatrick 1925, p. 71; Barraud & Covell 1928, p. 676 (buccal cavity), Martini 1930, p. 200.

† [This character was first discovered by Dyar and Shannon (1924), who stated in their key that the hairs on the underside of the wing at the base are on the first vein (stem-vein, remigium, or base of radius). The statement has been repeated by others, including myself and Barraud, but, as has been pointed out to me by Mr A. J. E. Terzi, it is the base of the subcosta which carries these hairs (as correctly noted by Dyar and Shannon in their *diagnosis* of the genus *Culiseta*); the stem-vein is hidden under the thickened base of the subcosta when the wing is seen from beneath —F. W. EDWARDS]

♀ *hypopygium*. in both subgenera the abdomen is blunt-tipped, VIII not retractile; cerci comparatively small; three chitinated spermathecae

PUPA.—Respiratory trumpet short, with large opening; dendritic tufts present on abdominal segment I, lateral hairs on VII and VIII comparatively small, paddles large and rounded, a single apical hair near termination of midrib.

Fig 19



♂ hypopygial structures of *Theobaldia*: a, *longiareolata*; b, *investiata*; c, *indica* Lettering as on p. 4

LARVA, 4th stage (fig 20)*—A single pair of hair-tufts on siphon at base; pecten and anal fan well developed; other characters given in descriptions below.

DISTRIBUTION and BIOLOGICS.—In the Indian region the genus is confined to the north-west and Himalayas, the adults suck human blood and may at times be troublesome. The eggs are laid in boat-shaped masses on water; the larvae live in ground-pools and sometimes in pools in beds of hill-streams.

* Edwards 1921 c, p. 287; 1932, p. 102.

The species of *Theobaldia* are few in number, but occur throughout the Palearctic and Nearctic regions, and also in Africa, Australia, New Zealand, being absent from the Neotropical and the greater part of the Oriental region.

Key to Adults.

- | | |
|---|-------------------------------|
| 1 Thorax decorated with longitudinal lines of pale scales, anterior edge of costa entirely pale scaled from base to tip of wing | <i>longiareolata</i> , p. 88. |
| Thorax without pale lines, costa either entirely dark or with a sprinkling of pale scales only | 2 |
| 2 Tarsal segments marked with broad basal white rings .. | <i>indica</i> , p. 93 |
| Tarsi entirely dark | <i>niveicincta</i> , p. 91 |

40. *Theobaldia longiareolata* (Macquart), 1838.

Dipt Exot 1, p. 34 (*Culex*) * TYPE-LOC : Canary Is TYPE (if in existence) Paris Museum

Culex spathopalpis Rondani, 1872, Bull Soc Ent Ital iv, p. 31.
TYPE-LOC : Italy TYPE Florence Mus

ADULT † —Wing-length in larger specimens 6.5–7 mm., width 1.5 mm

♀ —*Head* a border of fairly broad white scales to eye-margins, continued downwards between eyes, a double line of similar scales from nape to front of vertex and some similar scales along back of head and low down at each side, remainder covered with black scales, long yellowish hair-like false scales, and long upright scales with slightly expanded ends, the last chiefly on nape. Tori brown, with white scales on upper and inner surfaces, white scales on undersides of first two flagellar segments, remainder, as well as clypeus and proboscis, dark brown. Palpi between $\frac{1}{2}$ and $\frac{1}{4}$ length of proboscis, dark brown, heavily sprinkled with white scales, chiefly on upper surface, tips entirely white. *Thorax* : integument brown, a median line of white scales from front margin to scutellum, dividing either side of ante-scutellar space, another line on each side from front margin along lateral border, curving inwards about mid-way between front border and wing-base, then continued in straight line to lateral lobe of scutellum, another indefinite line of white scales

* [Macquart refers in 'Diptera Exotica' to his description of this species in Webb & Berthelot's 'Hist Nat des Iles Canaries,' ii, pt. 2, Zool., Dipt p. 99. The latter work, however, was not actually published until 1840 (*vide* Wiegmann's Arch 1841, p. 153), consequently the original reference is correctly given as above.]

† Edwards 1917, p. 228, and 1921 c, p. 287 (syn.), 1926 b, p. 408; Waterston 1918, p. 9, Barraud 1924 l, p. 139, Kirkpatrick 1925, p. 72; Ségué 1925 a, p. 79, Martini 1930, p. 214

laterally over wing-root; remainder covered with narrow, curved, light brown scales, bristles on dorsum short, those laterally over wing-roots longer and yellowish, white lanceolate scales and yellowish bristles on scutellar lobes. Postnotum and integument of pleuræ brown, some fairly broad yellowish scales on upper part of *ppn*, white scales on lower part, and also on sternopleura, mesepimeron, prosternum, coxæ, and lower part of *apn*, a number of yellowish bristles on upper part of last, 5 or 6 spiracular bristles and some hairs, a few rather short posterior pronotal, 4 lower mesepimeral and about 20-24 upper mesepimeral, a group of about 12 sternopleural near lower corner of mesepimeron. *Wings* front margin of costa continuously pale scaled from base to tip of vein 2 1, otherwise scales are dark, except for a few pale ones on remigium, a few hairs also on upper side of wing on this part, and more numerous hairs in same position on underside. *Legs* femora mottled or spotted with white or yellowish and brownish-black scales, the pale scales forming almost complete preapical pale rings on fore and hind femora, tibiae dark, fairly evenly spotted with pale scales on anterior surface; segment 1 of tarsi of all legs pale at base and extreme tip, otherwise brownish-black, streaked with pale scaling, segments 2 and 3 mainly dark with basal pale markings, and pale speckling in some specimens, 4 and 5 on fore and mid-legs dark, segment 4 on hind leg with basal pale marking, 5 dark. *Abdomen* tergites with basal white bands, otherwise entirely covered with a mixture of pale yellow and dark brown scales, variable in arrangement, in some specimens the tergites are almost entirely pale, in others the pale scaling is confined to the middle and posterior margin of each tergite; in others again the scaling is mainly dark, except for the basal white bands; these different forms possibly represent seasonal or local variations.

♂—Markings as in ♀, antennæ plumose and barely $\frac{2}{3}$ length of proboscis, palpi a little shorter than proboscis, last segment and apical part of penultimate widened and spatulate, dark brown, with white scaling on upper side at bases of segments 2-5 and on underside at tips of last two segments, otherwise dark brown. *Hypopygium* (fig. 19, a) of very distinctive structure.


PUPA*.—Respiratory horn short, wide, with large opening, apical part covered with network of small folds. One of the sublateral hairs on tergites V and VI very long and single, a similar hair on IV, but 2-branched; all hairs of posterior margin of VII rather short, lateral hair with several short branches; lateral hair on VIII with a number of moderately

* Theodor 1924, p. 341; Kirkpatrick 1925, p. 72.

long branches forming a tuft, another hair internal to this with 2-3 branches. Paddles large and rounded, posterior margin minutely serrate, a single terminal hair slightly internal to, and beyond, termination of midrib.

LARVA (fig 20, a-e) * —Length about 10 mm, including siphon, which is about 1.25-1.5 mm long, head very dark;

Fig 20



Larval structures of *Theobaldia* a-e, *longiareolata*, f-i, *niveitcenata*.
b, i, pecten-teeth, c, g, comb-teeth.

siphon pale brownish. Antenna dark brown, comparatively short, nearly straight and moderately stout, length about 0.32 mm., shaft smooth, with hair-tuft of 2-3 branches arising nearer apex than base. Hairs of mouth-brushes with minute serrations apically. Frontal hairs as figured, A 2-3 branched, B and C single, d single or double. Lateral tufted hairs on thorax well developed and arising from large chitinised tubercles.

* Langeron 1918, p. 98; Séguy 1924, p. 54; Kirkpatrick 1925, p. 75; Martini 1930, p. 215.

Lateral hairs on *abdomen* fairly long, especially on I-III, comb a triangular patch of about 45-50 scale-like fringed teeth, each ending in several sharp points. *Siphon* as figured, chitinisation at base towards posterior margin irregular in outline, no definite acous, pecten of 8-10 fairly large widely-spaced teeth, extending from base nearly to apex, several teeth nearest to base with 1-2 lateral denticles, others simple, usually a few additional small teeth on membrane near base of tube, hair-tuft at base of chitinised part of tube near posterior border, with 15-20 subplumose branches, subsiphonal tuft large, with 10-12 branches. *Anal segment* almost completely enclosed in chitinised ring, some large spines on posterior border towards dorsum, *lh* fine and divided into 3-4 short branches, *osc* of 2-3 long branches, *usc* divided into a number of branches all about same length. *Anal fan* well developed, about 16 hairs arising from fan-plate, each split into a number of fine branches, several branched hairs also arising from membrane between fan-plate and base of segment, dorsal anal papillæ about as long as anal segment, ventral pair shorter, both pairs pointed at tips.

HABITAT.—Pools and ponds

DISTRIBUTION.—Common in the North-West and Western Himalayas as far east as Naini Tal, but not found in other parts of the Indian region. NW FRONTIER: Bannu*, iv-v 1917 (*Sinton*), Malakand*, iv. 1931, Taru* (Peshawar), v. 1916 (*T. B. Fletcher*), Cherat*, Idak*, Dera Ismail Khan*. N. BALUCHISTAN: Quetta*, viii. 1911 (*Davys*), and vi. 1923 (*Browse*); Pishin*, vi. 1929, Chaman*, viii. 1931. SIND: Karachi*. WESTERN HIMALAYAS: Murree*, 1922 (*Gill*), Kasauli* and Sanawar*, v. 1911 & vi. 1912 (*Christophers*), and iv-vi. 1921, 1922, and 1924 (*Barraud*), Naini Tal (U.P.) 7,000' (*Giles*).

Outside India this species occurs in the MEDITERRANEAN REGION, ATLANTIC ISLANDS, PERSIA, EAST and SOUTH AFRICA.

41. *Theobaldia niveltsoniata* (Theobald), 1907

M.C. iv, p. 272 (*Pseudotheobaldia*). TYPE-LOC.. Dehra Dun, U.P., India, u. & iii. (*Thompson*). TYPES: 2 ♂♂ in Brit. Mus.

ADULT †.—Distinguished by the dark legs and broad white scutellar scales †. Wing about 7-8 mm.

♀.—*Head*: dorsal surface with narrow pale yellow scales, dark upright and narrow hair-like false scales, a patch of

† Edwards 1921 o, p. 287, Barraud 1924 l, p. 141, Martini 1930, p. 212

† [The European *T. glaphyoptera*, which is the nearest relative of *T. niveltsoniata*, and with which it was at one time confused, has narrow scutellar scales and a different hypopygium.]

broader yellow scales at each side. Tori dark brown, with yellow scales, chiefly on inner sides, flagellum brown, with pale pubescence and brown hairs. Clypeus and proboscis dark brown. Palpi about $\frac{1}{2}$ length of proboscis, brown, with a plentiful sprinkling of pale scales, tips pale. *Thorax*: integument of mesonotum dark brown, with a sparse covering of golden, narrow, curved scales, bristles fairly numerous but not long, yellow or brown according to the angle of view, scutellar scales white or creamy, fairly broad, bristles long and yellowish. Postnotum and integument of pleurae dark brown, some yellowish, fairly broad scales on upper part of *ppn*, pale broad scales on lower part and on propleura, sternopleura, mesepimeron, and below margin of mesonotum in front of wing-root, pleural bristles pale, about 8 spiracular, 4 or 5 small *ppn*, bristles on propleura, sternopleura, and upper part of mesepimeron fairly numerous and long, 2 or 3 lower mesepimeral. *Wings* veins clothed with long narrow dark scales, membrane darker in patches, chiefly in region of cross-veins and at bases of forks, giving the wing a slightly spotted appearance, c-vs 2-3, 3-4, and 4-5 closely approximated. *Legs* bronzy black, marked with white and pale yellow, all the femora with a pale knee-spot, a subapical pale line and spot on anterior surface of fore pair, sometimes confluent and forming a large pale patch, anterior surface of mid-pair dark, posterior surface of fore and mid-pairs pale from base nearly to tip, hind femur white except for a subapical black ring and a dark line along the dorsal edge. Tibiae dark when viewed from the front, with apical pale ring; pale for whole length when viewed from behind. Tarsi entirely dark, first segment indistinctly paler ventrally at base in some specimens. *Abdomen* tergites brownish-black, with silvery basal bands, in many specimens the bands are distinctly narrower in the middle of each tergite, sternites mainly covered with pale yellow scales and with small lateral basal silvery spots.

♂—Resembles ♀, but antennae plumose; palpi slightly longer than proboscis, dark brown, paler at joints, some long hairs on segments 3 and 4, shorter hairs on 5, which is widened and spatulate. *Hypopygium* (fig 19, b): very similar to that of *T. indica*, differing in shape of *lp* and in having fewer small spines on apical border of 8t.

PUPA.—Unknown.

LARVA (fig 20, f-1)—Length about 9 mm., including siphon, which is 1.3-1.5 mm long. *Antenna* pale brown, about 0.5 mm long, hair-tuft of 4-5 branches, arising at about middle of shaft, latter smooth. Median hairs of mouth-brushes minutely serrate. *Frontal hairs* as figured, *A*, *B*, and *C* all equally developed, each usually with 3-10 moderately

long branches. Lateral tufted hairs on *thorax* well developed and arising from chitinated tubercles. Some fairly long lateral hairs on abdomen. *Comb* of 45-50 scale-like fringed teeth in a triangular patch, subsiphonal tuft large, with 8-12 subplumose branches. *Siphon* about 3 times length of diameter at base, acus fairly large, pecten of 12-16 strong teeth with 3-4 lateral denticles, commencing from base, usually 3-4 smaller teeth in addition close together at base; distal to the teeth a row of about 15 hairs forming a continuation of the pecten and extending nearly to apex of tube; hair-tuft at base, near posterior margin, usually of 8-10 long branches. *Anal segment* enclosed in chitinated ring, without large spines on posterior margin; *lh* fine, with 2-3 branches, *osc* single and long, *isc* divided into a number of branches subequal in length, fan well developed, about 14 branched hairs arising from fan-plate, 2 or 3 branched hairs on segment between plate and base, both pairs of papillæ long and pointed, distinctly longer than anal segment.

HABITAT—Pools in beds of hill-streams

DISTRIBUTION—**WESTERN HIMALAYAS** Kasauli*, common from II-IV (*Christophers & Barraud*); Murree*, 1922 (*Gill*); Theog*, Hindustan-Tibet road, 8,000', 2 v 1907 (*Annamdale*), Nain Tal* (CMB Records), Muktesar*, v 1923 (*T. B. Fletcher*), Dehra Dun (type-loc. as given above). **EASTERN HIMALAYAS** north of Yatung*, Tibet, near Sikkim border, c. 12,000' (*H. W. Mulligan*).

Not known from elsewhere

42 *Theobaldia indica* Edwards, 1920 †

Bull Ent Res x, p 137 (♂ & ♀) TYPE-LOC - Bakloh, Gurdaspur dist., Punjab, 5,000', 28. II 1900 (*Lyndesay*) TYPE ♂ in Brit Mus

ADULT ♀ ‡.—Resembles *T. invertensata* in general appearance, but differs as follows:—Proboscis with plentiful sprinkling of pale scales on about basal $\frac{3}{4}$, palpi about $\frac{1}{2}$ length of proboscis; scutellar scales narrow, pale in colour, about 10 spiracular bristles. *Wings* with plentiful sprinkling of pale scales along costa, subcosta, vein 1, and usually a few at base of 5, membrane not definitely clouded, but aggregations of scales at base of 3 and at bifurcations of 2 and 4, giving the wing a slightly spotted appearance. *Legs* brownish-

† This species was recorded from India by Giles as *Culex annulatus* Schr., and by Theobald as *Theobaldia annulata* (Schr.), and was included under the latter name in Brunetti's and Senior-White's Catalogues. *T. annulata* is a *Palaeartio* species, and is not known from the Indian region at present; it differs from *indica* in having a pale ring at middle of segment I of hind tarsus.

‡ Barraud 1924 I, p. 142

black, with pale ochreous markings, a plentiful sprinkling of pale scales on femora and tibiae, pale dorsal marks at apices of latter, ventral surface of femora almost entirely pale, pale scales on hind tibiae tend to form stripes, tarsi dark, with rather broad basal pale rings to segments 2, 3, and 4, that on 4 very small on fore legs, segment 1 on each leg narrowly pale at base, and with sprinkling of pale scales on basal $\frac{1}{2}$ or more. Dorsum of *abdomen* brownish-black, with basal white bands of even width on tergites III-VI, some pale scales in middle of I, a pale basal band on II, produced in middle towards posterior margin, usually lateral pale patches on VII; venter covered with dull white scales and with indistinct lateral darker markings.

♂—Antennae plumose, proboscis sometimes pale for nearly whole length; palpi a little longer than proboscis, dark brown, pale yellowish at joints, and with extensive mottling of pale scales, some long hairs at apex of segment 3 and along 4, 5 less expanded than in the other Indian species. *Hypopygium* (fig. 19, c) very similar to that of *T. nivescentiata*, differing chiefly in the shape of the apical border of 8t and in form of 1p.

PUPA, LARVA, and EGG—Unknown.

DISTRIBUTION.—WESTERN HIMALAYAS and PUNJAB only: Bakloh (type-loc as above), Dalhousie, Gurdaspur dist., 6,000', 4. v. 1906 (*Barrow*), Kasauli*, Simla Hills, 6,000', iii & v. 1921, ii. 1924, ix. 1923 (*Barraud*), and ii. 1926 (*Sinton*), Muktesar*, Kumaon (U.P.) 7,500' (*T. B. Fletcher*); Ambala, Punjab, 9. iii 1905.

Not known from elsewhere.

Genus **ORTHOPODOMYIA** Theobald, 1904.


Entom xxxvii, p 237 Genotype, *O. albipes* Leno.

ADULT †—Mosquitoes of moderate size, resembling some *Anopheles* in having spotted wings (fig 21, a), speckled legs, and black and white rings on hind tarsal segments, or with the last three segments entirely white. The genus differs from other Culicini in the proportionate length of tarsal segments of fore and mid-legs; tarsal segment 1 rather longer than last four together, 4 very short in both sexes, much shorter than 5, and only a little longer than broad (fig. 21, b), 4 and 5 together about as long as 3, claws of ♀ simple, larger claw of fore leg of ♂ toothed. Antennae in both sexes about length of proboscis, those of ♂ plumose. Palpi of ♂ nearly

† Edwards 1932, p. 106. See also Theobald 1907, p. 527; Lang 1920, p. 109; Barraud 1927 a, p. 525, Barraud & Covell 1928, p. 676 (buccal cavity), Martini 1930, p. 193.

as long, or quite as long, as proboscis, the last one or two segments short and not swollen or very hairy, those of ♀ about $\frac{1}{3}$ length of proboscis. Wings rather narrow, with


Fig 21



Orthopodomyia anopheloides a, wing, b, tip of front tarsus of ♀.

very long fork-cells, vein *sc* long and terminating at about level of bifurcation of 2, vein 6 long, terminating much beyond fork of 5, squama fringed. Bristles on mesonotum well developed; 2-4 posterior pronotal, no spiracular, post-

Fig 22



Pupal structures of *Orthopodomyia*, showing dorsal chestotaxy of abdominal segments III-VII, etc.. a, *anopheloides* (with respiratory trumpet and paddles); b, *flavicosta*, c, *flavithorax*.


spiracular, or lower mesepimeral, but upper mesepimeral and sternopleural fairly numerous.

♂ *hypopygium*: apical border of tergite VIII produced into a median plate; coxite moderately long and narrow, with

basal lobe on inner surface and with style at apex carrying terminal appendage, phallosome small and simple, paraprocts with teeth at crown

♀ *hypopygium* cerci of only moderate length, three chitinised spermathecae present.

Fig. 23.



Larval structures of *Orthopodomyia* (head, tail-end, enlarged comb-scales) a, *anopheloides*; b, *flavicocta*, c, *flavithorax*.

PUPA (fig. 22) *—Respiratory horn of moderate length, with wide opening, dendritic tufts on abd seg. I well developed, one of the five submedian and sublateral hairs on tergites IV–VI long, tufted lateral hairs on VII and VIII fairly large. Paddles nearly twice as long as broad, rather flattened on outer side towards apex, apical $\frac{1}{2}$ very transparent, a short terminal hair, moderately developed midrib, an irregular dark marking across the base.

LARVA *, 4th stage (fig. 23)—Siphon without pecten, a single pair of hair-tufts at about middle of tube towards posterior margin, comb with 2 rows of teeth, some long single hairs on abdomen, antennal shaft smooth, tuft at $\frac{1}{4}$ or $\frac{1}{3}$ from base. In *O. anopheloides* and *O. flavicosta* there is a chitinated saddle on abd seg VIII, and in the former there is a well-developed saddle on VII also, but in *O. flavithorax* there is no very definite saddle on either segment, only some indefinite chitination on VIII.

Eggs †.—Those of Indian species are unknown.

DISTRIBUTION and BIONOMICS—Several species appear to have a very local and limited distribution, but *O. anopheloides* (Giles) is widely spread in India. Little is known of the habits of the adults, and they are not known to attack man. The larvae live in water in tree-holes, and are commonest during the monsoon. All the Oriental species occur in the Indian region, others are known from North and South America and Europe and one from Mauritius, but none from Africa or Australia.

Key to Adults.

- | | |
|---|---|
| 1 Three white spots on costa on apical $\frac{1}{2}$ of wing, in addition to 1–3 small spots on costa on basal $\frac{1}{2}$. Palpi of ♂ shorter than proboscis. | 2. |
| Two white spots on costa on apical half of wing (the spot near apex of wing often small); small spots on basal $\frac{1}{2}$ of costa present or absent. Palpi of ♂ as long as proboscis. | 5 |
| 2 Last three hind tarsal segments completely white. | 3. |
| Last three hind tarsal segments not completely white, second dark at base. | 4. |
| 3 Second hind tarsal segment dark at base, narrowly white at tip. | [<i>andamanensis</i> , p. 102.
<i>anopheloides</i> var. n. |
| Second hind tarsal segment narrowly white at base and usually also at tip. | <i>albipes</i> , p. 102 |
| 4 Segments 4 and 5 of hind tarsi completely white. | [<i>maculipes</i> , p. 101.
<i>anopheloides</i> var. |
| Segment 4 of hind tarsi with black sub-apical ring, segment 5 white. | [var. <i>maculata</i> , p. 101.
<i>anopheloides</i> , p. 98, and |

* Barraud 1932, p. 1013; Edwards 1932, p. 107.

† Howard, Dyar & Knab 1917, p. 878.

- 5 Mesonotal scales mainly rather bright golden; two pale spots at base of costa, but no scattered yellow scales on basal $\frac{1}{2}$ of that vein; often no white spot on vein 5 at fork in ♀ *flavithorax*, p 103.
- Mesonotal scales brownish, at most one pale spot at base of costa, but scattered yellow scales present on basal $\frac{1}{2}$ of that vein, a white spot on vein 5 at fork in both sexes *flavicoستا*, p 104.

Key to Pupæ

1. Longest hair on tergite VI reaching beyond posterior margin of VIII, and either single or 2-branched 2.
 Longest hair on VI not reaching beyond posterior margin of VII, and usually of 4 branches (fig 22, a) *anopheloides*
2. Longest hair on IV-VI single (fig 22, b) *flavicoستا*
 Longest hair on IV and V 3-branched, that on VI 2-branched (fig 22, c) *flavithorax*

Key to Larvæ (4th stage)

- 1 Abd. seg. VII with a large chitinous saddle, one branch of ω much longer than the others; a separate lateral chitinous strip at base of anal segment on each side (fig 23, a) 2.
 Abd. seg. VII without a chitinous saddle, all branches of ω about the same length; no separate chitinous strip at base of anal segment 3.
- 2 Comb-teeth of larger row with long lateral branches. Comb-teeth of larger row with short fringe only *anopheloides*
albipes
- 3 Larger comb-teeth ending in a single long point; 5 pairs of clypeal hairs equally developed, and with long plumose branches, hairs on sides and dorsum of thorax remarkably developed and with long plumose branches (fig 23, c) *flavithorax*.
 Larger comb-teeth ending in several sharp points, frontal hair d comparatively small, with only 3 or 4 fine branches, e fine and single or split, hairs on dorsum of thorax fine and inconspicuous, lateral hairs only being well developed (fig 23, b). *flavicoستا*.

43 *Orthopodomyia anopheloides* (Giles), 1903

- Journ Trop Med vi, 315 (*Mansonia*) (♂ & ♀). TYPE-LOC.: Delhi, Dun, U.P., India (*Wyville-Thomson*) TYPE: ♂ & ♀ in Brit. Mus.
- "*Orthopodomyia nigritarsis* var." Leicester, 1908, Cul. Malaya, p 177 TYPE-LOC "The Dindings," Pangkor Laut, F.M.S. (*Daniels*) TYPE: ♀ in Brit. Mus.
- var. *maculata* Theobald 1910, Rec Ind. Mus. iv, p 29 (♂). TYPE-LOC.: Maddathora, Travancore, xi. 1908 (*Annandale*). TYPE: ♂ in Ind. Mus.
- var. *maculipes* Theobald 1901, M.C. v, p. 470 (♂ & ♀) (*O. maculipes*). TYPE-LOC.: Andaman Is., vii. 1908 (*Lewis & Ray White*). CO-TYPE: ♀♀ in Brit Mus
- var n. *andamanensis* (see p 102) TYPE-LOC.: Andaman Is., viii. 1926 (*G. Covell*) TYPE: ♂ & ♀ in Brit. Mus.

ADULT * —Medium-sized species, wing 4-6 mm.

♀.—*Head*: dorsal surface thickly covered with large pale-coloured upright forked scales, some narrow yellow scales in front and along eye-margins, a few black and numerous flat white scales at sides, some white scales on torus and on 1st flagellar segment of antenna, palpi rather less than $\frac{1}{2}$ length of proboscis, black or dark brown, with white markings in middle and at tip. Proboscis with a moderately wide white band on apical $\frac{1}{2}$, and usually a white streak on upper surface between the band and tip. *Thorax*: mesonotum covered with narrow pale yellow and white scales, the latter arranged chiefly along sides and over wing-roots, three darker spots forming a triangle posteriorly, in some specimens the most anteriorly situated spot is very small or absent, in others there are several additional dark spots anteriorly. Scutellar scales long, white, and narrow, some on lateral lobes broader. Pleurae covered for the most part with flat dull white scales, arranged chiefly around the various groups of bristles, scales on upper part of *ppn* narrow and yellow. *Wings* (fig 2, a) veins clothed with black and white scales, latter arranged chiefly in four or five spots along anterior part of wing. Two small spots at base of costa, sometimes confluent, a subbasal spot on *c*, *sc*, and vein 1, a spot at *sc* junction including vein 1, a subapical spot including *c*, 1, and fork of 2, an apical spot including *c*, tip of 1, 2.1, and 2.2. Smaller spots at tips of 3, 4.1, 4.2, and 5.1, also at fork of 4 and on 2, 3, 4, and 5.1 in regions of cross-veins, on 1 and 2 at origin of latter, a small spot at extreme base of 5, with another immediately above it on 1 and *sc*. *Legs*: femora dark brown, speckled with numerous white and pale yellow scales, not arranged in definite spots, a rather wide subapical pale ring not always continued completely round leg, knees narrowly white. Fore and mid-tibiae sprinkled with pale scales anteriorly, almost entirely yellow posteriorly, except for subbasal and subapical dark rings, hind tibiae more extensively pale but dark at base; all tibiae usually white at tips. Fore tarsi dark, with a few pale scales at joints, mid-pair with well-marked narrow basal and apical white rings over joints, segments 4 and 5 mainly pale in some specimens, segment 1 of hind tarsi narrowly pale at base and tip, 2 dark at base, rather narrowly white at tip, 3 and 4 black, with broad basal and narrow apical white rings, 5 entirely white. *Abdomen*: dark brown, tergites II-VII with narrow basal white bands, not always visible in dried specimens, and small apical lateral white spots; a pair of

* Edwards 1913 b, p. 239, 1926 a, p. 117 (syn.); 1932, p. 108; Barraud 1927 a, p. 527 (syn.).

small submedian pale spots in middle of tergites IV-VII. Sternite II almost entirely pale, narrow basal pale bands on III-V.

♂—Resembles ♀, but antennal plumes dense, hairs long and arranged chiefly on dorsal and ventral aspects, pale yellow in colour, some white scales on inner side of first 4 or 5 flagellar segments. Palpi about $\frac{3}{4}$ length of proboscis, black-scaled, with white scaling at base and tip and a white ring in middle. Proboscis dark brown or black, with a white ring at about the middle and a white spot dorsally at tip.

PUPA * (fig. 22, a)—Distinguished from those of *O. flavithorax* and *O. flavicosta* by the length of the longest hair on tergite VI, this being comparatively short, and usually of 4 branches. The innermost submedian hair on III-VII is very small, whereas in the other two species it is larger and spine-like.

LARVA * (fig. 23, a).—*Head*, including antenna, deep brown or black, lighter areas on head around eyes. Frontal hair *A* with 9-11 branches, *B* and *d* with 6-8 (the latter well developed), *C* with 4-6, *e* fine and long, often split into two towards tip. Preclypeal spines fairly long, slender and tapering. Antennal shaft about 0.41 mm long, 4-6 times as long as greatest breadth, stout on rather more than basal $\frac{1}{2}$, tapering to tip; tuft of 6 to 8 branches at $\frac{1}{3}$ from base, apical spines all at tip of shaft. *Thorax* with lateral tufted hairs well developed and some long single hairs. *Abdomen* usually reddish in colour, first two segments with lateral tufted hairs, III-VIII with fine hairs, some of which are single and long, VII with a chitinated saddle covering rather more than the dorsal $\frac{1}{2}$ in side view, saddle on VIII nearly enclosing the segment. Comb with 7-10 larger teeth and 25-30 smaller fringed teeth; the largest teeth very long, with main terminal portion of moderate thickness and several long branches on each side which are slightly less stout. *Siphon* dark brown or black, lighter at apex, 1-1.1 mm long and about 5 times as long as diameter at base, tuft of 8 to 14 strong subplumose branches arising at about middle of tube, but very slightly nearer base than apex. *Anal segment* enclosed in a chitinated ring, a separate chitinated strip on each side at base; *osc* single and long, *isc* split into a number of branches, one much longer than the others, *lh* with 2 or 3 fine branches. Ventral papillae about $\frac{1}{2}$ length of dorsal, latter rather more than $\frac{1}{2}$ length of anal segment; fan well developed, hairs arising from a fan-plate and each splitting into a number of branches some distance from base.

HABITAT.—Tree-holes during the monsoon.

* Barraud 1932, p. 1015.

DISTRIBUTION.—Fairly common in the WESTERN and EASTERN HIMALAYAS from foothills up to 7,000' or more, in ASSAM along Brahmaputra Valley and in hills to the south, WESTERN GHATS from north of Bombay to NORTH KANARA (Yellapur)

Not definitely recorded from outside the Indian area, but Leicester's *O. albipes* var. *nigritarsis* from MALAYA is probably this species, as it agrees in the markings of the hind tarsi.

[var *maculata* Theo

ADULT.—Differs from typical *O. anopheloides* in the wider white band over joint between 2nd and 3rd hind tarsal segments, white area on 2nd segment occupying $\frac{1}{3}$ — $\frac{1}{2}$ its length instead of $\frac{1}{3}$ — $\frac{1}{2}$ as in the typical form, and white at base of 3rd segment occupying $\frac{2}{3}$ instead of barely $\frac{1}{2}$ the length of this segment

LARVA (Javan specimens)—Apparently indistinguishable from typical *anopheloides*

HABITAT.—Foul water in stumps of giant bamboo and in tree-holes.

DISTRIBUTION.—TRAVANCORE: type-locality, as noted above. CEYLON Peradeniya, Suduganga This variety perhaps replaces typical *O. anopheloides* in South India and Ceylon, all the specimens in the British Museum from Ceylon are of this form, as are those from JAVA (Buitenzorg R. W. Parne)]

var. *maculipes* Theo *

ADULT.—Very similar to typical *O. anopheloides*, but last two hind tarsal segments completely white. Segment 3 of hind tarsi marked with a narrow black subapical ring. Differs from *O. albipes* and *O. anopheloides* var. *andamanensis* in having only two, not three, hind tarsal segments completely white, and from *O. albipes* also in having segment 2 of hind tarsi dark at base.

PUPA.—Unknown.

LARVA †.—Very similar to, if not identical with, that of *O. anopheloides*.

DISTRIBUTION.—I have not seen any specimens agreeing with Theobald's description. It is known only with certainty from the ANDAMANS and SINGAPORE

* Theobald appears to have confused two forms under this name. Edwards (1913, p. 239) states that the specimens recorded by Theobald (1910, p. 473) from Ceylon were *O. anopheloides*. The type ♂ mentioned by Theobald, under the same reference, as being in the Indian Museum is not on record there. He was referring most probably, by mistake, to the type ♂ of *O. maculata*, which is there; it is also a specimen of *O. anopheloides*.

† Edwards 1928, p. 118 (Singapore specimens).

var n *andamanensis*.

ADULT.—Differs from typical *O. anopheloides* as follows.—Segment 2 of hind tarsi only very narrowly white at tip, 3–5 entirely white. Proboscis of ♀ with a wider pale band. Proboscis of ♂ nearly all yellow on apical $\frac{1}{3}$. Differs from *O. albipes* in having definite apical and basal banding on fore and mid-tarsi.

PUPA and LARVA.—Unknown. Until these have been discovered it is not possible to say whether this is a variety of *O. anopheloides* or of some other species, or possibly distinct.

DISTRIBUTION.—ANDAMAN IS *, as above, also ix. 1911 (*Christophers*). One ♂ from Sukna *, DARJEELING DIST., viii. 1928 (*Sobha Ram*), is of this form. This was collected at the same time and place as typical *O. anopheloides* and *O. albipes*. SINGAPORE (*Given*), 1 ♂, 1 ♀ in Brit Mus., in company with var *maculipes*.

44 *Orthopodomyia albipes*, Leicester in Theobald, 1904.

Entom xxxvii, p 237, Cul Malaya, p 176, 1908 TYPE-LOC.: Kuala Lumpur, F M S (*Leicester*) TYPE ♂ & ♀ in Brit Mus

* *Kerteszia mcgregori* Banks, 1909, Phil J Sci A, iv, p 548 TYPE-LOC. Philippine Is

ADULT.—Resembles *O. anopheloides* except as follows:—Segment 2 of hind tarsi narrowly white at base and usually very narrowly so at tip †, segments 3–5 entirely white. Pale markings on fore and mid-tarsi mainly basal and segment 5 mainly pale. Proboscis of ♀ with a wider pale band; palpi of ♀ with narrow white band in middle and narrow apical and subapical white markings close together. Proboscis of ♂ with large creamy subapical area nearly surrounding the organ.

[LARVA ‡.—A Malayan specimen (not isolated) in the British Museum differs from *O. anopheloides* in having the head pale, antennæ longer (8 times as long as their greatest breadth), and the larger comb-teeth without long lateral branches, but with a fine fringe of short hairs only, leaving the distal $\frac{1}{3}$ bare, siphon rather longer and more pointed.]

HABITAT.—Bamboos (*Leicester*).

DISTRIBUTION.—Known in India only from Sukna, DARJEELING DIST., 500', viii. 1928 (*Sobha Ram*), 3 ♀♀, 1 ♂, taken in company with *O. anopheloides*.

Outside India known from MALAY PENINSULA, BORNEO, and PHILIPPINE IS.

† [Often, but by no means always, without any white scales at tip in Malayan specimens.]

‡ Edwards 1926, p. 118.

45 *Orthopodomyia flavithorax* Barraud, 1927 *

Ind Journ Med Res xiv, p. 529 (♂ & ♀) TYPE-LOC Karwar,
N Kanara dist., ix 1921 (*Barraud*). TYPE ♂ & ♀ in Brit Mus

ADULT —Differs from *O. anopheloides* as follows —Mesonotal scales bright yellow or golden, wings without the subbasal and subapical white spots on *c*, etc, white markings on mid-tarsi at bases of segments only, pale scales on femora and tibiae arranged in definite spots, palpi of ♂ as long as proboscis, white scales on 1st flagellar segment only of antenna of ♂. proboscis of ♂ with a white marking on upper side rather beyond the middle, and usually another marking on underside nearer tip, the two markings sometimes forming nearly a complete band, another white marking on upper side near tip, proboscis of ♀ with a white band and yellow spots, variable in extent, sometimes the whole organ is strongly spotted with yellow, the white band being obliterated.

A rather variable species. Frequently in the ♀ there is a white spot on the stem of vein 5 at the fork. This spot is present in the ♂♂, including the type.

PUPA † (fig. 22 c).—One of the sublateral hairs on tergites IV-VI very long, these long hairs on IV and V with 3 branches, VI 2-branched (In *O. anopheloides* the corresponding hair on VI is comparatively short and usually 4-branched, in *O. flavicosta* it is long and simple, as are also the longest hairs on IV and V.)

LARVA † (fig. 23 c).—This has a very characteristic appearance, even to the naked eye, owing to the remarkable development of the hairs on the head and thorax. Hairs *A*, *B*, *C*, *d*, and *e* on the clypeus are all about equally developed and each has a number of subplumose branches which are as long as, or longer than, the clypeus. Antennal shaft about 0.58 mm long, about 6 times as long as greatest breadth, stout on basal $\frac{1}{3}$, much more slender apically, brown for the most part, paler at tip; tuft of about 5 subplumose branches, arising at about $\frac{1}{3}$ from base; apical bristles all very near tip of shaft. Preclypeal spines long, slender, tapering. *Thorax* densely clothed dorsally and laterally with long tufted hairs with subplumose branches. *Abdominal* segments with some very long single or 2-branched hairs, as well as shorter tufted hairs with subplumose branches,

* [*O. arboricollis* d'Emm. (endemic in Mauritius) shows much resemblance to *O. flavithorax*, but has all the wing-scales larger and broader, with many scattered white scales in addition to those forming definite spots; femora and tibiae heavily sprinkled with pale scales which are not aggregated into spots; no scattered pale scales on tarsi, 4th hind tarsal segment black, abdominal tergites basally banded.]

† Barraud 1932, p. 1015.

VII without a chitinated saddle, VIII with indefinite chitination not forming a definite saddle. Comb with 10-12 comparatively very large pointed teeth and 18 to 20 smaller delicately fringed teeth. *Siphon* 1.4-1.5 mm long and 5-6 times length of diameter at base, dark brown, a very narrow darker ring at base, paler at extreme tip, tuft of about 8 finely frayed branches, arising at about $\frac{2}{3}$ the length from base. *Anal segment* enclosed in a chitinous ring, no separate chitination at base, *osc* single and long, 6 or 7 times length of chitinated part of segment, *isc* split into a number of branches all about same length and about $2\frac{1}{2}$ times length of segment, ventral papillae about $\frac{1}{2}$ length of dorsal pair, latter only about $\frac{1}{2}$ length of segment, *lh* of 2 or 3 fine branches, fan well developed, about 14 hairs arising from fan-plate, each hair split some distance from base into a number of fine branches.

HABITAT.—Tree-holes during the monsoon.

DISTRIBUTION.—S W INDIA Malabar Coast *, x 1915 (Khazan Chand), Karwar * and Kadra*, N. Kanara, ix 1921 (Barraud) CEYLON Madola, near Opanayaka, ii. 1933 (Henry)

46 *Orthopodomyia flavicosta* Barraud, 1927

Ind. Journ Med Res xiv, p. 531 (♂ & ♀) TYPE-LOC Karwar, N Kanara dist, ix 1921 (Barraud) TYPE ♂ & ♀ in Brit. Mus

ADULT.—Very similar to *O. flavithorax*, described above, and differs from *O. anopheloides* in very similar details. From *O. flavithorax* it differs in the browner scales, in markings of proboscis, and in having only one small spot at base of costa, or none, the basal $\frac{1}{2}$ of that vein being thickly sprinkled with yellow scales, which may be so numerous as to form a streak. Proboscis in ♀ with a small white spot on upper side and usually some pale scaling on underside nearer to base (not nearer to tip, as in *O. flavithorax*); in the ♂ the white band is produced on underside for some distance towards tip. Palpi of ♂ as long as proboscis, and pale scales on femora and tibiae arranged in definite spots in both sexes, as in *O. flavithorax*.

PUPA † (fig 12).—Very similar to *O. flavithorax*, but longest sublateral hairs on tergites IV-VI single

LARVA † (fig 23 b).—Differs in several respects from that of *O. flavithorax*, with which it is found breeding in association, the hairs on the head and thorax being much less developed, and the larger comb-teeth being of different form. *Head* yellowish, slightly darker posteriorly. Antennal shaft about 0.5 mm long, tapering from base to apex, comparatively slender, about 11 times as long as greatest breadth.

† Barraud 1932, p 1016

paler on apical part than towards base, tuft of about 6 branches, arising at a point between $\frac{1}{2}$ and $\frac{1}{3}$ length of shaft from base. Frontal hair *A* usually of 9 branches (variation 9 to 12); *B* of 6 or 7 branches, *C* of 4 or 5, *d* of 3 or 4 fine branches much shorter than *B* or *C*, *e* fine and usually single, but may be split into 2 towards the tip. Preclypeal spines fairly long and tapering, pale in colour. *Thorax* with lateral tufted hairs well developed, but hairs on dorsum fine and inconspicuous. First two *abdominal* segments with lateral tufted hairs, following segments with fine hairs, some of which are long and single as usual, VII without a chitinated saddle, VIII with a definite saddle nearly enclosing the segment. Comb with 7 to 10 larger teeth and about 20 smaller fringed teeth, the larger teeth end in a number of sharp points and resemble a hand, there being usually 5 flattened leaf-like points. *Siphon* 1.2 mm long, 5 or 6 times length of diameter at base, tuft of about 6 branches, finer than in the other three species here described, arising very slightly nearer base than apex of tube. Anal segment very similar to that of *O. flavithorax*.

HABITAT—Tree-holes during the monsoon.

DISTRIBUTION—S W INDIA only. Karwar and Kadra, North Kanara, ix 1921 (*Barrard*).

Genus FICALBIA Theobald, 1903

M C iii, p. 296 Genotype, *F. minima* Theo

Mimomyia Theobald, 1903, M C iii, p. 304 Genotype, *M. splendens* Theo.

Etorleptomyia Theobald, 1904, 1st Rept Wello Lab p. 71 Genotype, *E. mediolineata* Theo

O'Reilia Ludlow, 1905, Can. Ent xxxii, p. 101 Genotype, *O'R. luzonensis* Ludl

Ludlowia Theobald, 1907, M C iv, p. 193 Genotype, *L. chamberlaini* (Ludl).

Radioculex Theobald, 1908, Rec Ind Mus ii, p. 295. Genotype, *R. clavipalpus* Theo

Dasyomyia Leicester, 1908, Cul Malaya, p. 102. Genotype, *D. fusca* Leic


Conopomyia id., ib. p. 113 Genotype, *C. metallica* Leic.

Etorlepidomyia Alcock 1911 (emend.), Ann & Mag. Nat Hist (8) viii, p. 249

This genus has recently been extended to include *Mimomyia* Theo., which was formerly classed separately. As regards the Indian region, five species are known, representing three different subgenera. These might reasonably be regarded as of generic rank, but this is a matter of opinion, and it seems inadvisable to change the classification again at present. The characters of the subgenera are indicated in the key on pp 107-8.

ADULT*.—Small, or rather small, mosquitoes, antennæ about length of proboscis, proboscis of ♂ distinctly widened towards tip, palpi of ♂ not more than about $\frac{1}{4}$ length of proboscis, *apn* well separated, no spiracular or postspiracular bristles, one lower mesepimeral or none, several strong *ppn*, other groups of bristles fairly well developed. Pleuræ not very scaly. Segment 1 of hind tarsi shorter than tibia, segment 4 of all tarsi of ♀ as long as, or longer than, 5, no pulvilli; claws of ♀ simple, fore and mid-claws of ♂ unequal and toothed. Vein 6 terminating nearer apex of wing than

Fig 24.



Pupal characters (respiratory horn and tip of abdomen or paddle) of
Ficalbia a, b, *chamberlaini*, c, d, *fusca*, e, f, *hybridula*, g, *luzonensis*

level of fork of 5, squama with complete fringe. Hypopygium of ♂ of simple structure, coxite fairly long, with a bristly basal lobe on inner side. Style long, tapering, with terminal appendage. Phallosome thinly chitinated and tube-like, without lateral plates or teeth. Paraprocts with several small teeth at crown. Ninth tergite with median lobe bearing hairs on each side. Hypopygium of ♀ with one to

* Edwards 1932, p. 108. See also Macfie & Ingram 1922, p. 182 (hypopygium), Barraud 1929 a, pp. 1054-6.

three spermathecae, according to species, segment VIII short and broad, cerci short

PUPA (fig. 24).—Respiratory horns remarkably long and narrow, resembling long thin tubes with opening at extremity (in the aberrant species *F. hybrida* (Leic) the apical part of the horn is modified for piercing tissues of plants and divided into leaflets) Dendritic tufts on abd. seg I not well developed or absent. Paddles rather narrow or very narrow, with serrations or spines along margins, at least towards apex, usually no terminal hair When the pupa is resting at the surface the respiratory horns are held forwards and the paddles flexed towards the dorsum of the abdomen

LARVA, 4th stage (figs 25-27).—The different species exhibit great diversity of form, but the following characters are common to the Indian species:—Head large, broader than long, antenna with large tuft and with subapical hairs some distance from apex Pleural plates of meso- and metathorax large, usually bearing a long spine. Abd seg VIII without a lateral plate, but with a varying number of usually simple teeth Siphon 4-6 times length of diameter at base, with few simple pecten-teeth or none, one pair of hair-tufts some distance from base with few branches Anal segment with complete chitinous ring and with moderately developed fan but no fan-plate, lateral hair usually long, both pairs of subdorsal hairs usually of several branches

Eggs—Unknown.

DISTRIBUTION and BIONOMICS.—Found chiefly in the southern and eastern parts of the Indian region, the species are not common, and little is known of their habits, they do not appear to attack man The genus is represented in Africa, and extends through the Oriental region to Australia. So far as is known, the larvæ live in large permanent pools containing much vegetation, or in fallow rice-fields

Key to Adults (species and subgenera).

- 1 Fork-cells short, *af* only about $\frac{1}{4}$ length of stem, palpi of ♂ slightly longer than proboscis, last segment thickened, 3 chitinated spermathecae in abdomen of ♀ (subgenus *Mimomyia*) 2.
- Fork-cells not shortened, *af* as long as, or longer than, stem; palpi of ♂ shorter than proboscis and slender, 1 chitinated spermatheca in abdomen of ♀ 3.
2. Mesonotum with conspicuous yellow scaling laterally; tarsi with yellow rings, segment 5 of hind tarsi entirely yellow *chamberlaini*, p 108.
- Mesonotum without conspicuous yellow scaling, tarsi not ringed; segment 5 of hind tarsi dark *hybrida*, p. 111.

- 3 Palpi of ♂ about $\frac{1}{2}$ – $\frac{2}{3}$ as long as proboscis,
1st flagellar segment of antenna of ♀
not much longer than 2nd (subgenus
Etorleptomysia) 4
Palpi of ♂ only about $\frac{1}{3}$ as long as proboscis
(fig 27, h), 1st flagellar segment of an-
tenna of ♀ about 3 times as long as
2nd (fig 27, g) (subgenus *Ficalbia*) 5
- 4 Wings speckled with light and dark broad
scales, tarsi mainly yellow, spotted with
black, abdomen pale, with median dorsal
dark stripe *luzonensis*, p 113.
Wings not speckled, scales all dark and less
broad, tarsi mainly dark, but pale
beneath, dorsum of abdomen dark *fusca*, p 115.
- 5 Tarsi with narrow pale rings, most distinct
on hind legs, dorsum of abdomen with
transverse pale bands *minima*, p 116

Key to Larvæ (4th stage).

- 1 Maxilla with a strong black apical spine
resembling clypeal spines, 1–3 stout
pecten-teeth on siphon 2
Maxilla without apical spine, no pecten-
teeth on siphon (subgenus *Etorleptomysia*) 4.
- 2 Antennal tuft near middle, clypeal spines
simple, siphonal tuft near middle (sub-
genus *Mimomyia*) 3
Antennal tuft far beyond middle, clypeal
spines barbed, siphonal tuft near base
(subgenus *Ficalbia*) *minima*
3. Siphon of moderate length and normal
shape, valves large, one pair each with a
long, fine hair *chamberlaini*
Siphon very short, broad at base but very
narrow at tip, valves very small, without
long hairs *hybrida*.
4. Subapical hairs of antenna simple, mentum
with central tooth, comb-teeth in a
regular row *luzonensis*
Subapical hairs of antenna dendritic,
mentum without central tooth, comb-
teeth in a patch *fusca*

47 *Ficalbia (Mimomyia) chamberlaini* (Ludlow), 1904

Can Ent xxxvi, p 297 (*Mimomyia*) (♂) TYPE-LOC. Bayamban,
Luzon, P I, v 1904 (*Chamberlaini*). TYPE ♂ in U.S. Nat Mus
Radroculex clavipalpus Theobald, 1908, Rec Ind Mus. ii, p. 295.
(♂ & ♀). TYPE-LOC. Calcutta, xi–xii 1907 (*Mus. collr.*)
TYPE ♂ & ♀ in Ind Mus
var *intermedia* Barraud, 1929, Ind Journ. Med. Res. xvi, p 1055.
(♂ & ♀). TYPE-LOC Gauhati, Kamrup dist., Assam, iv. 1927
(*Barraud*). TYPE ♂ & ♀ in Brit. Mus

ADULT*.—Easily identified by the markings The only

* Theobald 1907, p 194 (*Ludlowia*); Edwards 1911, p 244 (syn.).
Dyar & Shannon 1925, p. 82, Barraud 1929 a, p 1055

other Indian mosquito with which it could be confused is *Aedes (Banksinella) lineatopennis* (Ludl.), which has a similar ornamentation of the mesonotum but entirely dark tarsi * Wing about 2.7 mm

♀.—*Head*. dorsal surface covered with flat scales, usually appearing dull ochre-yellow, with a darker patch on either side towards front, in some specimens the scales appear mainly dark, some very small dark upright scales on nape. Antennæ and clypeus dark brown. Palpi dark brown at base, a little lighter in middle and at tips; about $\frac{1}{2}$ length of proboscis. Proboscis rather long, about length of abdomen, distinctly swollen apically in side view, dark bronzy brown, underside paler. *Thorax*. integument of mesonotum shining brownish-black, with narrow scales and bristles of same colour, a large well-defined patch of bright yellow scales over, and in front of, each wing-root. Scutellar scales narrow and dark brown. Postnotum dark brown, sides paler. Integument of pleuræ uniformly pale. *Wings*. dark scaled, fork-cells short, *af* much shorter than its stem. *Legs*: fore femur yellowish with bronzy sheen, darker apically, mid- and hind femora bronzy brown, paler on posterior surfaces. Tibiæ and tarsi with dark violet, bronzy, or brassy sheen, hind tibiæ pale at tips, narrow yellow markings at bases of segments 2 and 3 of tarsi of fore and mid-legs, similar, but more pronounced, markings on joints between tarsal segments of hind legs, segment 5 entirely pale. *Abdomen*. dorsum entirely covered with dark scales having a violet sheen, venter pale yellowish.

♂.—Antennal shaft yellow, plume-hairs brown and dense. Palpi slightly longer than proboscis, yellow, with a dark ring at joint between last two segments and a dark marking at tip of last segment, which is swollen. Proboscis yellow, with dark tip, widened on apical $\frac{1}{2}$. Other markings as in ♀.

var *intermedia* Barraud differs in having large lateral yellow patches on abdominal tergites, uniting in some specimens to form basal bands. This form appears to agree closely with Miss Ludlow's original description. It is possible that this form, which occurs in Assam (and presumably in the Philippines), may be specifically distinct from the *claripalpus* form described from Calcutta. The larva of the variety has not yet been obtained.


PUPA (fig. 24, a, b).—Respiratory horns very long, slightly expanded at apices, a small dendritic tuft on I, paddles

* [Two other Oriental species are nearly related to *F. chamberlaini*: (1) *F. metallica* Leic (Malay Penin. and Queensland), which differs in having a more or less continuous median yellow abdominal stripe; and (2) *F. aurea* Leic (Malay Penin and Borneo), which lacks the yellow borders to mesonotum and has the abdomen distinctly banded.]

wider than in *F. luzonensis* and *F. fusca*, brownish, with pale area on outer side and strong spines along apical margins

LARVA * (fig. 25, e-g)—*Head* rather large, dark brown or black. Antennal shaft long, very dark on basal $\frac{2}{3}$, with strong spinelets along its length, tuft at about middle point composed of a number of plumose branches, subapical bristles

Fig 25



Larval characters of *Ficalbia*, subgenus *Mimomyia*: a-d, *hybrida*; e-g, *chamberlaini* (d, comb-teeth)

long and placed some distance below tip, one long and one shorter apical bristle. Frontal hair *A* with a number of densely plumose branches, *B* with 2-4 long plumose branches, *C* with 6-8, *d* rather small, with about 5 fine branches; *e* fine and usually split into two some distance from base. Hair *d*

* Barraud 1923 h, p 504. See also Edwards & Given 1928, p. 349.

almost level with *B*, *C* slightly posterior to both, all three fairly close together. A very long 2-branched hair ventrally on head near base of antenna. Preclypeal spines long and black. Mouth-brushes fairly large, hairs simple. Lateral hairs on *thorax* well developed, arising from chitinated tubercles having small thorn-like processes. Some of the hairs very long, reaching forwards as far as tip of antenna. *Abdominal* hairs fairly long, especially lateral series on first two or three segments. Comb of 10–16 small fringed teeth, majority in one row, but two, usually separate from main rank, towards posterior border of segment. *Siphon* comparatively short, slightly curved forwards towards tip, 4–5 times length of diameter at base, tufts of about 4 branches, towards posterior border, at about middle of tube. Pecten of 1–3 strong simple teeth at base of tube. *Acus* small. Valves normal, but one pair with a very long fine hair about $\frac{1}{2}$ length of siphon. Anal segment enclosed in chitinous ring, much narrower ventrally than dorsally, posterior border spiny, *lh* very long and subplumose, *osc* and *isc* with a number of branches. Both pairs of papillæ rather short and pointed, fan moderate, no fan-plate.

HABITAT—Pools and ponds, especially those containing much vegetation.

DISTRIBUTION—From the PUNJAB to ASSAM and BURMA, and through PENINSULAR INDIA to CEYLON, but seldom abundant.

Apart from the PHILIPPINE ISLANDS, it has not been recorded from elsewhere.

48 *Ficalbia (Mimomyia) hybrida* (Leicester), 1908*

Cul Malaya, p. 115 (♂ & ♀) (*Conopomyia*). TYPE-LOC. Kuala Lumpur, Malay Penin (Leicester) (♂), Singapore (Finlayson) (♀). TYPE ♂ & ♀ in Brit Mus.

Ludlowia minima Ludlow, 1907, Can. Ent. xxxix, p. 413 (♂ & ♀). TYPE-LOC. Carandagan, P. I., 1906 (Duncan). TYPE ♂ & ♀ in US Nat. Mus.

Ficalbia ludlowi Brunetti, 1920, p. 173 (nom. nov. for *L. minima* Ludl.).

No 1 *Undetermined sp.* Leicester, 1908, Cul Malaya, p. 260 (♀). TYPE-LOC. Kuala Lumpur, Malay Penin (Leicester). TYPE: ♀ in Brit Mus.

ADULT †—A rather small, obscure species, sometimes mistaken for *Uranotaenia* on account of the short anterior

* Owing to the inclusion of *Mimomyia* in *Ficalbia*, the name of this species is now *hybrida*. It has been referred to previously by most writers as *M. minima* (Ludl.).

† Theobald 1910 b, p. 191; Edwards 1922 d, p. 468 (syn.); Barraud 1929 a, p. 1054; Bonne-Wepster 1932, p. 69.

fork-cell, but distinct on generic characters Wing about 3 mm

♀—*Head* dorsal surface covered with flat greyish or yellowish scales, some very small black upright scales on nape. Antennae, clypeus, and palpi brown, palpi about $\frac{1}{2}$ length of proboscis, 1st flagellar segment of antenna fully twice length of 2nd. Proboscis brown, darker on apical $\frac{1}{3}$, which is distinctly swollen. *Thorax* mesonotal scales dark brown. Postnotum almost black Integument of pleurae pale brown. *Wings* scales dark and moderately broad, fork-cells quite short *Legs* dark brown, femora lighter, tibiae and tarsi with pale yellowish sheen when seen in certain positions, tibial bristles distinctly long *Abdomen* dorsum almost black, with narrow basal pale yellow bands widening out at lateral margins of tergites Sternites pale yellow

♂—Markings as in ♀ Antennal plumes dense, pale brown. Palpi slightly longer than proboscis, deep brown, with some mottling of paler scales, last segment swollen, rather hairy, and very dark in colour.

[PUPA * (fig. 24, e, f).—Respiratory horns long, basal $\frac{3}{4}$ annular, distal $\frac{1}{4}$ smooth, modified for piercing by being deeply split at tip into two spine-like portions. Dendritic tuft on I small Paddles moderately broad, wholly dark; with short, sharp spines along apical margins, deeply emarginate at tip, without terminal hair

LARVA *† (fig. 25, a-d).—*Antennae* long, with a dark ring at base, otherwise pale, strong spinules extending as far as subapical hairs, which are long, tuft just before middle of antenna, portion beyond subapical hairs very long and slender, apical hairs both short. *Frontal hairs* A, B, and C long, with about 12, 3, and 5 branches respectively, d short, with about 5 branches, e single. Preclypeal spines long and black; similar spines present on maxillae as in other species of this subgenus *Comb* of about 10 finely fringed teeth in a fairly regular row. *Siphon* short, conical, its distal $\frac{1}{4}$ very narrow, with remarkably small valves which do not bear any obvious hairs, a single fine hair at mid-length representing the tuft, 2 strong, simple pecten-teeth. Anal segment much as in *F. chamberlaini*

HABITAT.—Among *Pistia* in a fresh-water pond together with *Mansonia uniformis*. The larvae presumably insert the narrow tip of the siphon into the *Pistia* roots to obtain air;

* Bonne-Wepster 1932, p. 71, re-described above from Javan specimens presented to the British Museum by Mrs. Bonne.

† [The larva recorded by Edwards and Given (1928, p. 349) as possibly "*M. minima* (Leic.)," i. e., *minima* Ludl = *hybrida* Leic., proves, on re-examination, to be that of *F. luzonensis*]

the whole siphon is similar in shape to that of *Mansonia*, although the valves are not specially modified as in that genus]

DISTRIBUTION — ASSAM Gauhati*, Kamrup dist., iii. 1925 (*Barraud*), Golaghat*, Sibsagar dist., v. 1925 (*Barraud*). BENGAL Dacca*, xii. 1911 (*S. K. S.*), Santragachi*, Howrah, viii. 1926 (*Senior-White*), Garden Reach*, Calcutta, xi & xii 1930 & 1931, and viii 1931 (*Senior-White*). BIHAR Kierpur (*Parva*)

Known also from MALAYA and PHILIPPINES

49 *Ficalbia* (*Etorleptomyia*) *luzonensis* (Ludlow), 1905.

Can Ent xxxvii, p 101 (♀) (*O'Reilly*) TYPE-LOC Bayambang, Luzon, P I, ix 1904 (*Chamberlain*) TYPE. ♀ in US Nat Mus
Etorleptomyia completiva Leicester, 1908, Cui Malaya, p 178 (♂).
TYPE-LOC Singapore (*Finlayson*) TYPE. ♂ in Brit Mus

ADULT †.—Very distinct by colouring of wings and tarsi and the median black stripe on abdomen ‡. Wing about 2.4 mm

♀ — *Head* dorsal surface covered with narrow white scales and white-tipped upright scales, latter chiefly towards nape, some deep brown flat scales at sides. Tori deep brown speckled with pale scales, flagellum of antenna brown. Clypeus deep brown with pale sheen. Palpi about $\frac{1}{2}$ length of proboscis, black, with white-scaled tips. Proboscis black at base, remainder yellow, speckled with black scales, labella light, apical $\frac{1}{3}$ of proboscis slightly enlarged. *Thorax* mesonotum covered with black and pale yellow or white narrow scales, latter arranged chiefly in a broad median band and in patches towards wing-roots. Scutellar scales black and light brown, intermixed, all fairly broad. Postnotum dark brown, lighter in middle and at sides. Integument of pleuræ black anteriorly, posterior part of sternopleura and mesepimeron very light. *Wings* speckled with pale yellow and black broad scales, many of which are asymmetrical, forks long, especially *af*. *Legs* femora dark on anterior surface, speckled with pale scales, paler posteriorly. Tibiæ speckled with black and pale yellowish scales. Tarsi yellow, with black markings, segment 1 on all legs with strong speckling of black scales in middle and usually a dark subapical ring, remainder yellow, segment 2

† Theobald 1907, p. 507, Edwards 1913 b, p 238 (syn.), Barraud 1929 a, p. 1057.


‡ [The only nearly related Oriental species, *F. elegans* Taylor (Queensland and Sumatra), differs in having abdomen all dark above and hind tarsi dark, with yellow rings over joints. The African *F. mediolineata* has a median pale line on the abdomen.]

yellow, with subbasal and subapical dark rings; 3, 4, and 5 usually with a single medial dark ring. *Abdomen*: dorsum with a median dark stripe running whole length, pale yellow scales laterally, the pale areas often speckled with black scales. Sternites mainly pale yellow.

♂—Markings as in ♀. Antennæ with dense plumes of brown hairs, arranged chiefly on two sides. Palpi slender, $\frac{2}{3}$ length of proboscis, slightly swollen at tip, actual tip white, part immediately behind this black. Proboscis considerably swollen on apical $\frac{1}{2}$.

[PUPA * (fig 24, a, b)—Respiratory horns extremely long, with annular thickening, except on the distal 6th, which is smooth and with slit-like opening, but not split as in

Fig 26



Larval characters of *Ficalbia*, subgenus *Etorleptomysa*:
a-c, *luzonensis*, d, e, *fusca*.

F. hybrida. No dendritic tuft on I. Lateral and sublateral hairs on II-VII very long and branched; sublateral pair on VII extremely long, lateral hair on VIII large, 8-10 branched. Paddles very long and narrow, with a dark spot at the tip, otherwise pale, short stout spines along both outer and inner edges, but stronger on outer, a fine terminal hair in a slight notch.

LARVA * (fig. 26, a-c).—Antennæ long, entirely pale, shaft spicular, a large tuft at middle, subapical hairs long, simple, portion of antenna beyond them only $\frac{1}{2}$ as long as middle section, one of the two apical hairs long, almost equalling

* [Senior-White 1925, p 220. Redescribed above from specimens from Ceylon (*Senior-White*) and Hong Kong (*Dr. R. B. Jackson*).]

the subapical pair *Frontal hairs* much as in *F. hybrida*. Preclypeal spines long and moderately stout, not blackened, but with a strong barb on each side near base, no spines on maxillæ. Meso- and metapleural plates each with a long spine. *Comb* of 10-12 teeth in one regular row. *Siphon* long and narrow, slightly widened before middle and considerably narrowed on distal $\frac{1}{4}$, when flattened appearing rather strongly curved, no pecten, a bifid hair at $\frac{1}{4}$. *Anal* segment very short, lateral hair long and single, as in *F. chamberlaini* and *F. hybrida*, very few hairs in fan.

HABITAT—Not definitely recorded, but presumably woody ponds on swamps with *Pistia*, as in the case of the allied African species.]

DISTRIBUTION—BENGAL Calcutta*, x 1910, 1 ♂ (*Shan ma*); Garden Reach*, viii 1931, 1 ♂ (*Senior-White*), U P. Saharanpur*, ix 1927, 1 ♀ (*Simton*). CEYLON Colombo*, 1913 (*James*), Kankawala village, Matale dist, larva of 1 ♀ (*Senior-White*).

Also known from MALAYA, BORNEO, SIAM, TONKIN, PHILIPPINE IS., and HONG KONG.

50. *Ficalbia* (*Etorleptiomyia*) *fusca* (Leicester), 1908.

Cul. Malaya, p. 102 (♂ & ♀) (*Dasymyia*). TYPE-LOC. Malay Penin., five miles from Kuala Lumpur (♂), Bukit Kutu (♀) (*Leicester*). TYPE. ♂ & ♀ in Brit Mus.

ADULT†—Small, obscure, dark species. Wing about 2 mm.

♀—**Head**: dorsal surface covered with flat dark purplish scales, a small patch of silvery scales on vertex and a narrow silvery line along eye-margins, a few small dark upright scales on nape. Antennæ dark brown, some silvery white scales on inner surface of torus. Clypeus, palpi, and proboscis brownish-black, palpi about $\frac{1}{3}$ length of proboscis, the latter slightly swollen on apical $\frac{1}{4}$. **Thorax**: mesonotal scales dark brown, bristles very long and numerous. Scutellar scales flat, small, and dark brown. Pleuræ dark brown, with small patches of silvery scales. **Wings**: scales comparatively large, dark in colour, many asymmetrical, as in *Mansonia*; fork-cells moderately long. **Legs**: dark brown, femora paler beneath, last two segments of hind tarsi distinctly pale, but not white. **Abdomen**: dorsum deep brown, with purplish sheen, small basal white lateral spots. Sternites pale yellowish.

♂.—Antennæ plumose, hairs dark brown. Palpi $\frac{1}{4}$ length of proboscis, very slender, and without hair-tufts. Proboscis much enlarged on apical $\frac{1}{4}$. Other details as in ♀.

† Edwards 1928 b, p. 57; Barraud 1929 a, p. 1057.

[PUPA † (fig. 24, c, d) —Respiratory horns extremely long and slender, basal $\frac{1}{2}$ blackened and densely covered with minute spicules (a unique feature of this species), distal $\frac{1}{2}$ bare and showing two false joints; below the horn a long bristle, and behind and above it a shorter stout spine-like bristle. No dentritic tuft on I; sublateral hair on VII not longer than on VI, lateral hair on VIII very large, with 12 or more branches. Paddles entirely pale, similar in form to *F. luzonensis*, but still narrower, almost linear.

LARVA † (fig. 26, d, e) —*Antennae* moderately long, pale, shaft with a few fine hairs but no spicules, tuft very large and well before middle; subapical hairs very long and branched in a dendritic manner, one of the apical hairs similar. *Frontal hairs* A, B, and C very large, plumose, A and C with about 25 branches, B with about 12. *Preclypeal spines* long, rather slender, simple, apical spine of maxilla short and weak. *Mentum* small and without median tooth. *Mesopleural plate* rather small, without spine, *metapleural* large, with 2 short spines. *Comb-teeth* about 30, in a triangular patch, those towards the base small. *Siphon* long and rather slender, not at all widened before middle, slightly narrowed and curved forwards on distal $\frac{1}{2}$, a bifid hair at $\frac{1}{2}$. Anal segment much as in *F. luzonensis*, *lh* may be double.

HABITAT —“Rot-hole in forest tree at ground-level, associated with *Heizmannia funerea*, the water was muddy, apparently from storm water”]

DISTRIBUTION —Only one ♂ known from Indian region ANDAMAN IS *, ix 1911 (*Christophers*). The specimen agrees exactly with Leicester's description.

Also known from MALAYA

51 *Ficalbia* (*Ficalbia*) *minima* (Theobald), 1901.

MC II, p. 262 (*Uranotaenia*) (♂) TYPE-LOC. Quilon, Travancore, III. 1900 (*James*) TYPE. 2 co-type ♂♂ in Brit. Mus.

Mimomyza minuta Theobald, 1908, Rec Ind Mus II, p. 301 (♂). TYPE-LOC. Sylhet, Assam, XI. 1904 (*Hall*) TYPE. ♂ in Ind. Mus. Theobald, *ib* 1910, IV, p. 30 (♀) TYPE-LOC.: Calcutta, VII. 1907 (*N. Annandale*) ALLOTYPE ♀ in Ind Mus

ADULT †.—A small species which might be mistaken for a *Uranotaenia*, but distinct on generic and subgeneric characters. Wing barely 2 mm.


♀—*Head* clothed for the most part with flat creamy scales (in some specimens appearing darker), some black

† [Edwards & Given 1928, p. 349. Redescribed above from the Singapore material.]

‡ Theobald 1903 a, p. 297, Edwards 1913 b, p. 238, Barraud 1929 a, p. 1057

upright scales on nape. Antennæ and proboscis brown, latter slightly enlarged towards tip, 1st flagellar segment long (fig 27, g). Proboscis as long as, or longer than, abdomen. Palpi very short, hardly exceeding clypeus. *Thorax* integument of mesonotum shining dark bluish, few dark scales, fairly numerous long black bristles. Postnotum deep brown. Pleuræ pale brown, with several small darker areas anteriorly. *Wings* veins clothed with broad dark scales, both forks fairly long. *Legs* femora mainly pale yellow, darker dorsally on apical $\frac{1}{2}$, knees of mid- and hind pair narrowly silvery

Fig 27



Ficalbia minima Thea. . a-f, larval characters (b, pecten-teeth, c, comb-teeth, d, preclypeal spines), g, base of antenna of ♀, h, proboscis and palp of ♂.

Tibiae and tarsi dark brown, with pale yellowish scaling over the joints between segments. *Abdomen*: dorsum dark brown, narrow basal creamy bands, widening out at sides in most specimens. Venter pale brown.

♂.—Very similar to ♀. Antennæ plumose. Palpi only slightly longer than in ♀. Proboscis very much swollen apically (fig. 27, h).

PUPA.—Unknown.

[LARVA † (fig. 27, a-f) — *Antennae* rather long but stout, shaft spicular, mostly white, but with a narrow blackish ring at base and another beyond the tuft, latter far beyond the middle, large, externally placed, with about 20 simple branches, subapical hairs both long, rather stout, simple, pale, one apical hair long (but shorter than the subapical), stout, blackened, portion of antenna between subapical and apical hairs short and stout. *Frontal hair* A with 8–10 plumose branches, moderately long, B with 2 strong plumose branches, C with 6–8 plumose branches, about as long as A, B and C nearly side by side and not far apart, d small, 2–3-branched, a little in front of C. Preelypeal spines stout and rather strongly barbed, especially on inner side. Mesopleural plate small, without spine, metapleural plate not much larger, with short, inconspicuous spine. *Comb* of about 6–8 teeth in one row. *Siphon* rather short, index 2.5–3, two strong simple pecten-teeth, tuft large and inserted near base. Valves large, unmodified. *Anal segment* without spines on posterior margin, *lh* rather short, 5–7-branched, papillae equal, short.]

DISTRIBUTION.—Type-localities, as given above, also BENGAL: Calcutta*, vii & viii 1907 (Annandale), xi 1910 (H. N. Sharma), viii 1911 (Graveley), Sonarpur*, x 1925 (M O T Iyengar), Dacca*, xi 1911 (S K S). ORISSA: Puri*, i 1911 (Annandale & Graveley), Ranihat*, i. 1922 (S Sundar Rao). ASSAM: Nalbari*, Kamrup dist, ix. 1928 (Sobha Ram), Golaghat*, Sibsagar dist, xii 1924 (Barraud).

Outside India has been recorded from BORNEO and HONG KONG

Genus **MANSONIA** Blanchard, 1901.

C R. Soc Biol. Ind, p 1045 Genotype, *Culex titillans* Walk

Taeniorhynchus L. Arnbalzaga (nec Weiland), 1891, Rev Mus La Plata, 1, p. 374 Genotype, *Culex titillans* Walk.

Panophites Theobald (nec Gould), 1901, M C n, p 173. Genotype, *Culex titillans* Walk

This genus is now divided into four subgenera—*Mansonia*, s str., *Rhyncotenia* Bréthes, *Coquillettia* Dyar, and *Mansonioides* Theo. The last two only are represented in India, the other two being purely American. There have been a number of changes in the classification of the genus, and for some years, up to 1930, it was generally known as *Taeniorhynchus* L. Arnib. I am in accord with the views of Edwards and others that this name is preoccupied by *Taeniorhynchus*

† [Described from specimens sent by Dr R. B Jackson from Hong Kong. The larva closely resembles that of the African *F. (Ingramia) malfeys* Newst. as figured by me in 1912.—F. W. EDWARDS.]

Weinland. The reader is referred to Edwards (1932, pp 114-120) * for synonymy and definitions of the four subgenera. The two subgenera included here may be recognized by the characters given below, as they show a number of structural differences they are more conveniently dealt with separately.


Subgenus **COQUILLETIDIA** Dyar, 1905

Proc Ent Soc Wash. vii, p 47 Genotype, *Culex perturbans* Walk

Chrysocoenops Theobald, 1907, M C iv, p 491 (in part)

ADULT —The three Indian species are yellow or yellowish-brown mosquitoes of moderate size, resembling some species of *Aedes* (*Aedimorphus*) in the form of the ♂ palpi, which are

Fig 28



♂ hypopygial structures of *Mansonia*, subgenus *Coquillettia* a, *crasipes*; b, *novochracea*, style, c, *M. (C.) ochracea* Theo Lettering as on p 4

slightly longer than the proboscis, the last two segments subequal in length, turned downwards, and bearing hairtufts. The general structure is very much as in *Aedes*, but there are no postspracular bristles (this appears to be the only constant character applicable to both sexes distinguishing the subgenus from *Aedes*). Distinct from *Culex* in absence of pulvilli. Tarsal claws of ♀ all simple, those of fore and mid-legs of ♂ unequal, the larger toothed, wing-scales narrow and lanceolate; pleurae with only a few small patches of scales.

* See also Barraud & Covell 1928, p 676 (buccal cavity).

PUPÆ of Indian species unknown, those of non-Indian species have the tip of the respiratory horn modified into a spine for piercing tissues of plants, no dendritic tuft on I, no bristles on following segments, paddles fairly long and narrow, emarginate at tips, no fringe or apical hair

LARVÆ of Indian species not known with certainty. A larva of probably *M. crassipes* or *M. ochracea* has been described by Edwards & Given (1928, p. 347). This and other non-Indian species of the subgenus have a very short siphon, the valves being modified for piercing tissues of plants, the part of the antenna beyond the subapical bristles is very long and flexible and has been said to have a prehensile function.

DISTRIBUTION and BIONOMICS.—Only one species is at all common in India and little is known of the habits, but the ♀♀ probably attack man. The larvæ and pupæ live in mud amongst the roots of aquatic plants and seldom come to the surface. About 27 species of the subgenus are known, and representatives occur in the Palearctic, Ethiopian, Oriental, and Australian regions, some extending to the Pacific Islands and New Zealand, one species is known from North America.

Key to Adults

- 1 Yellowish-brown; purplish scales on first 4 or 5 abdominal tergites, wing-scales mainly dark. *crassipes*, p. 120.
- Yellow or bright golden, abdomen yellow-scaled, wing-scales all yellow. 2
- 2 Integument of mesonotum and pleuræ uniformly pale. *ochracea*, p. 122.
- Integument of mesonotum with short brown stripes in front of each wing-base, pleuræ with brown stripes on upper part and a small brown marking below mesepimeron. *novochracea*, p. 121

52 *Mansonia (Coquillettidia) crassipes* (van der Wulp), 1892

- Bijl. Fauna Mid Sumatra, Dipt. p. 9 (♀) (*Culex*) TYPE-LOC. Soeroelangoen, Sumatra. TYPE ♀ in Leiden Mus.
- Tæmarhynchus brevicellulus* Theobald, 1901, M. C. II, p. 212 (♂ & ♀). TYPE-LOC. Selangor, Malay Penin. (*Wray*) (♀); Thayotmyo, Burma (*Watson*) (♂). TYPE ♂ & ♀ in Brit. Mus.
- Chrysocoenops pygmaeus* Theobald, 1908, Rec. Ind. Mus. II, p. 300 (♀). TYPE-LOC. Sylhet, Assam (*Hall*). TYPE ♀ in Ind. Mus.
- Theobald, *ib.* 1910, IV, p. 25 (♂). TYPE-LOC. Purnea, Bihar (*Pavani*). TYPE ♂ in Ind. Mus.
- Chrysocoenops fuscopteron* Theobald, 1911, Tijds. v. Ent. IV, p. 239 (♀). TYPE-LOC. New Guinea (*Dr. Koch*). TYPE ♀ in Amsterdam Mus.

ADULT*—Distinguished by its yellowish-brown thorax and dark wings and legs. Wing 4.5–5 mm.

* Leicester 1908, p. 163, Theobald 1910 b, p. 435, Edwards 1913 b, p. 230 (syn.), 1924, p. 365, and 1925 a, p. 260, Dyar 1925, p. 42, Barraud 1927 d, p. 556; Bonne-Wepster 1930, p. 396.

♀—*Head* covered with narrow and upright forked yellow scales, broader yellow scales at sides, bristles yellow, antenna, palpi, and proboscis brown, last with violet sheen, palpi $\frac{1}{4}$ – $\frac{1}{3}$ length of proboscis. *Thorax*. integument yellowish or brown, with sparse covering of narrow yellow scales and bristles, scutellar scales hair-like, yellow, usually sparso, integument of pleurae brown or dark yellow, a patch of silvery flat scales on upper part of mesepimeron, another on lower part of sternopleura. *Wings* dark scaled, a few yellow scales at extreme base in some specimens, membrane with pronounced purple, blue, and green reflections. *Legs*. femora yellowish at base, purplish on apical $\frac{1}{2}$, hind femur with some silvery scales on outer side in middle, tibiae and tarsi brown, with purple or brassy sheen. *Abdomen* first four or five tergites usually covered with purple scales, following segments yellow-scaled, but markings variable, some specimens having very few yellow scales on dorsum, in others they are much more numerous.

♂—Markings as in ♀, palpi longer than proboscis by about length of last segment, latter about $\frac{1}{2}$ length of penultimate segment, both moderately hairy and turned downwards. *Hypopygium* (fig. 28, a) shape of style characteristic.

PUPA—Unknown

LARVA not known with certainty (*vide* remarks above under subgenus).

DISTRIBUTION—Common in the PUNJAB and eastwards to ASSAM and BURMA, and southwards through PENINSULAR INDIA to CEYLON

The range extends through MALAYA to AUSTRALIA and FIJI

53 *Mansonia* (Coquillettia) *novochracea* (Barraud), 1927.

Ind Journ Med Res. xiv, p. 558 (♂) (*Taniorhynchus*). TYPE-LOC. Nongpoh, Khasi Hills dist, Assam, vii. 1932 (Barraud). TYPE ♂ in Brit Mus (Allotype ♀ from Margherita, Lakhimpur dist, Assam, v. 1920 (T. B. Fletcher), in M.S.I. coll Kasauli.)

ADULT.—A medium-sized bright yellow mosquito with some dark markings on thoracic integument*. Wing 4–4.5 mm or rather more

♀.—General coloration bright yellow, *head* covered with narrow and upright yellow scales, tori yellow, flagellum of antenna and hairs brown, palpi and proboscis yellow, latter dark at tip, palpi rather more than $\frac{1}{4}$ length of proboscis. *Thorax*. integument of mesonotum yellow, a short brown

* [The nearly related *M. gibli* Taylor (*conopas* Theo.), which is widely distributed from Malaya to Queensland, differs in having dark spots on posterior $\frac{1}{2}$ of mesonotum; lateral lobes of scutellum dark; hind femur without black ring at tip, etc.]

stripe laterally on each side in front of wing-roots, a fairly dense covering of narrow yellow scales and bristles, scutellar scales and bristles yellow, former hair-like and few; integument of pleuræ yellow, a dark stripe from the posterior margin of *ppn* to posterior margin of mesepimeron, another small dark area below lower margin of mesepimeron, very few pale flat scales. *Wings* entirely yellow scaled. *Legs*: yellow, with blackish-brown markings at tips of femora (on hind femur forming a narrow apical black ring), bases and apices of tibiae, and apices of tarsal segments, dark markings on tarsi of fore and mid-legs narrow, on hind legs wider, occupying fully $\frac{1}{2}$ of segments 2 and 3, femur and tibia of mid-leg speckled with brown scales. *Abdomen*: dorsum yellow scaled, small dark apical markings on tergites laterally, venter yellow.

♂—Markings as in ♀, antennal shaft and plume-hairs yellowish, segments distal to plumes brown, palpi longer than proboscis by nearly length of last segment, last two segments subequal in length, with tufts of yellowish brown hairs, scales brown except for yellow ring at base of penultimate segment, the long segment of palpi yellow scaled, with some brown scales and long yellow hairs at tip. *Hypopygium*. very similar to that of *M. crassipes*, but style of different form (fig 28, b).

PUPA and LARVA—Unknown.

DISTRIBUTION—Known only from the type-localities.

54. *Mansonia* (Coquillettia) *ochracea* (Theobald), 1903

M C III, p 263 (*Toemorrhynchus*) (♀) TYPE-LOC Kuala Lumpur, Malay Penin (*Durham*) TYPE ♀ in Brit Mus

Mansonia chrysogona Knab, 1909, Ent News Phil xx, p. 366 (♂ & ♀) TYPE-LOC Parang, Mindanao, P I, v 1908 TYPE ♂ & ♀ in U.S. Nat Mus

ADULT †—Very similar to *M. novochracea*, described above, but differs in absence of dark markings on integument of mesonotum and pleuræ and slightly in markings of legs of ♀, the fore and mid-tarsi being yellow or brown without any distinct banding, hind tarsi mainly brown, with yellow scaling at bases of first one or two segments, banding more distinct in ♂. Palpi in ♀ dark at tips. Style of ♂ hypopygium more elbowed than in *M. novochracea* (fig 28, c).

PUPA and LARVA—Unknown.

DISTRIBUTION—ASSAM. Gauhati *, Kamrup dist, v. 1918 and v 1920 (*T. B. Fletcher*), Margherita *, Lakhimpur dist,

† Leicester 1908, p. 164, Edwards 1922 d, pp 457 and 469 (syn.); Barraud 1927 d, p 559, Bonne-Wepster 1930, p 389

v 1920 (*T. B. Fletcher*), Cherrapunji, Khasi Hills dist,
 x 1920 (*Senior-White*) BENGAL Bossandhur, Khulna dist,
 viii 1919 (*J. T. Jenkins*)


Recorded also from MALAYA, SIAM, PHILIPPINES, and CHINA
 (Hangchow, Feng-Swen Li)

Subgenus *MANSONIODES* Theobald, 1907.

M.C iv, p 498 Genotype, *M. septemguttata* Theo

ADULT —Size moderate and build robust, general coloration yellowish-brown or black, legs with many pale markings and white rings on tarsi, wings speckled with light and dark broad scales, many of which are asymmetrical, palpi of ♂

Fig. 20




♂ hypopygial structures of *Mansonia*, subgenus *Mansoniodes*:
 a-c, *uniformis*; d, e, *indiana*, f-h, *annulifera* Lettering as
 on p. 4.

longer than proboscis, penultimate segment turned upwards, no distinct hair-tufts, terminal segment minute and turned slightly downwards; pleural bristles well developed, usually 10-20 postspiracular, 9-15 *ppn*, 12-18 upper mesepimeral, 4-9 towards lower part of mesepimeron; tergite VIII of ♀ with a number of chitinous hooks or teeth, the arrangement of which is of value in identification, lobes of sternite VIII

of characteristic shape in different species, two chitinised spermathecae Coxite of ♂ hypopygium short and wide, a large process arising from base on inner side, style of characteristic form, phallosome without lateral plates, a few small teeth at apex, paraprocts with a few teeth at crown and several minute hairs. Figures are given of the hypopygial details of the ♀♀ of all the Indian species. J. Bonne-Wepster (1930, pp. 197-212) first drew attention to the importance of the arrangement of chitinous hooks on tergite VIII of the ♀♀, it has been found also that the shape of the lobes of sternite VIII is of diagnostic value (Edwards, 1930, p. 541)

Fig 30



Tergite VIII (with teeth) and sternite (bare) of ♀, *Mansonia*, subgenus *Mansonioides* (flat preparation) a, *uniformis*, b, *indiana*, c, *annulifera*, d, *annulipes*.

An examination of these structures, and of the hypopygia of the ♂♂, has proved of great value in making correct identifications. This method should always be used in cases of doubt (see fig 30).


PUPA—Very similar to that of subgenus *Coquillettidia*, referred to above, tip of respiratory horn modified into a spine for piercing the tissues of plants; but one or two pairs of stout bristles present on hind margins of abd. seg. II-VII.

LARVA, 4th stage* (fig 31)—The larvae of three species have been isolated in India by Mr M. O. T. Iyengar at

* Ingram & Macfie 1917, p. 137.

Trivandrum, Travancore, and the skins of these and of pupæ, together with the corresponding adults, have been kindly loaned to me for description. The following characters are common to all three species, all are very similar in general structure.—Antennal shaft rigid and divided into three parts of about equal length by the hair-tuft and subapical bristles

Fig 31



Larval and pupal structures of *Mansonia*, subgenus *Mansonioides*: a-c, *indiana*; d, e, *annulifera*; f-h, *uniformis* (b, clypeus; c, h, respiratory horn with median ridge of thorax).

(in *Coquillettia* the apical part is very long and flexible), shaft with a number of small spines, subapical bristles long and reaching beyond end of antenna. Mouth-brushes fairly large, hairs simple. Preclypeal spines rather long and conspicuous. Frontal hairs A, B, C forming a concave row at about middle of clypeus, hair d directly in front of C, but much further forward, and anterior to level of A; C and d

all very small and difficult to see, *B* also small, *A* larger, and usually with 6-8 rather broad plumose branches. Mentum small, with comparatively large teeth, a large central one, with 4-5 smaller on each side. Hairs of thorax and abdomen only moderately developed. Comb usually of 2 very long blunt teeth. Siphon very short and conical, valves black and modified for piercing tissues of plants, and with 2 pairs of long spines, a single pair of branched hairs on tube near base of valves. Anal segment longer than wide in side view, as long as, or longer than, siphon, and completely ringed by plate. Both pairs of subdorsal hairs with a number of branches, these and fan-hairs all about same length. Fan-plate well developed. Usually 4 small branched hairs ventrally between fan-plate and base of segment. Lateral hair some distance from posterior margin and towards dorsum, of about 4 short, fine branches.

DISTRIBUTION and BIONOMICS *—Four species are known in the Indian area, two being common and widespread. The ♀♀ readily attack man, and are at times very troublesome. At least one species (*annulifera*) is a transmitter of *Filaria* (Iyengar, 1932) in India, and most probably other species are instrumental in the spread of this disease. About nine species in all are known, and are confined to the Old World, of two species found in Africa, one (*uniformis*) has an extensive distribution through the Oriental region to Japan and Australia; the remaining species are found in the Malayan and Australian regions. The larvæ and pupæ attach themselves to stems and rootlets of floating vegetation, and seldom come to the surface except when the pupa is ready for emergence. They should be sought for in ponds and swamps, particularly those much overgrown with vegetation.

Key to Adults

- | | | |
|---|----|------------------------------|
| 1 Mesonotum marked with distinct round spots of white scales | 2 | |
| Mesonotum not so (though there may be small indefinite white spots or patches) | 3. | |
| 2 Yellowish-brown, mesonotum marked with 4 (or more) distinct round white spots, rather broad white scales on mid-lobe of scutellum | | <i>annulifera</i> , p. 127. |
| Brownish-black, mesonotum marked with 2 (or 3) round white spots; scales on mid-lobe of scutellum narrow | | <i>longipalpis</i> , p. 128. |
| 3 Mesonotum marked with a pair of sublateral greenish stripes on a brown ground | | <i>uniformis</i> , p. 129. |
| Mesonotum dark brown, not marked with greenish stripes, some white scales, tending to form indistinct spots or patches, in some specimens | | <i>indiana</i> , p. 130. |

* Ingram & Macfie 1917, p. 137, Schwatz 1930, p. 311.

55 *Mansonia* (*Mansonioides*) *annulifera* (Theobald), 1901.

M C II, p. 183 (*Panophiles*) (♀). TYPE-LOC: Madras, XII 1899 (*Goodrich*), and Quilon, Travancore (*James*) TYPE ♀ in Brit. Mus.

Panophiles seguni Laveran, 1901, C R Soc Biol III, p. 991 TYPE-LOC Hanoi, Tonkin TYPE non-existent

Mansonioides septemguttata Theobald, 1907, M C IV, p. 499 (♀). TYPE-LOC Sarawak (*Barker*). TYPE ♀ in Brit Mus

ADULT *—Small yellowish-brown species. Wing about 4 mm.

♀—*Head*: some white scales along eye-margins and at sides, vertex and nape with narrow yellow and numerous pale upright scales, tori and clypeus pale brown, flagellum of antenna brown, with lighter rings, palpi about $\frac{1}{3}$ length of proboscis, yellow basally, broadly white apically, proboscis pale brown or yellow, usually darker at base and on apical $\frac{1}{3}$. *Thorax*: mesonotum with narrow yellow or golden scales, two conspicuous round white spots near front margin, another pair posterior to these, and three other spots, less distinct, at about level of wing-roots, scutellar scales white, broad on mid-lobe, narrow on lateral lobes, some narrow white scales on *ppn* and patches of broader white scales on pleurae. *Wings*: speckled with yellowish and dark brown, broad, asymmetrical scales, fairly evenly mixed. *Legs*: yellowish, marked with numerous snowy white rings, about 5 rings on each femur, knees also white, tibiae with 4-5 rings, tips also more or less distinctly white, segment 1 of tarsi of all legs with medial white ring, and basal ring also on mid- and hind legs and on 4 and 5 of hind legs. *Abdomen*: scaling on dorsum brown, or yellowish, and white, the last arranged chiefly in lateral patches towards apical border of each segment; lateral chitinous hooks on tergite VIII widely and evenly spaced (fig. 30, d), much as in *indiana*, but less curved, postgenital plate not divided to the base into two lobes as in *indiana*, the apex only being divided into two small lobes.

♂—Palpi $1\frac{1}{3}$ length of proboscis, yellowish, with two white rings on long segment, a white ring at base of penultimate, the small terminal segment entirely white. Scaling of head, body, wings, and legs as in ♀. *Hypopygium* (fig. 29, f-h): style divided nearly to the base into two arms, process arising from base of coxite, with 3 spines at apex.

PUPA.—Respiratory horn narrower than in *uniformis* or *indiana*, being 8-9 times length of greatest width; terminal spine about $\frac{1}{3}$ of total length. Posterior borders of II-IV with only one pair of long spines, V-VII with two pairs

* Leicester 1908, p. 174; Edwards 1913 b, p. 230 (syn.); Barraud 1927 d, p. 559; Bonne-Wepster 1930, p. 199.

of similar spines Paddles less emarginate at tips than in allied species

LARVA (fig 31, d, e) —Differs from those of allied species in having basal $\frac{1}{2}$ of antenna darkened (in *uniformis* and *indiana* there is a dark ring at base and another at level of origin of tuft)

DISTRIBUTION.—Common in BIHAR, ORISSA, BENGAL, ASSAM, BURMA, EAST and SOUTH-WEST COASTS of Indian Peninsula, and found in CEYLON Specimens have been examined from Fyzabad, U P, and from Central Provinces, but, from available records, the species does not appear to occur west of a line drawn from the western boundary of Nepal to Bombay

Recorded from MALAY PENINSULA, SIAM, PHILIPPINES, and MALAY ARCHIPELAGO

56 *Mansonia (Mansonioides) longipalpis* (van der Wulp), 1892

Bijl Fauna Mid Sumatra, Dipt p 9 (*Culex*) (♀) TYPE-LOC.: Sumatra Co-TYPE ♀♀ in Leiden Mus

Culex annulipes Walker (nec Meigen), 1857, Proc Linn Soc Lond 1, p 6 (♀) TYPE-LOC Singapore (A R Wallace) TYPE ♀ in Brit Mus

ADULT * —Differs from *M. annulifera* in larger size and darker coloration † Wing about 4.5 mm

♀ —**Head**. narrow white scales to eye-margins, vertex and nape with dark brown narrow and upright scales, palpi about $\frac{1}{3}$ length of proboscis, dark brown, with conspicuous whitetips, proboscis yellow on middle $\frac{1}{3}$, otherwise dark brown, with scattered yellow scales **Thorax** mesonotal scales dark brown, a pair of round white spots composed of broad scales towards front, another median white spot posteriorly of narrow white scales, some narrow white scales on front margin forming three patches, some broad white scales laterally in front of wing-roots, and sometimes some narrow white scales at sides more anteriorly, scutellar scales silvery white and all narrow, three patches of broad white scales on pleuræ, and both narrow and broad white scales on *ppm*. **Wings**: very similar to those of *M. annulifera*. **Legs**: dark brown, with numerous white rings having much the same arrangement as in *M. annulifera*, but white rings on femora less complete, except one preapical **Abdomen** dorsum dark brown or black, tergites I and II with a few yellow scales

* Leicester 1908, p 172, Edwards 1922 d, p 457, 1925 a, p 260, 1930 b, p 542, 1932, p. 120 (syn.), Barraud 1927 d, p 561; Bonne-Wepster 1930, p 206

† [*M. bonnea* Edw (Malay Peninsula and Archipelago) is closely allied, differing almost solely in hypopygium. The two species occur together in the Malayan region and, perhaps, also in India]

at base in middle; II, III, V, and VI with small lateral patches of white and yellow scales, IV with usually some white scales on apical border. Chitinous hooks on tergite VIII much as in *M. uniformis*, but there is a more pronounced gap between the lateral and median series, lobes of sternite VIII rather pointed.

♂—*Palpi* white-ringed as in *M. annulifera*, but ground-colour darker. Markings otherwise as in ♀ *Hypopygium* resembles that of *M. uniformis*, differing in form of phallosome, a pointed spine at tip of process of coxite.

PUPA and LARVA.—Unknown.

DISTRIBUTION (as checked by author from examination of hypopygia).—ASSAM Margherita, Lakimpur dist., v 1920 (*T. B. Fletcher*), Nongpoh, Khasi Hills dist., vii 192 and ix 1926 (*Barraud*), Cinamara, Jorhat, Sibsagar dist., 1932 (*D. Manson*). EASTERN BENGAL Chittagong Hill Tracts, Rangamati, viii 1922 (*Barraud*). Some previously published records have been found to be incorrect; others are considered doubtful.

Known from MALAY PENINSULA and ARCHIPELAGO and SIAM and recorded from PHILIPPINES, records from New Guinea and Australia are doubtful*.

57 *Mansonia (Mansonioides) uniformis* (Theobald), 1901.

M. C. II, p. 180 (*Panoplites*) (♀) TYPE-LOC Quilon, Travancore (*James*), and Taiping, Malay Penm. (*Wray*) CO-TYPE ♀♀ in Brit. Mus.

ADULT †—Brownish species, with less defined rings on femora than in the last two. Wing about 4.5 mm.

♀.—*Head*: vertex and nape with yellow decumbent and brown upright scales, those along eye-margins lighter, some broad pale scales low down at sides; tori yellow, flagellum of antenna brown, with faintly lighter rings, clypeus pale brown; palpi about $\frac{1}{2}$ length of proboscis, with scattered brown and yellow scales, those at tips lighter but not conspicuously white as in the two species described above, proboscis mainly yellow, with the apical $\frac{1}{2}$ dark brown, in many specimens narrowly dark at base also. *Thorax*: scales of mesonotum light brown and greenish, latter forming a pair of broad sublateral stripes from the front to above wing-roots; scutellar scales narrow and pale yellow; integument of pleuræ brown, several patches of broad white scales and some narrow pale scales on *ppn.* *Wings*: scaling very similar.

* [Such may refer to the allied *M. septempunctata* Theo., which has been confused with *M. longipalpis*.]

† Leicester 1908, p. 171, Edwards 1930 b, p. 542; Barraud 1927 d, p. 562; Bonne-Wepster 1930, p. 204.

to the two species described above. *Legs* four or five oblique pale markings on outer side of hind femur, similar but less definite markings on fore and mid-pair, pale tibial markings yellowish-tinged and run together, markings of tarsi as in *M. annulifera* but with yellowish tinge, not snow-white. *Abdomen*: dorsum dark brown, with lateral white and yellow patches and yellow apical bands in some specimens. Lateral chitinous hooks on tergite VIII curved and slightly separated from median series, lobes of sternito VIII larger and more rounded than in *M. indiana* (fig. 30, a).

♂—*Palpi* rather more than $1\frac{1}{2}$ times length of proboscis, brown, with two pale rings on basal $\frac{1}{2}$ and a narrower pale ring at base of penultimate segment, terminal segment minute and entirely pale, antenna about length of proboscis, with long brown plume-hairs, greenish stripes on mesonotum often less distinct than in ♀, otherwise markings are similar. *Hypopygium* (fig. 29, a-c) style wide, with pointed tip, a large process arising from base of coxite on inner side, ending in an appendage which is notched at tip.

PUPA (fig. 29, h)—Respiratory horn about 6 times length of greatest width, terminal spine between $\frac{1}{3}$ and $\frac{1}{4}$ total length. Hind margin of abd seg IV-VII with two pairs of strong spines, VII having also a smaller third pair, II-III with only one pair. The spines are usually longer than in *indiana*, one pair on tergite V reaching as far as hind border of VII. Paddle as in fig. 31, f.

LARVA (fig. 31, f, g).—Antenna with a dark ring at base and another at level of origin of hair-tuft. For other characters see under subgenus above.

HABITAT—Swamps and pools, especially those thickly overgrown with vegetation.

DISTRIBUTION—Common in most parts of India (including ASSAM, BURMA, PENINSULAR INDIA) and CEYLON, less common in the north-west, but specimens have been received from the United Provinces, Punjab, Smd, and Bombay. Often abundant and a troublesome blood-sucker.

The range of this species extends from AFRICA, through the ORIENTAL REGION, to JAPAN and AUSTRALIA.

58 *Mansonia (Mansonioides) indiana* Edwards, 1930*.

Bull. Ent. Res. xxi, p. 541 (♂ & ♀) TYPE-LOC.. Bandjaran, S Preanger, Java (*S. L. Brug*) TYPE ♂ in Brit. Mus.

ADULT.—Very similar to *M. uniformis* described above, but differs in the absence of greenish stripes on the mesonotum.

* See also Bonne-Wepster 1930, p. 210 (as *T. africanus* Theo.) *M. indiana* has previously been confused with *uniformis* or taken to be *africanus*, but the three are distinct on hypopygial structures.

the scaling of that part being usually almost uniformly dark brown, in some specimens there are some pale scales towards the lateral margins of mesonotum, and these may show up as faint light spots or areas in certain positions, but there are no definite round white spots as in *M. annulifera* and *M. longipalpis*. Chitinous hooks on tergite VIII of ♀ widely spaced and curved, without a definite gap between these and the median teeth (fig 30, b). Lobes of sternite VIII slightly emarginate and smaller than in *M. uniformis* (compare fig 30, a & b).

♂—*Hypopygium* (fig 29, d, e). style in side view (mounted flat) fairly wide, with slightly hooked tip, process arising from base of coxite moderately wide and long, with several hairs at tip, no definite appendage.

PUPA (fig 31, e).—Very similar to that of *uniformis* in shape of respiratory horn and in arrangement of bristles on abdominal segments, but these bristles usually shorter. Margin of head-capsule near base of antenna markedly crenulate, much more so than in *uniformis*.

LARVA (fig. 31, a, b).—No differences have been detected between the larvæ of this species and of *uniformis* in the material available. (For general structure see under subgenus above).

HABITAT.—As given for *uniformis*.

DISTRIBUTION (as checked by author by examination of hypopygia).—ASSAM: Nongpoh, Khasi Hills dist., and Dimapur, Sibsagar dist., vii. 1922 (*Barraud*), Gauhati, Kamrup dist., iii 1925 (*Barraud*), and x. 1928 (*C. S. Swaminath*). BENGAL: Calcutta, Garden Reach, iii 1931 (*Senior-White*). NILGIRI HILLS. Coonoor (no date). TRAVANCORE: Trivandrum, 1932 (*M. O. T. Iyengar*). BURMA (no date or locality).

Genus AËDOMYIA Theobald, 1901

M. C. II, p. 218 (*Aëdomyia*). Genotype, *A. squamipennis* L. Arriv.
Aëdomyia Edwards (amend.), Bull. Ent. Res. iii, 1912, p. 24.

ADULT* (fig. 33).—The single Indian species is a small mosquito with comparatively short legs, ornamented with a dense covering of white, black, and yellow scales, tufts of suberect scales at tips of mid- and hind femora. Antennæ about $\frac{2}{3}$ length of proboscis, those of ♀ with very short hairs, flagellar segments short, thick and rounded, and all about equal in length; those of ♂ plumose, last two segments thickened, but only slightly longer than others; [penultimate segment without whorl of long hairs at base, differing in this

* Edwards 1932, p. 121. See also Barraud 1927 a, p. 523; Barraud & Covell 1928, p. 676 (buccal cavity).

respect from all other genera of Culicini, as well as from *Anopheles* and *Megarhinus*] Palpi short in both sexes.

Other characters, as well as those of larva and pupa, and notes on distribution, are given below

59. *Aedomyia venustipes* (Skuse), 1889


Proc. Linn Soc N S W. (2nd series), iii, p 1761 (*Aedes*) (♀). TYPE-LOC. Elizabeth Bay, Sydney, Australia TYPE ♀ in Sydney Mus.

Aedeomyia catastricta Knab, 1909, Ent News, xx, p. 387 (♀) TYPE-LOC. Philippine Is TYPE ♀ in U.S Nat Mus

Aedeomyia squammipenna Theobald, 1901, M.C ii, p 210 (in part) (nec *Aedes squampennis* L Arnib)

ADULT * (fig. 33).—Fairly easily recognized by generic characters given above. Wing about 3 mm.

Fig 32.




Aedomyia venustipes Skuse. a & c, antennæ of ♂ and ♀, b, wing

♂♀.—Palpi rather less than $\frac{1}{2}$ length of proboscis, black-scaled, with white markings in middle and at tips and scattered

* Leicester 1908, p. 182 (*squammipenna*); Edwards 1924, p. 364 (syn.), 1929 b, p 326; Barraud 1927 a, p. 523

white scaling on upper surface; proboscis black, with three white rings. A broad median stripe of ochre-yellow scales on thorax, spreading out posteriorly in front of level of wing-roots; no spiracular or postspiracular bristles. *Wings*: densely clothed with dark brown, yellow, and white broad scales, the last arranged chiefly in three spots on costa and in smaller spots at forks of veins 2, 4, and 5. *Legs*: dark brown, speckled and spotted with yellowish and white scales, white basal and apical rings over the first two or three tarsal

Fig 33



Larval structures (portion of comb, siphon, anal segment, antenna)
of *Aedomya venustipes* Skuse

joints on fore and mid-legs and over all tarsal joints on hind legs, segment 5 being entirely white. *Abdomen*: dorsal surface mainly covered with ochre-yellow scales, patches of white scales at sides, venter brown, with many white scales, forming apical bands in some specimens.

♂—*Hypopygium**: coxite short, simple; style with comb-like appendage at tip.

PUPA.—Respiratory horn of moderate length, with wide opening, the sides of which are reticulated with fine lines;

* Edwards 1917, p. 229.

dendritic tufts on abd seg I small, one of the sublateral hairs on IV-VII long, with several shorter branches from base, lateral hairs on VII and VIII short, thick, and spine-like, with 2 or 3 branches, paddles moderately long and wide, with long terminal hair

LARVA * (fig 33).—*Antenna* with unusually long, wide, hairy shaft, with 3 long subplumose hairs at tip, tuft of about 10 subplumose hairs arising at about middle of shaft. *Thorax* with extremely long lateral, tufted, plumose hairs arising from chitinated plates, which bear thorn-like processes also Teeth of *comb* unusually long and bristle-like, in one rank along posterior margin of thinly chitinated plate, 10-15 on each side of segment *Siphon* short, whole surface setose, one pair of subposterior tufted hairs, with 5-6 extremely long branches, arising nearer apex than base, another pair, with shorter branches, on posterior valves, a pair of single hairs towards apex anteriorly, tip of siphon with a pair of strong curved spines, acus present, pecten absent *Anal segment* (fig. 33) with both *asc* and *osc* and fan-hairs plumose along one side

HABITAT—Weedy pools, more usually those containing much vegetation

DISTRIBUTION—Common in SOUTH INDIA, along the east coast of the Peninsula, in ORISSA, BIHAR, BENGAL, ASSAM, and BURMA, and is found in CEYLON It does not appear to occur west of a line drawn from the western boundary of Nepal to Bombay The author has seen single specimens from Bhopal, Central India, and Poona

Beyond the limits of the Indian region this species occurs throughout the Oriental region and in AUSTRALIA

Genus *ÆDES* Meigen, 1818

Syst Besch 1, p 13 Genotype, *A cinereus* Mg. (For Oriental synonymy see under subgenera.)

This genus, as at present constituted, includes more than 400 species, of which more than 100 are found in India, and it is certain that a further number await discovery. The Indian species are arranged in the following pages under subgenera, and a key is given to these. The adults vary much in size and ornamentation. As the group as a whole is an important one, and as difficulty may be experienced in deciding whether a particular specimen belongs to it or not, it may be useful to set forth the characters distinguishing *Ædes* (in the adult stage) from other genera represented in India

* Barraud 1923 h, p. 505 (*catasacta*).

The genus *Aedes* differs from *Uranotænia*, *Harpagomyia*, *Topomyia*, and *Hodgesia* in the venation of the wing (vein 6 ending nearer apex of wing than level of fork of 5), and in presence of fringe of hairs on squama of wing, from *Theobaldia* and *Tripteroides* in the absence of spiracular bristles and presence of postspiracular bristles, from *Theobaldia* also in absence of hairs from underside of wing at base, from *Culex*, *Mansonia* (*Coquillettidia*), *Ficalbia*, and *Aedomyia* in presence of postspiracular bristles, in addition, from *Culex* in absence of pulvilli, from *Ficalbia* in not having the proboscis enlarged at tip, and from *Aedomyia* in structure of antennæ, from *Heizmannia* and *Hæmagogus* in form of *apn* lobes (these being normal in *Aedes* and not enlarged or approximated behind the head), from *Heizmannia* also in the postnotum being without hairs, from *Orthopodomyia* in the relative lengths of tarsal segments, segment 4 on front legs of ♀ not being shortened, from *Mansonia* (*Mansonioides*) in wing-scaling (in *Aedes* the scales are never very large and asymmetrical) and in the structure of the hypopygium in both sexes. As regards the remaining genus, *Armigeres*, which is evidently very closely allied to *Aedes* (subgenus *Stegomyia*), but retained as a separate genus chiefly on account of the structure of the larvæ, the only difference in the adults appears to be in form of the proboscis, which in *Armigeres* is somewhat curved downwards towards the tip, the curved part being slightly compressed laterally. In *Aedes* the proboscis is straight (but tends to curve upwards in dried specimens), and is not laterally compressed.

The palpi of the ♂♂ are of variable length and form, those of the ♀♀ not usually more than $\frac{1}{4}$ length of proboscis. Antennæ of ♂ nearly always distinctly plumose, with the last two segments elongate; those of ♀ with moderately long hairs and all flagellar segments of about equal length. Bristles on mesonotum usually well developed. Structure of hypopygium of importance in both sexes in differentiating subgenera and species (see keys and under subgenera).

PUPA.—Respiratory trumpet usually fairly short, with round, slightly oblique opening. Dendritic tufts on abdominal segment I usually well developed; hairs on posterior margins of II–VII nearly always fine and inconspicuous, either branched or single, not bristle-like; rather small branched hairs on outer posterior corners of VII–VIII. Paddles usually fairly wide and rounded or slightly elongate, hind margin not markedly serrated and usually without conspicuous fringe, a small terminal hair, which may be branched. This stage of the Indian species has been very little studied at present. So far as is known the subgeneric and specific differences are slight.

LARVA.—Mouth-parts not modified for predaceous habits except in subgenus *Mucidus*. Median hairs of mouth-brushes in many species slightly hooked, and with more or less obvious serrations along one side apically. Arrangement and form of the four important pairs of frontal hairs very variable and of importance in identification. Thorax with moderately developed hairs and, in a few species, with large spines, or thick barbed hairs, on dorsum. Segments I-VIII of abdomen without chitinous plates, except in a few species of subgenus *Stegomyia*, in which there are lateral plates on VIII. Comb of teeth present on VIII, variable in number and arrangement. Siphon usually $2-3\frac{1}{2}$ times length of diameter at base, but longer in a few species, only one pair of hair-tufts on tube, towards posterior border. Pecten present, variable in number and shape of individual teeth. Anal segment with chitinous saddle, rarely enclosed in chitinous ring. Lateral hair usually fairly long. Outer subdorsal hairs single and long, inner pair shorter and usually divided into 2 or more branches. Anal fan moderately developed or rather small.

EGG.—With thick shell, spindle-shaped or elliptical, often with fine sculpturing on surface.

DISTRIBUTION and BIONOMICS (see remarks under subgenera).—Eggs, so far as known, laid singly, very resistant to desiccation. Several species are known to be capable of transmitting disease. Many are day-flying, and ♀♀ are active blood-suckers. Larval habitat very variable in different subgenera.

Keys to Subgenera

♀♀

- | | | |
|--|---|----------------------|
| 1. Segment VIII narrow and completely retractile, cerci long and narrow and projecting from ring of segment VII | 2 | |
| Segment VIII broader and not completely retractile, cerci shorter and broader (but sometimes rather long in subgenus <i>Aedes</i>) | 6 | |
| 2. Large yellow, white, and brown mosquitoes with outstanding scales on body and legs, giving them a shaggy or mouldy appearance, wing-membranes clouded in region of c - vs. 2-3, 3-4, 4-5, these three being almost in a straight line | | MUCIDUS, p. 144 |
| Smaller mosquitoes, not of shaggy appearance; wing-membranes not clouded | 3 | |
| 3. Dark species, with conspicuous lateral yellow scaling on mesonotum | | BANKSINELLA, p. 269. |
| Ornamentation otherwise | 4 | |
| 4. First hind tarsal segment as long as tibia, dark brown species, without ornamentation | | RHINOSKUSEA, p. 216. |

- First hind tarsal segment shorter than tibia, as usual, body and legs usually ornamented 5 [p 246
 5. No lower mesepimeral bristles . . . AEDIMORPHUS (part).
 One or two lower mesepimeral bristles usually present OCHLEROTATUS, p 147.
 6 Tarsal claws toothed, at least on fore and mid-legs 7
 Tarsal claws all simple 11
 7 Sternite VIII large and prominent, ornamentation various FINLAYA, p 153
 Sternite VIII smaller and less prominent; dorsal surface of head with many flat scales 8
 8 Scutellar scales all narrow, dark species, with little ornamentation AEDES (part), p. 277.
 Scutellar scales broad and flat, at least on mid-lobe, species with conspicuous ornamentation 9.
 9 Scutellar scales broad and flat on mid-lobe, narrow on lateral lobes [p 265.
 Scutellar scales broad and flat on all lobes. A (*Aedim*) *nummatus*,
 10 Proboscis dark, apn lobes with broad flat scales 10 [p 217.
 Proboscis with a white ring, apn with narrow scales only STEGOMYIA (part),
 11 Black species, with snow-white markings and white basal bands on tarsi [p 212.
 Species with little or no ornamentation CHRISTOPHERSIOMYIA,
 12 Scutellar scales all narrow [p 217.
 Scutellar scales broad and flat STEGOMYIA (part),
 13 First hind tarsal segment as long as tibia, proboscis unusually long 12
 First hind tarsal segment shorter than tibia, as usual, proboscis not unusually long 13
 14 Head with white stripe either side of middle line AEDES (part), p 277
 Head all dark dorsally 13
 RHINOSKEUSA, p 216.
 14
 DICEBOMYIA, p 271
 CANCRAEDES, p 297.

♂♂ (as far as known)

1. Palpi at least $\frac{1}{2}$ length of proboscis, usually longer 2.
 Palpi short, not more than $\frac{1}{2}$ length of proboscis 9.
 2 Palpi about $\frac{1}{2}$ length of proboscis, thin, and without hair-tufts A (*Steg*) *albolineatus*.
 Palpi more than $\frac{1}{2}$ length of proboscis, with or without hair tufts. [p 243, A (*Aedim*)
 3 Phallosome divided into lateral plates, bearing chitinous teeth 3 [*nummatus*, p 265
 Phallosome not divided into lateral plates, tube-like or scoop-shaped and without chitinous teeth 4.
 4 Palpi of only 2 distinct segments, the 2nd upturned and hairy; dark species, with conspicuous lateral yellow scaling on mesonotum 7.
 Palpi of 3 distinct segments, ornamentation otherwise BANKSINELLA, p 269.
 5

- 5 Palpi slender, upturned, without distinct hair-tufts, black species, with conspicuous snow-white ornamentation STEGOMYIA, p. 217
 Palpi not upturned and with hair-tufts, ornamentation otherwise 6.
- 6 Scales on *ppn* broad and flat, lower mesepimeral bristles present (1-3) (except in *periskeletus*) DICEROMYIA, p. 271.
 Scales on *ppn* narrow, no lower mesepimeral bristles AEDIMORPHUS, p. 246
- 7 Large yellow, white, and brown mosquitoes with outstanding scales on body and legs, giving them a shaggy and mouldy appearance, wing-membrane clouded in region of c - vs 2-3, 3-4, and 4-5, these three being almost in a straight line MUCIDUS, p. 144
 Smaller mosquitoes, not of shaggy or mouldy appearance, wing-membrane not clouded. 8
- 8 Coxite with projecting basal lobe carrying a long spine, and sometimes an apical lobe also OCHLEROTATUS, p. 147
 Coxite without such lobes (if small basal lobe present there is no long, differentiated spine) FINLAYA, p. 153.
- 9 Ornate species with pale band on proboscis CHRISTOPHERSIOMYIA,
 No marked ornamentation, proboscis without a pale band [p. 212.]
- 10 Proboscis very long, segment 1 of hind tarsi as long as tibia 10
 Proboscis not unusually long, segment 1 of hind tarsi shorter than tibia RHINOSKUSEA, p. 216
- 11 Scutellar scales all flat 11
 Scutellar scales all narrow CANORAÆDES, p. 297.
 AÆDES, p. 277

Key to Larvæ (4th stage).

As *Aedes* larvæ do not possess well-marked subgeneric characters, one key to all the known larvæ is given. [The known larvæ of *Heizmannia* are also included, as this genus does not differ obviously from *Aedes* in the larval stage]

- 1 Mouth-brushes adapted for predaceous habits, both *oso* and *isc* hairs single (unbranched), anal fan of close-set branched hairs extending along whole length of anal segment ventrally [p. 146.
 Mouth-brushes not adapted for predaceous habits, *isc* usually of 2-10 branches, but, if both pairs are single, anal fan does not extend along whole of ventral surface of segment A (*M.*) *scatophagoides*, 2.
2. Abd seg VIII with a semicircular chitinous plate on each side, from posterior margin of which the comb-teeth arise 3.
 Abd seg. VIII without lateral chitinous plates 5
- 3 Comb of 3 teeth with strong basal lateral denticles A (*S*) *desmotes*, p. 227
 Comb of 5 teeth without basal denticles. 4

4. Siphonal hair-tuft of 2-3 fine branches,
 1/4 of 2 subplumose branches, pecten-
 teeth without denticles Δ (S.) *annandalei*, [p 220.
 Siphonal hair-tuft of 3 subplumose
 branches, 1/4 of 2 simple branches,
 pecten-teeth with lateral denticles Δ (S.) *mediopunctatus*, [p 231
- 5 Thorax with 2 pairs of large, single, black
 spines on dorsum, comb-teeth in a patch,
 siphon normal 6
 Thorax without such spines 7
- 6 Siphonal hair-tuft modified into a stout
 spinulose bristle, which may be bifid Δ. (P') *elsae*, p 181.
 Siphonal hair-tuft of 3-4 subplumose, or
 barbed, branches Δ (P') *shortii*, p 184
- 7 Siphon with a ring of spines near apex and
 a median patch of similar spines on
 anterior surface Δ (Aedim) *cæcus*, [p 258.
 Siphon without such spines 8
- 8 Comb of 5-10 large teeth in a single row,
 or of 6-14 teeth in an irregular row, or
 more or less in 2 rows 9.
 Comb of 20-70 teeth, usually small,
 arranged in several rows, or more or less
 in a triangular patch 33
9. Antennal shaft with small spicules or
 spines 10
 Antennal shaft smooth 22
- 10 The 4-5 pecten teeth nearest base of siphon
 simple and larger than more distal teeth,
 which have lateral denticles Δ (F') *novonivensis*, [p 212.
 All pecten-teeth simple *Heizmannia funerea*,
 Basal pecten-teeth with lateral denticles,
 and smaller, or at least not larger than,
 more distal teeth 11 [p 302.
- 11 The 2-3 pecten teeth furthest from base of
 siphon more widely spaced than those
 nearer base, and usually distinctly larger
 than basal teeth 12.
 All pecten-teeth evenly spaced and all
 about same size, those furthest from base
 being but little larger than remainder 16.
- 12 Frontal hairs B and C both with several
 branches 13,
 Frontal hair B single (unbranched) 15
- 13 All pecten-teeth with lateral denticles,
 except most distal tooth in some in-
 dividuals 13
 The most distal 2-3 pecten-teeth without
 lateral denticles Δ (Aedim) *varans**, [p 254.
14. Antennal shaft with few minute spicules,
 and tuft of 3 fairly stout branches,
 nearer base than apex, comb usually
 of 10 teeth, pecten-teeth with long
 sharp point, frontal hairs B and C each
 usually 3-branched Δ. (Aedes) *indicus*, [p. 284.

* As figured by Howard, Dyar, and Knab for *Aedes sylvestris*

- Antennal shaft with fairly numerous strong spinelets, and with tuft of about 5 branches at middle, comb usually of 6-8 teeth, pecten-teeth smaller, frontal hairs *B* and *C* each with usually 5 or more branches [p. 270.
A. (B) lineatopenus,
 15 Frontal hair *C* with 3-5 branches, siphon [p. 257
 2-2½ times length of diameter at base *A. (Aedim) syntheticus*,
 Frontal hair *C* single (unbranched), [p. 254.
 siphon 2½-3 times length of diameter at base *A. (Aedim.) vexans**,
 16 Frontal hairs *A*, *B*, *C*, and *d* all well 17
 developed, and each with a number of 21
 fairly long branches
 Hair *d* much smaller than other 3 pairs,
 and with few short fine branches
 17 Comb of 8-14 teeth arranged irregularly, 18
 more or less in 2 rows, siphonal tuft of 19
 about 3 fairly long branches at ¼ length 20
 of tube from base, pecten of 16-21 21
 teeth
 Comb similar, but pecten of 5-8 teeth
 Comb of 8-19 teeth arranged in 1 close-
 set row, siphonal tuft of 4-6 branches at
 about middle of tube 18
 18 Comb of 16-19 teeth, each with fringe [form), p. 209.
 from base to apex, siphon about 2½ *A. (F.) nveus* (typo-
 times length of diameter at base . . .
 Comb of 8-12 teeth, each fringed not more
 than half-way from base to apex; siphon
 3-5 times length of diameter at base . . 19.
 19 Siphon about 5 times length of diameter [p. 211.
 at base *A. (F.) nveoulos*,
 Siphon not more than about 3½ times 20
 length of diameter at base
 20 Comb-teeth fringed for about ¼ length
 from base to tip, longer pecten-teeth
 about ¾ length of larger comb-teeth,
 siphon about 1 mm. long
 Comb-teeth fringed at base only, longer
 pecten-teeth about ¾ length of larger
 comb-teeth; siphon about 0.7-0.8 mm
 long.
 21 Comb of about 14 teeth in a single row, [p. 210.
 i.e. 2-branched; pecten of about 16 com- *A. (F) albonveus*,
 paratively small short teeth [p. 206.
 Comb of about 15-20 teeth irregularly *A. (F) albolateralis*,
 arranged, i.e. of about 12 branches, [p. 214
 pecten of about 20 comparatively large *A. (C.) thomsoni*,
 and long teeth
 22 The most distal pecten-tooth widely spaced *A. (O) caspius*, p. 149.
 from main rank and attached between
 hair-tuft and apex of siphon *A. (S) vittatus*, p. 245.
 Pecten-teeth all in one rank, none widely
 spaced 23.
 23 Comb-teeth with strong basal lateral
 denticles *A. (S) aegypti*, p. 223

* Specimens sent by Dr. P. A. Buxton from Samoa

- Comb-teeth without basal lateral denticles, but usually finely fringed 24
- 24 Frontal hair *C* many-branched, *B* single, comb-teeth in a regular row, pecten with only 5-8 teeth, *isc* 8-10-branched [p 244.
A. (S.) albolineatus,
[*ensis*, p 307.
Heizmannia himalay-
- Frontal hairs *B* and *C* each 3-branched, comb-teeth irregular, pecten with only 4-5 teeth 25.
- Frontal hair *C* single, *B* single or bifid, pecten-teeth more numerous
- 25 Comb of 6-10 teeth in an irregular row; pecten of about 18 teeth in a close-set rank, antennal hair of 3-4 branches *A. (O) pulchritarsis*,
[p 151
- Comb-teeth in a single regular row, pecten usually of less than 14 teeth, antennal hair usually single 26
- 26 Pecten-teeth with lateral denticles along one side from base to apex 27
- Pecten-teeth with basal lateral denticles only, *isc* with at most 3 branches, usually single 30
- 27 *isc* of 3-6 branches 28
- isc* usually single, sometimes split into two [p. 237.
A (S) pseudalbopictus,
- 28 Mesopleural tubercle with short spine, siphonal tuft represented by a single barbed hair at about middle of tube *A (D) uyengari*, p 275.
- Mesopleural tubercle with long spine, siphonal tuft usually of 2 branches, nearer apex than base of tube 29 [p 276
- 29 *isc* of 3 branches *A (D) macropterus*,
- isc* of 6 branches *A (D) reginae*, p. 277
- 30 Siphon about 3 times length of diameter at base *A (S.) novalbopictus*
[p 237, *A (S)*
- Siphon 2-2½ times length of diameter at base 31 [p 239
[*flavopictus*,
[p 239
- 31 No definite fan-plate on anal segment, a clear area in chitinisation around base of *lh* [p 242.
A (S) unilineatus,
- A definite fan-plate on anal segment, no clear area in chitinisation around base of *lh* 32.
- 32 Antennal shaft about 6 times length of greatest width [p 241
A. (S) scutellaris,
- Antennal shaft about 10 times length of greatest width [p 235.
A. (S) albopictus,
33. One or two pairs of frontal hairs single and very long, longer than whole head 34
- None of the frontal hairs longer than whole head 36.
- 34 Frontal hairs *B* and *C* both single and very long *A (F) gilli*, p 198.
- Frontal hair *B* very long and single, *C* much shorter and divided into several branches. 35
- 35 Dorsal pair of anal papillæ only about length of anal segment, ventral pair shorter, both pairs pointed *A (F) greeni*, p. 185
[p 195.
- Dorsal pair of anal papillæ longer than anal segment, the ends rounder *A (F) christophersi*,


- 36 Several of the more distal pecten-teeth bare, without lateral denticles or fringe, and usually larger than the more basally situated teeth . . . 37
 All pecten-teeth with lateral denticles or fringed, and usually all about same size . . . 42
- 37 Pecten with 2-3 large simple distal teeth markedly detached from the more basal teeth, pecten confined to about basal $\frac{1}{4}$ of tube . . . 38
 Pecten either without markedly detached distal teeth or, if with such teeth, the pecten extends for much more than basal $\frac{1}{4}$ of siphon . . . 39
- 38 Siphon very long, 6-7 times length of diameter at base, antennal tuft of about 20 branches, pecten of 9-13 teeth . . . [striatus, p 262]
A. (Aedon) pallido-
 Siphon about 3 times length of diameter at base, antennal tuft of 8-10 branches, pecten of 14-19 teeth . . . [calatus, p 260]
A. (Aedon) peper-
- 39 Dorsal-lateral hair [hair 7] of metathorax with 3-4 strong, almost spine-like and barbed branches, 4-5 simple pecten-teeth between hair-tuft and apex of siphon, last 2-3 very large . . . *A. (F) saricola*, p 101
 Dorsal-lateral hair of metathorax not unusually developed, at most 3 simple pecten-teeth between hair-tuft and apex of siphon . . . 40.
- 40 Pecten of 16-19 teeth, two simple ones beyond tuft, *lh* 3-4-branched . . . [p 187.
A. (F) chrysolineatus,
 Pecten of about 10 teeth, *lh* single or double . . . 41.
- 41 Siphon shorter, one or two simple pecten-teeth beyond tuft . . . [p 190
A. (F) formosensis,
 Siphon longer, no pecten teeth-beyond tuft . . . *A. (F) harveyi*, p. 189
- 42 Siphonal tuft very near apex of siphon and of 3 long branches, antennal tuft of 3-4 branches, nearer base than apex . . . [p 210.
A. (R) longirostris,
 Siphonal tuft not usually more than $\frac{2}{3}$ of length from base of tube, antennal tuft usually at about middle of shaft or nearer apex than this . . . 43
- 43 Frontal hair *d* unusually well developed and as large as *A*, *B*, and *C*, these four hairs each with a number of branches, pecten-teeth fringed on both sides and at tip, as in *Uranoctonus* . . . [p. 173.
A. (F) albocinctus. . . 44
- 44 Pecten of about 12 teeth, either unusually long and slender, and fringed along one side, or unusually short and stout, with one strong basal denticle . . . 45
 Pecten-teeth nearly always more than 12 in number, and not formed as above . . . 47.
- 45 Pecten-teeth long and slender and fringed along one side from base to apex . . . 46.

- Pecten-teeth unusually short and broad,
with one strong basal denticle, antennal
shaft smooth [p 171.
A. (F) uncinatus,
- 46 Antennae long and blackish, no obvious
stellate hairs on thorax or abdomen,
margin of anal saddle almost smooth [p 206
A. (Aedim) nummatus,
- Antennae short and pale, conspicuous
many-branched stellate hairs on thorax
and abdomen, margin of anal saddle very
spiny. *A. (F) poecilus*, p 158
- 47 Frontal hairs *B*, *C*, and *d* all placed well
forward towards front margin of clypeus,
and standing more or less in transverse
line, and each with 3 or more branches 48
Frontal hairs *B*, *C*, and *d* placed further
back towards middle of clypeus, *B* and *C*
usually single and standing one in front
of the other 50
- 48 Antennal shaft with small spicules, *asc*
usually 4-branched [p 200
A. (F) pulchiventer,
- Antennal shaft smooth, *asc* usually 2-
branched 49
- 49 Abdominal segments with stellate hairs,
comb of 50-60 teeth [p 177
A. (F) macdougalli,
- Abdominal segments without stellate hairs,
comb of 40-45 teeth [p 179
A. (F) pseudotarnatus,
- 50 Frontal hairs *B*, *C*, and *d*, all close
together, antennal shaft comparatively
long [p 204
A. (F) dissimilis,
- Frontal hairs *B* and *C* well separated,
antennal shaft not so long 51
51. Frontal hair *B* with about 4 branches
(compare also *A. (O) pullatus*, p 152). [p 192
A. (F) oreophilus,
- Frontal hair *B* single 52
- 52 Antennal shaft with small spines, antennal
hair with 4 or more branches
Antennal shaft smooth, antennal hair
usually single 53
- 53 Preclypeal spines comparatively stout and
blunt-ended 54
Preclypeal spines inconspicuous, rather
long, and tapering to a fine point 57
- 54 Frontal hair *d* nearer to *C* than to *B* 55
Frontal hair *d* nearer to *B* than to *C*, or
the 3 hairs placed one behind the other
at fairly even wide intervals 56
- 55 *asc* of about 8 fairly long subequal branches
asc of 2 long branches *A. (P) klazani*, p 168
A. (F) assamensis,
- 56 Frontal hair *d* near to *B* than to *C*, *asc*
of 3-4 fairly short branches [p 160
A. (F) decanus, p. 103
- Frontal hairs *B*, *C*, and *d* placed one
behind the other at fairly even, wide
intervals, *asc* of 2-6 long fine branches [p. 165.
A. (F) feegrader,
57. Median hairs of mouth-brushes with very
small serrations; antennal shaft from
0.41-0.51 mm long [(type-form), p. 161
A. (F) gubernatoris
- Median hairs of mouth-brushes with com-
paratively very large serrations, an-
tennal shaft from 0.32-0.35 mm long [p. 161
A. (F) kotensis,
A. (F) gubernatoris

Subgenus **MUCIDUS** Theobald, 1901.M.C. 1, p 268 Genotype, *Culex alternans* Westwd.*Pardomyia* Theobald, 1907, M C 1v, p 280. Genotype, *P. ausanilla* Theo*Ekrinomyia* Leicester, 1908, Cul Malaya, p 71 Genotype, *E. aureostrata* Leic

ADULTS*.—Large mosquitoes, with outstanding yellow, white, and brown scales on body and legs, giving them

Fig 34.



Larval and adult structures of *Aedes*, subgenera *Mucidus* and *Ochlotatus* a-g, *A. (M.) scatophagoides* (a, wing of ♀, b-f, larval details, e, pecten-teeth, f, comb-teeth, g, harpago of ♂), h, *A. (M.) laniger*, harpago of ♂, i-k, *A. (O.) caspius*, ♂ coxite, with appendages, harpago, and proctiger in side view

* Barraud 1929 a, p 1052, Edwards 1932, p. 133; Barraud & Covell 1928, p 676 (buccal cavity)

a mottled and mouldy appearance. Some of the white scales on mesonotum remarkably long and twisted, resembling strands of cotton-wool*. Wings mottled with yellow, brown, and creamy scales, membrane usually slightly clouded in region of cvs 2-3, 3-4, and 4-5, these three being approximated and almost in line, though 4-5 may be slightly nearer tip of wing than the other two. *ppn* bristles numerous, usually about 20.

Hypopygium in both sexes as in *Ochlerotatus*.

LARVA.—Mouth-brushes forming strong, matted tufts, the hairs slightly reduced in number (50-70) and thickened, but not specially modified, serrations of inner margin about as strong as in many other species of the genus. Both *isc* and *osc* single. Anal fan of close-set branched hairs extending along whole length of segment ventrally (an unusual character in Culicine larvæ), compare the similar condition in the predaceous subgenus *Lutzia* of *Culex*. Frontal hairs all single, their arrangement unusual, *d* being apparently considerably external to *B* and *C*.

DISTRIBUTION and BIONOMICS.—Only about seven species are known, and these are confined to the tropics of the Old World, from West Africa to Melanesia. Two are known in India, one having a very wide distribution, the other being rare. The ♀ usually carries the abdomen curved forwards beneath the thorax. The larvæ live in open natural pools and are predaceous on other mosquito larvæ.

60 *Aedes* (*Mucidus*) *scatophagoides* Theobald, 1901.

M C 1, p. 277. TYPE-LOC.: Myingyan, Burma (*E. N. Watson*), and Moradabad, N.W.P. (*Major Close*). TYPE: 2 ♀♀ co-types in Brit Mus.

ADULT ♂.—A large species, distinguished on subgeneric characters given above, and by leg-markings and relatively short hind tarsi. Wing 5.5-6.5 mm.

♀.—Proboscis mainly yellow, with a variable amount of white scaling in middle. Palpi rather more than $\frac{1}{4}$ length of proboscis, shaggily clothed with dark brown, yellow, and white scales. Thorax with light matt grey integument, mesonotal scales light brown and white, latter in long tufts. Costa of wing mainly yellow in some specimens, in others

* [The peculiarities of scaling mentioned above are found in the two Indian species, but the third Oriental species (*A. aurantius* Theo., widely spread from Malaya to Queensland, and quite possibly occurring in Burma) has normal scaling of the head, body, and legs, and very different colouring. It is a dark brown mosquito with tip of abdomen broadly golden; hind tarsi dark with narrow yellow rings and fifth segment conspicuously whitish; wings nearly bare.]

† Edwards 1911, p. 246, and 1922 *d.*, p. 450; Barraud 1929 *a.*, p. 1053.

dark scaled from base for some distance, but there is usually a large yellow streak towards apex of wing extending on to vein 1. *Legs* brown, with numerous white rings, including three on each tibia (at base, middle, and tip) and two on the first segment of each tarsus (at base and middle), segments 2-5 of hind tarsus each with a white basal ring. First hind tarsal segment barely $\frac{3}{4}$ length of tibia. *Abdomen* with brown and white scales, latter predominating, especially towards tip.

♂—Resembles ♀ except that antennæ are densely plumose, the hairs being yellow, and arising chiefly from two sides of the shaft, not in regular whorls. Palpi longer than proboscis by about length of last two segments, which are enlarged and spatulate, apical $\frac{1}{2}$ of palp with dense hair-tufts. Mesonotal scales often lighter than in ♀, sometimes almost entirely white. *Hypopygium* (fig 34) coxite moderately long, with basal lobe bearing a number of hairs. Style fairly long, with rather short terminal appendage. Harpago well developed, with curved blade about the length of stem. Phallosome scoop-shaped, without lateral plates or teeth.

LARVA * (fig 34)—*Head* comparatively large. Mouth-brushes adapted for seizing prey, and consisting of a number of slightly hooked lamellæ, with serrations along one side apically. Frontal hairs all simple and inconspicuous, arranged as shown in fig 34, d. Antenna comparatively short, shaft smooth, hair-tuft of 3-4 branches at only a short distance from tip. Mandibles very strong, teeth black. Mentum with 13-15 strong teeth, in almost a straight row. Lateral tufted hairs of *thorax* moderately developed, some hairs on dorsum single and long. *Comb* of 30-35 teeth in a patch, each tooth ending in several points. *Siphon* about 2 mm. long, and about 5 times length of diameter at base. Pecten of 22-25 teeth, extending along nearly basal $\frac{1}{2}$ of tube, each with one or two lateral denticles and a long fine point. Hair-tuft of 6-10 fairly strong barbed branches at about middle of tube. *Anal segment* enclosed, except ventrally, within chitinousation, some strong spines on posterior margin towards dorsum. Both *isc* and *osc* usually single and fairly long. Anal fan of numerous (about 28) rather short branched hairs, extending along entire length of segment ventrally (a somewhat unusual feature in larvæ of Indian mosquitoes). *Papillæ* pointed and about length of segment.

HABITAT—Open natural pools, chiefly during rainy season.

DISTRIBUTION—Widely distributed from RAJPUTANA to BURMA, and through CENTRAL PROVINCES and MADRAS to Ceylon.

Occurs throughout tropical Africa, but apparently not further east than Burma.

* Christophers 1906, p. 13, Bedford 1919, p. 739.

61. *Aedes (Mucoidus) laniger* (Wiedemann), 1821

Culex laniger, Dipt. Exot. p. 9 (♀) TYPE-LOC Java TYPE-♀ in Copenhagen Mus

ADULT ♀*—Very similar in appearance to *M. scatophagoides*, but no white median ring on front or middle tibiae, that on hind tibia less distinct and may be absent; all tibiae more broadly white at tip, front and middle tarsi wholly ochreous-brown, hind tarsi with segment 1 fully as long as hind tibia, brown, except for a narrow white ring at base, 2 wholly brown, 3-5 whitish, with tips narrowly brown

♂—Differs from that sex of *M. scatophagoides* as in ♀, palpi less spatulate apically. Hypopygium (fig. 34) very similar to that of *M. scatophagoides*, but both stem and blade of harpago longer, and blade less curved

LARVA—Unknown

DISTRIBUTION. ASSAM †(?) two ♂♂ in M.S.I. collection (*Christophers*) CEYLON Trincomali, iv 1915 (*C. F. Baker*)

Known also from Malayan region as far east as Philippine Islands.

Subgenus *OCHLEROTATUS* Lynch Arribalzaga, 1891.

Rev. Mus. La Plata, n. p. 143 Genotype, *O. confirmatus* L. A. (= *scapularis* Rond.)

ADULT ‡.—Apart from the structure of the ♂ hypopygium, this subgenus is difficult to separate from *Aedimorphus*, the structure of the ♀ hypopygium, the general appearance, and scale-structure being similar. In most species, however, lower mesepimeral bristles are present. Soales on vertex of head and on scutellum always narrow. Proboscis rather slender, and longer than fore femur. Palpi of ♂ about length of proboscis, last two segments more or less thickened, hairy, and turned slightly downwards. Plume-hairs on ♂ antenna directed mainly dorsally and ventrally. Segment VIII of ♀ abdomen completely retractile, cerci long and narrow, and usually, in dried specimens, projecting from the hollow ring of VII, as in *Aedimorphus*. Coxite of ♂ hypopygium with basal lobe, and usually apical lobe also, basal lobe carrying a long spine. Style long and slender, with terminal appendage. Harpago present, consisting of stem and blade,

* Edwards 1922 a, p. 451; Barraud 1929 a, p. 1053, Leicester 1908, p. 69 (as *M. mucidus*).

† Specimens examined by Author. Probably from Assam, but no label attached giving locality.

‡ Edwards 1932, p. 135, Barraud 1928 a, p. 655; Barraud & Covell 1928, p. 676 (buccal cavity).

as in *Mucidus* and *Finlaya*. Phallosome not divided into lateral plates and without teeth, as in *Finlaya*.

LARVA—Those of only two Indian species are at present known. In both the comb consists of 6–20 teeth irregularly arranged, pecten of 18–20 teeth in a close-set rank, none of the distal teeth being widely spaced, as is the case in most species of *Aedymorphus*.

DISTRIBUTION and BIONOMICS—Although over 100 species of the subgenus have been described, only a very few are known from the Indian region at present; these are all confined to the North-West Frontier and Kashmir, and represent intrusions from the Palearctic region.*

The larva of *A. (O.) caspius* (like those of the majority of Palearctic species) lives in ground-pools, but that of *A. (O.) pulchritarsis* has been found in tree-holes. The adult ♀♀ readily attack man and are often troublesome.

62 *Aedes (Ochlerotatus) caspius* (Pallas), 1771.

Reise versch. Prov. Russ. Reichs 1, p. 475 (*Culex*) (♀ ?) TYPE-
Loc. : shores of Caspian sea. TYPE non-existent.

(For synonymy and varieties vide Edwards, 1932.)

ADULT †.—Distinguished by speckled wings and creamy rings over tarsal joints. Wing about 4 mm.

♀—**Head**. dorsum thickly covered with white and fawn, narrow and upright scales, the white scales continued forwards between eyes. Tori with white scales on inner side. Clypeus pale brown. Palpi mottled with light and dark scales. Proboscis either mainly pale or mottled, but tip usually dark, for a varying distance in different specimens. **Thorax**: mesonotum densely clothed with fawn-coloured scales and with white scales forming a pair of more or less defined submedian stripes, running from front, back to scutellum, latter with narrow pale scales and yellow bristles. Broad white scales covering larger part of pleurae. **Wings**:

* In addition to the three species described here, it is certain that others occur in Baluchistan and parts of northern Kashmir. A few ♀♀ have been received from there, collected by Colonel L. G. Browne, I.M.S., at elevations from 7,300' to 14,000' above sea-level. Some of these specimens resemble *A. (O.) cataphylla* Dyar, but in the absence of ♂♂ an identification cannot be made with certainty, and it seems inadvisable to add this species to the Indian list until more material has been obtained. [Two females of another species, probably undescribed, were collected at an altitude of over 15,000' in Kashmir by G. Evelyn Hutchinson in August 1932. These have almost all the abdominal scales cream-coloured, and are peculiar in having no lower mesepimeral bristles, the tarsi are dark, also costa of wing.]

† Edwards 1921 c, p. 299, and 1926, p. 422; Kirkpatrick 1925, p. 97, Barraud 1928 a, p. 668, Martini 1930, p. 271.

speckled with light and dark scales, the white scales usually predominating in Indian specimens *Legs.* very pale, speckled with dark scales, tarsi with apical and basal pale rings, segment 5 of hind tarsi pale *Abdomen* tergites thickly covered with pale fawn, white, and dark brown scales in varying proportions and arrangement Sternites entirely pale.

♂.—Markings as in ♀ Palpi a little longer than proboscis, distinctly thickened on apical $\frac{1}{2}$, marked with white, fawn, and brown scales. last two segments with tufts of yellow hairs Antennal plumes dense, yellow *Hypopygium* (fig 34, 1) coxite with pronounced basal lobe carrying a strong bristly spine with hooked tip, one short spine, and a small number of flattened straight bristles, no very pronounced apical lobe Stem of harpago of moderate length and hairy, blade about same length as stem

PUPA *.—With the characters noted above for the genus Lateral tuft on seg. VIII short, with about 8 branches, paddle wholly pale, with fringe extremely short and fine, terminal hair $\frac{1}{2}$ length of paddle, and may be single, double, or triple

LARVA †.—*Head* moderately chitinised, some darker patches posteriorly Antenna moderately long, slightly curved, shaft with spinelets, those on apical $\frac{1}{2}$ strong, this part of antenna also darker than basal $\frac{1}{2}$, tuft of about 6 branches, arising at about middle of shaft Frontal hair *A* with 8–10 branches, *B* and *C* usually single and subplumose, *C* standing almost directly behind *B*, *d* small, with several fine branches Mouth-brushes normal. Preclypeal spines long and slender Mentum triangular, about 10 moderately large teeth either side of central one Lateral hairs of *thorax* and *abdomen* well developed *Comb* of 15–20 teeth, each ending in a number of sharp points *Siphon* about 1 mm. long, pale in colour, with narrow dark ring at base *Aous* fairly large. Pecten of about 20 fairly long, pointed teeth, each with 3–6 lateral denticles near base Hair-tuft of 4–6 branches at rather beyond middle point from base *Anal segment* with chitinised saddle covering dorsum and sides and with large fan-plate ventrally Surface of saddle (and of siphon) with numerous minute sharp points, *lh* single, *isc* with a number of subequal branches Fan hairs numerous and moderately long. Anal papillae $\frac{1}{2}$ – $\frac{1}{3}$ length of segment, tips pointed (length subject to variation)

HABITAT.—Open natural pools, both fresh water and brackish

* Theodor 1924, pp 341–5; Kirkpatrick 1925, p 83

† Lang 1920, p 60; Wesenberg-Lund 1921, p 39, Kirkpatrick 1925, p 82, Martini 1930, p 272, Montschadsky 1930, p 596

DISTRIBUTION—KASHMIR Baltistan, Bragica, 7,300', viii 1923 (*Browse*) N.W. FRONTIER Peshawar, iii. 1913 (*Howlett*), x 1914 (*Fletcher*), Taru, Peshawar dist., v 1916 (*Fletcher*), Nowshera, Peshawar dist., vi. 1911 (*Smith*), Rawalpindi, vi-vii 1917 (*Hodgart*) Following from Cent Mal Bureau records.—Bannu, vii 1925, Chaman, vi 1928; Dera Ismail Khan, 1929

Known also from EUROPEAN COASTS, CENTRAL EUROPE, and eastwards to GOBI DESERT, NORTH AFRICA, PERSIAN GULF, and MESOPOTAMIA.

63 *Aedes (Ochlerotatus) pulchritarsis* (Rondani), 1872.

Bull Soc Ent Ital iv, p 31 (*Culex*) TYPE-LOC Italy TYPE Florence Museum

[ADULT*.—Distinguished by the dark wings and pure white rings extending over the hind tarsal joints, the 5th segment being entirely white † Wing 3-4.5 mm.

♀.—*Head* a median line of long and rather wide white lanceolate scales on vertex, continued forwards between eyes and around eye-margins, white upright scales in centre of nape, a large patch of black scales on either side of middle line of vertex Palpi about $\frac{1}{2}$ length of proboscis, black, with conspicuously white-scaled tips Proboscis black, with a few grey scales over middle $\frac{1}{2}$ on underside *Thorax*: mesonotum with a variable amount of golden scales towards front in middle, pale scales along margins and over wing-roots, remainder dark brown. Scutellar scales narrow and mostly white, *apn* with white lanceolate scales, *ppn* with some dark scales anteriorly, flat white scales posteriorly, 4-5 lower mesepimeral bristles *Wings*: dark scaled, plume-scales long and narrow *Legs* mostly black, with conspicuous white knee-spots and white rings on tarsi, narrow on front and middle pairs, broader on hind pair, 5th tarsal segment of all legs white. *Abdomen*. black, with white basal bands on tergites II-VI, sternites similarly coloured.

♂.—No specimens available. *Hypopygium* with *bl* small, bearing one distinct spine

LARVA †.—See under var. *asiaticus*.

HABITAT.—Tree-holes

DISTRIBUTION Chiefly SOUTHERN EUROPE, but extending into TURKESTAN and KASHMIR.

* Edwards 1921, p 1031; 1926, p. 429, and 1932, p. 141 (syn. and vars), Barraud 1928 a, p. 668, Martini 1930, p. 316.

† [In ornamentation this resembles some species of subgenus *Finlaya*, especially *A. (F) greeni*, but may be distinguished by the long cerci and small sternite VIII in ♀]

‡ Montschadsky 1926, p 151, and 1930, p. 609; Vogel 1929, p. 161; Martini 1930, pp 317-319.

The species is usually scarce and still but little known, but appears to be subject to considerable local variation. Two different varieties, or possibly distinct species, occur within the Indian area and are described below]

[var *asiaticus* Edwards, 1926.

Riv Malar v, p. 430 (14 x 1926). TYPE-LOC Drosh, Chitral,
x 1915 (*W O Walker*) TYPE ♀ in Brit Mus

stegomyia Stackelberg & Montschadsky, Bull Ent Res xvii,
p. 151 (15 x 1926) TYPE-LOC Katta-Kourgan, Turkestan
TYPE Leningrad (?)

ADULT ♀.—Mesonotum mainly dark brown, with a double median line of white scales on anterior $\frac{1}{2}$, narrowly bordered externally with yellow scales, on each side of the posterior end of this stripe a rather large patch of whitish scales. White basal bands on tergites II-VI complete, that on VI very narrow. Femora and tibiae with fairly numerous scattered white scales, hind femur (except for this speckling) dark on nearly the whole outer surface]

LARVA *.—Antennal shaft smooth, hair-tuft of 3-4 hairs at about middle. The four pairs of frontal hairs, *A* to *d*, all with moderately long branches, and all placed rather far forward on clypeus. Hairs on *thorax* and *abdomen* well developed, those of dorsum of abdomen of stellate form. *Comb* of 6-10 large pointed teeth, fringed on basal $\frac{1}{2}$, arranged in an irregular row. *Siphon* dark, about 1 mm. long by 0.3 mm. broad at base, slightly tapering from middle to apex. *Pecten* of about 18 teeth in close-set row, each with 3-6 basal lateral denticles. *Siphonal* tuft 3-4-branched, at about middle of tube. *Anal segment* with small chitinous saddle, *lh* single, fairly long, *sc* of about 5 subequal branches of moderate length. Fan of a small number of 2-branched hairs arising from plate. *Papillae* very long, $\frac{1}{4}$ length of whole body and about twice length of siphon, sausage-shaped, with rounded ends.

DISTRIBUTION.—Apart from type-localities, recorded only from Old Bokhara (*Montschadsky*).

var. *versicolor* Barraud, 1924 †.

Ind. Journ Med. Res xii, p. 73 (*Finlaya versicolor*) (♀) TYPE LOC..
Yusimarg, Kashmir, 7500', viii. 1923 (*T. B. Fletcher*). TYPE
♀ in Brit Mus

ADULT ♀.—Mesonotum with a broad median yellowish patch on anterior $\frac{1}{2}$, not including a white central line,

* *Montschadsky* 1926, p. 151, and 1930, p. 609.

† [This differs very little from var. *berlandi* Ségué, 1921, found in the neighbourhood of Paris. The French form (according to a somewhat damaged ♀ in the British Museum) appears to have the white rings of the hind tarsi broader.]

indefinite patches of yellowish scales on each side of the median patch behind. White basal bands on tergites II-VI complete and very distinct. Femora and tibiae without any scattered pale scales, hind femur on outer side with basal $\frac{1}{2}$ whitish, apical $\frac{1}{2}$ black; a dark dorsal line runs almost to base. White rings on hind tarsi narrow, even that on 1-2.

LARVA.—Unknown

DISTRIBUTION.—Known definitely only from type-locality.

[64 *Aedes (Ochlerotatus) pullatus* (Coquillett), 1904

Proc Ent Soc Wash vi, p 168 (*Culex*) (♂♀) TYPE-LOC Kaslo, British Columbia (*H & Dyar*) TYPE U S National Mus
Aedes pullatus Howard, Dyar, and Knab, 1917, Mosq. N. & C Amer iv, p 738

ADULT*.—Characterized by dark tarsi and hypopygial structure. Wing about 5 mm

♂—*Head* with pale yellowish narrow scales. Palpi thin, slightly shorter than proboscis, last two segments and the slightly swollen tip of long segment with a moderate number of long hairs. *Thorax* clothed with pale yellowish scales, few or no darker ones on mesonotum. Lower mesepimeral bristles few in number (apparently only one on one side, none on the other, in the Kashmir specimen). *Wings*: dark scaled. *Legs* mainly dark, femora pale beneath and with some pale speckling. *Abdomen* blackish above, tergites with whitish basal bands widening out at sides. *Hypopygium*. coxite with well-marked apical lobe, rather numerous long hairs on outer side and near tip on inner side, but no definite hair-tufts, basal lobe very small, with one strong spine directed at right angles to axis of coxite, and at some distance from the spine a pair of curved bristles, on a small eminence. Harpago with rather long curved stem and broad blade; basal part of stem pubescent, with a more or less prominent angle or shoulder on inner side at some distance from base.

♀ (American specimens).—Colouring as in ♂, rather more pale scales on tibiae and at bases of tarsi. Often dark stripes present on mesonotum.

LARVA (American specimens).—Antennae covered with spicules, tuft just before middle. Frontal hair *B* with about 4 branches; *C* with about 8, placed behind *B*; *d* small. Comb with numerous teeth. Siphon about 3×1 , tuft near middle, pecten of numerous and uniformly spaced teeth. Anal papillae very long and pointed.

* Edwards 1921 c, p. 316, and 1926 b, p. 449.

HABITAT (in America).—Pools formed by melting snow, in mountains in spring.

DISTRIBUTION.—KASHMIR: Gulmarg, vi 1931 (*T. B. Fletcher*), 1 ♂ *

Also widely spread in mountains of CENTRAL and SOUTH EUROPE and WESTERN NORTH AMERICA.]

Subgenus FINLAYA Theobald, 1903.

M.C. iii, p 281. Genotype, *F. poicilia* Theo

Finlaya Giles (emend.), 1904, Journ Trop Med. vii, p 366

Danileva Theobald, 1904, Entom xxxvii, p 78 Genotype,

D. albocinctata Theo

Hulecoetomyia Theobald, 1904, Entom xxxvii, p. 162. Genotype,

H. trilineata Theo.

Popea Ludlow, 1906, Can Ent xxxvii, p 95 Genotype, *P. lutea* Ludl

Phagomyia Theobald, 1905, Gen Insect., Fam Culicidae, p 21

Genotype, *P. gubernatoris* (Giles)

Lepidatomyia id., ib p 22 Genotype, *L. magna* Theo

Pseudocarrollia Theobald, 1910, Rec Ind Mus iv, p 13 Genotype, *P. lophocentralis* Theo

ADULT †—Rather small or medium-sized mosquitoes of very varied ornamentation and scaling. Some species resemble those of the subgenus *Stegomyia*, being black with white markings, these may be distinguished on the points given under that subgenus. In others the mesonotum is ornamented with fine lines of white or yellow scaling, and in some there is a pale band, or pale scaling to a greater or less extent, on the proboscis. Some other species are much less ornamented. Proboscis usually longer than fore femur. Palpi of ♂ from rather more than $\frac{1}{2}$ to fully as long as proboscis. Palpi of ♀ from $\frac{1}{2}$ to $\frac{1}{3}$ length of proboscis. Antenna of ♂ much as in *Stegomyia*. Tarsal claws of fore and mid-legs toothed in both sexes. No lower mesepimeral bristles. In the ♀ segment VIII is only slightly retractile and laterally compressed, the sternite large and usually without scales, and the cerci quite short. Coxite of ♂ hypopygium long and usually without lobes. Style fairly long and simple, with usually a long terminal appendage. Harpago well developed, with stem and flattened blade (except in *dissimilis* and *subsimilis*, in which the stem is short and the harpago represented

* [This Kashmir specimen has rather broad and flat scales on *apn*, *ppn*, and mid-lobe of scutellum, these being normally quite narrow, but the hypopygial structure is almost or quite identical with North American *pullatus*. Until more material is forthcoming it is impossible to determine whether the Kashmir form is a distinct local variety.]

† Edwards 1932, p. 147. See also Barraud 1923 g, p 476, and 1924 a, p. 845.

by a stout bristle). In a few species (*albolateralis*, *niveus*, *pulchriventer*, etc.) there is a row or tuft of very large scales on ventral border of coxite towards apex. Phallosome tube-like or scoop-shaped, not divided into lateral plates, sometimes with serrations on apical rim, but no definite chitinated teeth as in *Stegomyia*. Paraprocts with 2-3 teeth at crown and a few minute hairs just below (characters absent in *Stegomyia*).

LARVA (4th stage) —Antenna usually fairly short and with or without spicules on shaft. Antennal hair single or with a few branches. Frontal hairs very variable in arrangement and character. Occasionally the four important pairs are all well developed and many-branched (*albolateralis*, *albo-cinctus*, *suffusus*). In another group, hairs *A*, *B*, and *C* have several rather short branches, *B* and *C* being placed well forward towards the front margin of clypeus (*pulchriventer*, *elsae*, *pseudotaenatus*, *shorti*, etc.). In *christophersi*, *gilli*, and *greeni* hairs *B* and *C* are placed further back and close together, and one or both may be very long, as long as the whole head. In a number of other species (*gubernatoris* group) hairs *B* and *C* are placed one behind the other, both usually single, and of moderate length. Chaetotaxy of thorax and abdomen of variable development, in a few species large spines are present on dorsum of thorax. Comb nearly always a triangular patch of numerous scale-like teeth, but in *albolateralis*, *niveus*, and *suffusus* there are fewer large teeth arranged more or less in a single row, as in *Stegomyia*. Siphon of only moderate length, with a single pair of hair-tufts near middle. Acus nearly always present. Pecten well developed, usually of close-set teeth with lateral denticles, in a few species there are some larger simple teeth towards apex of tube. A chitinated saddle on anal segment; anal fan only moderately developed.

DISTRIBUTION and BIOLOGICS —The subgenus has a world-wide distribution, but attains its greatest development in the Oriental region, where about 50 species are known out of a total of about 92. In the Indian region about 38 species occur, and it is most probable that this number will be increased by further collecting. About half the known species are found in the Himalayas. The larvae live either in tree-holes or bamboo-stumps during the monsoon periods, or in rock-pools and potholes in stream-beds. Some of the commoner species, such as *albolateralis*, are very persistent blood-suckers in the day-time, and cause much annoyance in certain forest localities. No species is known to transmit disease.

Key to Adults

- 1 Wings elaborately spotted and speckled with black and white scales
Wings not spotted or speckled, wing-scales all dark, except for a short line of pale ones at base of costa in some species 2.
- 2 Tarsi entirely dark 3.
- 2 Tarsi with white markings 7
- 3 Mid-femur with a median silvery mark on anterior surface
Mid-femur without such mark *poecilius*, p 157
4
- 4 Mesonotum with a large snowy-white patch in front, which may, in the ♀, be more or less divided into lateral patches
Mesonotum with ochreous, yellow, or golden scales [pp 205-212.
noveus group, 5.
- 5 Abdominal sternites with orange patches.
Abdominal sternites without orange patches *pulchriventer*, p 199. 6
- 6 Mesonotum, in ♀, black, with ochreous scales arranged in lines, in ♂ entirely, but sparsely, covered with pale scales
Mesonotum covered with ochreous scales, with a pair of indistinct submedian dark lines *oreophilus*, p 192
suffusus, ♀, p 194
7. Hind tarsi with one or more white rings at bases of segments only 8
Hind tarsal segments with both apical and basal white rings 23
- 8 Venter of abdomen with orange markings
Venter of abdomen without orange markings *auronitens*, p 202. 9
9. Hind tarsi with only one white ring. *unacinctus*, p. 170 10
10. Hind tarsi with three white rings 11.
10. Hind tarsi with four white rings 20.
- 11 Mesonotum marked with narrow lines of golden scales 12.
- 11 Mesonotum marked otherwise 16
12. Proboscis with a pale ring or with pale scaling on underside 13.
12. Proboscis entirely dark *saxicola*, p. 191. 14
13. Proboscis with pale scaling both on upper and undersides 15.
13. Proboscis pale on underside only 16.
- 14 Proboscis pale on basal $\frac{1}{2}$ both on upper and undersides *pallirostris*, ♀, p. 190.
- 14 Proboscis extensively pale on underside and with white scaling forming a narrow band on upper side *chrysolineatus*, p. 185.
harveyi, p. 188.
15. Mid-femur dark on anterior aspect *formosensis*, p. 189.
15. Mid-femur with a pale line on basal $\frac{1}{2}$ anteriorly *stevensoni*, p 174.
16. Mesonotum blackish, with a white spot in front 17.
16. Mesonotum otherwise

17. All femora with small white knee-spots, scales on dorsum of head mainly narrow *christophersi*, p 195.
 All femora dark at knee, scales on dorsum of head mainly broad and flat 18.
18. Posterior, or under, surface of fore tibia conspicuously pale for whole length . . . *gilli*, p. 196
 Fore tibia dark, except narrowly at base 19
19. Mid-femur dark on anterior surface, except at extreme base *sumlensis*, p 198
 Mid-femur with a well-defined white streak on anterior surface, ventrally, on basal $\frac{1}{4}$ *albocinctus*, p. 172
20. Hind tarsi with white ring on segment 4 very wide and covering nearly the whole segment *subsumilis*, p 203
 Hind tarsi with white ring on segment 4 not very wide 21
21. Mesonotum with a large area of white scaling in front *albotæniatus*, p 174
 Mesonotum dark, or with pale scaling forming lines 22
22. Mesonotum dark on anterior $\frac{1}{2}$, or with an indistinct median yellow line [ranus, ♀, p 176.
 Mesonotum with a median and sublateral lines of white or creamy scales. *albotæniatus* var *miki-*
shoriti *, p. 183.
23. Mesonotum with a white patch in front 24
 Mesonotum without a white patch in front 33
24. Venter of abdomen with very long outstanding tufts of scales 25
 Venter of abdomen with only moderately developed tufts of outstanding scales or with none 26
25. *ppn* bare *khazani*, p. 168
ppn with white scales *prominens*, p 169
26. *ppn* with only a small patch of white scales on posterior border 27
ppn with a large patch of white scales 29.
27. Venter of abdomen with moderately developed tufts of outstanding scales 28
 Venter of abdomen without tufts of outstanding scales *cogilli*, p 165
28. Fore tibia dark on posterior surface, except for an apical white ring, scutellum densely clothed with flat white scales, fore femur dark on upper, or dorsal, surface at apex *lophoventralis*, p 167.
 Fore tibia pale posteriorly for whole length, scutellum much less densely clothed with white scales, fore femur with a white spot at apex on upper, or dorsal, surface
29. Scutellar scales brownish-black. 30
 Scutellar scales mainly white 32
30. Head without a median white-scaled line in ♀ *feegraderi*, p 164.
 Head with a median white-scaled line 31

* Hind tarsi with some white scales at apices of first 3 segments not forming definite basal and apical rings

- 31 Abdomen with small outstanding tufts of scales on venter and roughened scales on dorsum *assamensis*, ♀, p. 166.
 Abdomen without obvious tufts of outstanding scales *deccanus*, p. 163
32. Venter of abdomen with small outstanding tufts of scales, mesonotum almost entirely covered with white scales *assamensis*, ♂, p. 166
 Venter of abdomen without tufts of outstanding scales, white scales of mesonotum confined to an anterior patch and one in front of each wing-root *gubernatoris*, p. 159.
- 33 Mesonotum marked with lines of white, yellowish, or golden scales on a dark ground 34.
 Mesonotum covered with ochreous, brown, or golden scales, not arranged in lines 38.
- 34 Proboscis entirely dark 35.
 Proboscis with pale scaling on under-surface 37.
- 35 Mesonotum with a line of creamy scales each side, continued over wing-root, no median line, but three small patches of pale scales on anterior margin *sinioni*, p. 200
 Mesonotum with median and lateral well-defined lines of white or golden scales, no patch of scales on anterior border of mesonotum 36
36. Pale lines on mesonotum white or creamy; femora with white longitudinal lines for whole length *pseudotaenratus*, p. 178
 Pale lines on mesonotum golden, femora not lined with white *greeni* var. *kanaranus*, [p. 185.
- 37 Proboscis pale on underside for basal $\frac{2}{3}$, pale lines on mesonotum very narrow, clearly defined and golden *macdougalli*, p. 177
 Proboscis pale beneath from near base to tip, except for a small interruption at about $\frac{1}{3}$ from base; lines on mesonotum pale yellow, the median one formed of two lines of scales placed close together. [shorti, p. 183.
clavæ, p. 180, cf also
38. Last segment of hind tarsi white dorsally, mesonotal scales bright golden *greeni*, p. 184
 Last segment of hind tarsi dark, mesonotal scales brown in ♀, pale yellow in ♂. *inguinatus*, p. 169.

65. *Aëdes* (Finlaya) *pocellus* Theobald, 1903.

M.C. III, p. 283 (*Finlaya pocellia*) (♀) TYPE-LOC : Penang, Malay Penin. x. 1901 (*Freer*). TYPE : ♀ in Brit Mus.

ADULT *.—Very distinct on account of the spotted wings and presence of numerous spots and bands on the legs. In general appearance it somewhat resembles *Aëdomyia venustipes* (Skuse). Wing about 2.5–3 mm.

* Edwards 1924, p. 380, and 1926 a, p. 104.

♀—*Head* a few white scales in middle of vertex and two patches of white scales at each side, remainder covered with black scales, rather numerous upright scales on nape extending forwards some distance. *Tori* with white scales on inner sides, some black scales on first flagellar segment, antenna otherwise brown. *Clypeus* and *palpi* black, latter with white scales at apices. *Proboscis* black, with a clear white ring rather beyond the middle and a small white ring at base of labella. *Thorax*: *mesonotum* black, with a covering of greyish and white scales, some flat silvery scales laterally in front of wing-roots. *Scutellar* scales black on lateral lobes, black and white on mid-lobe. *Integument* of *pleuræ* black, with patches of white scales, some of these forming a more or less continuous line from *meseponeron* to *apn*, some black and white scales on *ppn*. *Wings*: speckled with rather broad black and white scales, white scales usually forming two or three small spots on costa. *Legs*: black, with numerous white rings, all femora and fore and mid-tibiæ each with about six white rings and narrowly pale at tips, hind tibiæ with about four similar rings. *Tarsi* with three or four white rings, segment 5 also entirely pale. *Abdomen*. brownish-black, a pair of submedian white spots in middle of each tergite, laterally a white longitudinal mark on tergal borders and a white spot near hind margins, on sternites V–VII there are outstanding tufts of dark scales.

[♂ (specimen from Philippine Is in British Museum) — Colouring much as in ♀. *Palpi* about as long as *proboscis*, dark brown, with five white rings and narrow white tip; few long hairs on last two segments. *Hypopygium*: *coxite* with cluster of about 8 very large but pointed scales on inner side at about mid-length, *style* rather short, widened beyond middle, *blade of harpago* very long, scarcely widened in middle, ending in a long point.]

[LARVA * — Pale in colour, including head and siphon. *Antenna* short, shaft smooth, with a single or bifid hair beyond middle. *Frontal hairs* *A* 5-branched, *B* 2-branched, *C* 2–3-branched, placed well behind *B*, *d* 3-branched, nearly as long as *B*. *Thorax* and *abdomen* with numerous many-branched stellate hairs, but no special spines on plates on thorax. Comb of numerous teeth in a patch. *Siphon* short and rather broad, its surface covered with minute hairs, no *acus* (an unusual feature in this subgenus); *pecten*

* [Brug 1931 b, p. 23. Redescribed above from some of Brug's Javan specimens in the British Museum. The larva of the allied *A. (F.) kochi* of New Guinea and Samoa, of which *A. poscilus* was at one time regarded as a variety, differs in several respects, having clypeal hair *C* long and single, siphon without fine hairs on surface, pecten-teeth simple and needle-like.]

of only 6-10 rather large teeth, which are finely fringed all along one side and at the tip, no simple teeth at end of row and none detached, tuft of 3-5 branches just beyond middle of tube. *Anal segment* with moderately large saddle with spiny margin, *isc* with 4-5 branches, *lh* long, 2-branched, papillæ longer than segment, equal and pointed.]

HABITAT (in Java) —Leaf-axils of *Colocasia indica* and *Crunum* sp

DISTRIBUTION —N. BENGAL: Sureil and Tindharia, Darjeeling dist., 5,000' (*M. O T Iyengar*) BURMA Sandoway*, x 1924 (*Civil Surgeon*)

Recorded also from MALAY PENINSULA and AROHIPHLAGO, and PHILIPPINES

66 *Aedes* (Finlaya) gubernatoris (Giles). 1901

Journ Bomb Nat. Hist Soc xiii, p 607 (*Culex*) (♀) TYPE-LOC Allahabad, Govt House, vii (*Giles*) TYPE ♀ in Brit Mus

Lepidotomyia magna Theobald, 1905, Gen Ins., Fam Culicidæ, p 22 (♂ & ♀) TYPE-LOC : Bombay, viii 1902 (*James*) TYPE ♂ & ♀ in Brit Mus

var. n. *kotensis* (see under "Larva") TYPE-LOC Kotli, near Kalka, W Himalayas


ADULT † (Pl I, figs. 5, 6) —A fairly large black and white species. One of the commoner tree-hole breeding mosquitoes of Western India. Wing about 4.5-5 mm

♀.—*Head*: a median pale line on vertex, a narrow pale border to eye-margins widening out at sides, a dark patch on either side of median line. In some specimens the dark patches are faint or absent and the head then appears mainly pale. Fairly numerous yellowish upright scales on nape extending forwards some distance. Antennæ, clypeus, palpi, and proboscis black. Palpi about $\frac{1}{2}$ length of proboscis, latter about same length as fore femur. *Thorax*: a large patch of white scales on front of mesonotum, the posterior margin of which is usually deeply emarginate, as shown in Pl. I, figs. 5, 6; a large oval white patch in front of each wing-root, and usually a thin median line posteriorly, dividing other side of ante-scutellar bare space, remainder of mesonotum covered with brownish-black scales. Mid-lobe of scutellum covered with lanceolate creamy or white scales, usually very few narrow hair-like white scales on lateral lobe. Postnotum and integument of pleuræ brownish-black, *apn* and larger part of *ppn* with broad white scales, several large and small patches of flat broad white scales

† Theobald 1901 a, p. 314 (*Stegomyia*), 1907, p. 223 (*Phagomyia*), and p. 250 (*L. magna*); Edwards 1913 b, p. 226, and 1917, p. 214 (syn.); Barraud 1923 d, pp. 214 and 216 (*F. magna*), 1923 g, pp. 482-3.

on pleuræ. *Wings* dark scaled. *Legs*: black, with white markings, fore femur black anteriorly, white on nearly basal $\frac{1}{2}$ beneath, mid-femur black, with white ventral streak on anterior surface on basal $\frac{1}{2}$, and white marking at knee, posteriorly broadly pale, except dorsally, on nearly apical $\frac{1}{2}$, hind femur white, except for narrow basal dark ring and wider subapical ring. Fore tibia usually very narrowly pale at base, broadly pale at tip and entirely pale for whole length beneath, mid-tibia black, with pale marking

Fig. 35



♂ hypopygial structures of *Aedes*, subgenus *Finlaya*. a, *gubernatoris*; b, *cacharamus* (style), c, *khazani* (style and harpago), d, *unicinctus* (phallosome, dorsal surface to left). Lettering as on p 4

at base and larger pale marking at tip, hind tibia usually entirely black. Fore tarsi usually very narrowly pale at base of first two segments, sometimes some pale scales at apex of segment 1, this segment also pale beneath for whole length, other parts black, markings of mid-tarsi subject to some variation, basal and apical white rings on segment 1 and medial dark ring variable in size, sometimes absent, when whole segment is white, a basal white ring on segment 2, usually occupying about $\frac{1}{2}$ the segment or more; there may,

or may not, be pale scaling over joint between 2 and 3; hind tarsi with rather narrow white basal and apical rings to segment 1, a basal white ring on 2, and usually some pale scales at joint between 2 and 3. *Abdomen*: black, with usually median basal yellowish or white markings, not forming complete bands, always lateral basal white patches on tergites I-VII and white basal bands on sternites, sternite VIII large, without scales, cerci short.

♂—Palpi black, about length of proboscis, last two segments with moderately long outstanding hairs. White scaling on mesonotum usually more extensive than in ♀, and head-scaling generally lighter. *Hypopygium* (fig 35, a) ♂ with 9-10 rather long hairs on each lobe, coxite long and narrow, nearly 4 times length of greatest width, style with terminal appendage, latter more than $\frac{1}{2}$ length of former; phallosome scoop-shaped. Harpago expanded in middle, blade much longer than stem, paraproct toothed at crown, with a few minute hairs below.

LARVA *, type-form (fig 36, a, e).—Arrangement of frontal hairs shown in figure (a). Antennal shaft smooth, about 12 times as long as wide, antennal hair single, arising at nearly $\frac{3}{4}$ from base. Lateral hairs of thorax well developed and strong, no large spines or special structures. Abdominal segments with fairly long 2-branched hairs. *Comb* of 30-40 scale-like teeth, each tooth rather broad at tip and ending in a number of points. *Siphon* $1\frac{1}{2}$ to twice length of diameter at base. *Pecten* of 14-17 strong teeth, each usually with 2 denticles near base. Hair-tuft of about 6 plumose blade-like branches, somewhat resembling a palmate hair (characteristic of this and several allied species), base of hair-tuft nearer apex than base of tube, sometimes between the pectens, but usually just beyond. *Anal segment* enclosed, except ventrally, within a large chitinated saddle; *lh* short and strong, unbranched; *sc* of 8-10 rather short, subequal black branches, *osc* single and long, as usual. Fan of 10-12 strong black hairs of only moderate length, with primary and secondary branches. Both pairs of papillæ shorter than saddle, bluntly pointed.

Another form of the larva, var. n. *kottensis*, is known. The adults resulting from these larvæ appear to be indistinguishable from the type-form. The points distinguishing these larvæ and those of *A. (F.) deccanus* and *A. (F.) assamensis*, which are very similar, are given in the summary on p. 163. The larvæ of several other closely allied species are at present unknown.


* Barraud 1923 e, p. 220 (*F. magna*); Senior-White 1927, p. 68.
DIPT.—VOL. V.

HABITAT.—Tree-holes, also rock-pools (?) *

DISTRIBUTION.—Common in UNITED PROVINCES, PUNJAB, and BOMBAY PRESIDENCY.

Records from other parts require confirmation*. The species appears to be absent from Bengal, Assam, and Burma. Further work is required upon this difficult group of species.

Fig 36



Larval structures of *Aedes*, subgenus *Finlaya* a, *gubernatoris* (head, siphon, anal segment, enlarged pecten-tooth and comb-scale; e, hair of mouth-brush), d, *gubernatoris* var. *kotienensis*, two hairs of mouth-brush, b, b', *decanus*, c, c', *assamensis* (head; portion of anal segment with *asc*, hair d omitted in c).

* [A series of specimens in the British Museum from Ceylon (Sudunganga, bred from rock-pools, *Senior-White*) closely resemble *A. gubernatoris* except that scales of mid-lobe of scutellum are purer white and broader. Larvæ of this form have not been preserved.]

Summary of characters distinguishing the larvae of several species of the *gubernatoris* group.—

- Preclypeal spines conspicuous, of moderate length, strong and rather blunt. Antennal shaft 0.28 mm long. Median hairs of mouth-brushes with moderately large teeth. Frontal hair *d* posterior to *B*, and nearer to that hair than to *C* (fig 36, b), usually slightly longer than in *gubernatoris*, and usually of only 3-4 branches (fig 36, b¹). Branches of siphonal tuft as in *gubernatoris* *deccanus*
- Preclypeal spines stout and blunt (fig 36, c). Antennal shaft 0.2-0.25 mm long. Median hairs of mouth-brushes with comparatively large teeth. Frontal hair *d* very small and nearer to *C* than to *B*, *isc* of only 2 long branches (about 1 mm). Branches of siphonal tuft less widened than in *gubernatoris* *assamensis*
- Preclypeal spines, antennal shaft, and branches of siphonal tuft as in *assamensis*. Frontal hairs *B*, *d*, and *C* standing one behind the other at fairly even, wide intervals, *isc* of 3-6 long fine branches *feegrades*
- Preclypeal spines inconspicuous, slender, and tapering to a fine point. Antennal shaft from 0.41 to 0.51 mm long. Median hairs of mouth-brushes with very small teeth. Frontal hair *d* slightly posterior to and near to *B*, *isc* with 6-8 branches, longest about 0.5 mm long. Branches of siphonal tuft widened, somewhat resembling leaflets of a palmate hair (fig. 36, a, e) *gubernatoris* (type- [form]).
- As above, but antennal shaft shorter, from 0.32-0.35 mm long. Median mouth-brush hairs with comparatively very large teeth (fig 36, d) *gubernatoris* var *kotiensis*.

67. *Aedes* (Finlaya) *deccanus* Barraud, 1923

Ind Journ. Med Res. xi, p. 317 (*Finlaya*) (♂ & ♀). TYPE-LOC.: Belgaum, Bombay, Decan, viii 1921 (*Barraud*). TYPE. ♂ & ♀ in Brit. Mus.

ADULT * (Pl I, fig 4).—Very similar to *assamensis* and *feegrades*, differing from the former only in the absence of obvious tufts of outstanding scales on the abdomen, and from the latter in the presence of a median white line on the vertex in ♀.

LARVA † (fig. 36, b).—Very similar to *gubernatoris*, but differs constantly, as indicated in the notes under the description of that larva.

HABITAT.—Tree-holes.

* Barraud 1923 g, p. 483.

† Barraud 1923 e, p. 221.

DISTRIBUTION.—BOMBAY: Deccan, Belgaum * (as above), Tavargatti *, and Nagargah *, viii 1921 (*Barraud*), Bombay Harbour, Trombay, and Elephanta Island *, vii 1921 (*Barraud*) WESTERN HIMALAYAS: Koti *, near Kalka, viii. 1923 (*Barraud*).

Not recorded from outside the Indian area.

68. *Aedes* (Finlaya) *feegradei*, sp. n

TYPE-LOC Rangoon, Burma, 1930 (*Feegrade*) TYPE ♂ & ♀
in Brit. Mus.

ADULT.—Very similar to *cogilli*, *deccanus*, and *assamensis*, differing from the first in having black scales on all scutellar lobes and *ppn* almost entirely covered with flat white scales, from *deccanus* and *assamensis* in absence of a median white line on head in ♀, rather smaller prealar white patches, and rounder white area on front of mesonotum. Wing 3.5–4 mm.

♀.—*Head*—a narrow pale border to eye-margins widening out at sides, vertex without a median pale line, entirely covered with flat black scales, dark upright scales on nape. Antennae, clypeus, palpi, and proboscis black, palpi about $\frac{1}{2}$ length of proboscis, latter slightly longer than fore femur. *Thorax*. mesonotum with fairly large patch of silky white scales in front, posterior margin of this area convex, not emarginate, its sides continuous with a large patch of flat white scales covering *ppn*, a small area of narrow white scales in front of each wing-root and, adjoining this, a small patch of broad white scales on margin of mesonotum behind anterior spiracle, remainder of mesonotum covered with brownish-black scales, no pale scales posteriorly in front of scutellum. Scutellar scales black on all lobes and fairly broad. Postnotum and integument of pleurae black. Broad white scales on *apn*, 2 patches on sternopleura, 1 on upper part of mesepimeron, small round spots of white scales on propleura and on coxae of fore and mid-legs. *Wings* dark scaled. *Legs* fore femur black, except narrowly beneath on basal $\frac{1}{2}$, mid-femur white ventrally on both surfaces on basal $\frac{1}{2}$ and at knee, otherwise black; hind femur with narrow black basal ring, a wider black ring on apical $\frac{1}{2}$, otherwise white, but actual tip of knee dark. Fore tibia black, with a few pale scales at base beneath and a subapical white marking not forming a complete ring above, mid-tibia black, with subapical dorsal white mark, hind tibia entirely black, except for yellowish appearance at tip on inner side. Fore tarsi black; mid-tarsi with rather narrow basal and apical white rings, latter continuous with basal white ring on segment 2, otherwise black, hind tarsi with small dorsal white mark at base of segment 1, a fairly broad white ring over joint between 1 and 2, otherwise black. *Abdomen*.

black, with large basal lateral silvery patches on tergites I-VII, a small silvery dorsal mark on VIII, sternites with basal white bands, small tufts of outstanding black scales on IV-VII, VIII large, without scales, cerci small. On tergites V-VII the black scales in mid-dorsal line project slightly.

♂.—Palpi black, slightly shorter than proboscis, with few outstanding hairs on terminal segment. In type ♂ and one other there is a median white line on vertex flanked by dark patches, a pale border to eye-margins, widening out considerably at sides. White area on mesonotum larger than in ♀ and continued posteriorly in three faint lines to scutellum. Scales on scutellum black as in ♀, and other ornamentation similar.

LARVA.—Very similar to *assamensis*, but frontal hairs *B*, *d*, and *C* standing one behind the other at fairly even, wide intervals, and *isc* of 3-6 long, fine branches. Siphonal tuft divided into 7-8 strong barbed branches, less widened than in *gubernatoris* and *deccanus*.

HABITAT.—Tree-holes.

DISTRIBUTION.—Known only from type-locality.

69. *Aêdes* (Finlaya) *cogilli* Edwards, 1922.

Bull. Ent. Res. xii, p. 92 (♀). TYPE-LOC. Karwar, N. Kanara, ix.-x. 1902 (*Cogill*) (♀), Karwar, ix. 1921 (*Barrand*) (♂). TYPE: 2 co-type ♀♀ and allotype ♂ in Brit. Mus.

ADULT ♀† (Pl. II, fig. 3).—A medium-sized black and white species resembling *gubernatoris*, but differing as follows.—Head without a median white line, eye-margins with narrow border of white scales, *ppn* bare, except for a small patch of flat white scales on lower posterior margin. White patch on front of mesonotum somewhat large, rounder, and more silky; prealar white patches smaller. Scutellum with mid-lobe densely covered with flat white scales, flat black scales on lateral lobes. Leg markings very similar to those of *deccanus*, but scaling of head and scutellum different.

♂.—Dorsum of head covered with flat dark scales, a narrow border of white scales to eye-margins widening out at sides; palpi distinctly shorter than proboscis, terminal segments with few hairs. *Hypopygium*: coxite with small round lobe at base of dorsal border, bearing a collection of long hairs. Blade of harpago tapering, and hardly expanded in middle.

LARVA.—Unknown.

HABITAT.—Tree-holes and hollow bamboos.

DISTRIBUTION.—Karwar* (as above). BOMBAY DECCAN: Nagargah*, viii. 1921 (*Barrand*).

Not known from elsewhere.

† Barrand 1923, p. 484.

70. *Aedes (Finlaya) assamensis* (Theobald), 1908.

Rec Ind Mus II, p 290 (*Stegomyia*) (♀) TYPE-LOC : Assam, Sylhet, iv 1905 (*Hall*) TYPE ♀ in Ind Mus

ADULT * (Pl. I, figs. 11, 12, and Pl II, fig. 1).—Very similar to *deccanus* and *feegraderi* in the ♀, and to *gubernatoris* in ♂, but differs from these as indicated in key It is rather smaller than the three referred to *Hypopygium* does not show any very marked differences from allied species

LARVA † (fig 36, c).—Very similar to *deccanus* and *gubernatoris*, but differs from both in arrangement of frontal hairs, form of preopyleal spines, and length of antenna, also in having *isc* with but 2 branches, which are more than twice length of segment (compare fig 36, a & c, and *vide* remarks under larva of *A (F) gubernatoris*)

HABITAT.—Tree-holes

DISTRIBUTION.—Common in ASSAM and in NORTH and EAST BENGAL It almost certainly occurs in Burma, but there are no records

Known also from COCHIN CHINA (*Borel*) and JAVA (*R. W. Paine*, 1931, ♂ & ♀ in Brit. Mus)

71 *Aedes (Finlaya) cacharanus* Barraud, 1923.

Bull Ent Res XII, p 406 (*Finlaya*) (♂ & ♀) TYPE-LOC : Haflong, Cachar Hills, Assam, vii 1922 (*Barraud*) TYPE ♂ & ♀ in Brit. Mus

ADULT ‡ (Pl I, figs 7, 8) —Very similar to *cogilli*, *feegraderi*, and other black and white species of the subgenus, but may be distinguished on the following combination of characters :—*Head* with a narrow median white line White patch on front of mesonotum round as in *cogilli*. Prealar white patches fairly large, a small patch of flat white scales on posterior margin of *ppn*, all femora with small white knee-spots, fore tibiae entirely pale beneath for whole length, except at extreme tip, all tibiae with a white mark dorsally near tip; fore tarsi dark, or with a few pale scales at base of segments 1 and 2, mid-tarsi with basal white ring to segment 1, and a fairly broad apical white ring continued on to segment 2 as basal white ring; similar rings on hind tarsi, and some pale scales at joint between segments 2 and 3

♂.—Palpi about length of proboscis, terminal segments with moderate hair-tufts. Mesonotum almost entirely covered with white scales. Upper surface of head mainly white. Other markings as in ♀. *Hypopygium* : style (fig. 35, b)

* Barraud 1923 b, p 408, and 1923 g, p. 486; Borel 1928 b, p. 54.

† Barraud 1923 e, p. 222, Borel 1928 b, p. 56.

‡ Barraud 1923 g, p. 485

with very long terminal appendage, $\frac{3}{4}$ — $\frac{3}{2}$ length of style. Other structures very similar to allied species

LARVA.—Unknown.

HABITAT.—Tree-holes.

DISTRIBUTION.—ASSAM Hailong* (as above), Nongpoh*, Khasi Hills dist., vii. 1922 (Barraud) N BENGAL: Meenglas, Jalpaiguri, vii 1922 (M. O. T. Iyengar)

Not recorded from elsewhere

72. *Aedes* (Finlaya) *lophoventralis* (Theobald), 1910.

Rec Ind Mus iv, p 13 (*Pseudocarrollia*) (♀) TYPE-LOC: Purnea, Bihar (formerly Bengal), viii 1907 (C. Parva). TYPE ♀ in Ind Mus

ADULT † (Pl. II, fig 5)—A rather large and ornate black and white species Wing 3.5–4 mm

♀.—*Head*: a narrow white border to eye-margins, remainder covered with black flat scales and numerous dark upright scales extending from nape on to vertex. Eyes touching in front. Tori, clypeus, palpi, and proboscis black, flagellum of antenna dark brown. *Thorax*: a large round patch of silky white scales covering anterior part of mesonotum, a small white area in front of each wing-root, otherwise scales are brownish-black. Flat white scales on scutellar lobes. A line of broad silvery scales on *apn* and a small patch on lower part of *ppn*, three small and two larger patches of silvery scales on pleurae. *Wings*: dark scaled. *Legs*: black, with white markings. Fore femur dark above, pale at base beneath; mid-femur with a white ventral streak on anterior surface on basal $\frac{1}{2}$ and white mark at knee, narrowly pale posteriorly at base; hind femur with narrow basal and much wider subapical black rings, otherwise white. Tibiae entirely dark except for white dorsal mark at tip on all legs. All tarsi with pale scaling or a pale ring at base of segment 1, and a pale ring over joint between 1 and 2; on mid-leg this ring is sometimes very wide, including the whole of 2, and extending on to 3, or the middle of segment 2 may be dark, with another white ring over joint between 2 and 3. *Abdomen*: black; tergites with lateral basal or medial silvery markings not usually produced on to dorsum; sternites III–V with basal white bands, outstanding tufts of black scales on sternites V–VII only (this seems to be characteristic of the species.)

♂.—Palpi black, a little shorter than proboscis, last two segments turned downwards and with small tufts of black

† Theobald 1910 b, p. 186 (*Pseudocarrollia*); Edwards 1917, p. 214 (*Ochlerotatus*); Barraud 1923 g, p. 487.

hairs. Antennæ strongly plumose. First two segments of mid-tarsi usually mainly white. Scale-tufts on sternites as in ♀. *Hypopygium* coxite shorter and more rounded at apex than in *gubernatoris*, otherwise very similar to that species

LARVA —No specimens available for description

HABITAT —Tree-holes, sometimes water-butts

DISTRIBUTION —Common in NORTH BIHAR from March to October, and known also from EASTERN BENGAL (Chittagong, vii. 1922 (*Barraud*)), but not recorded from elsewhere

73. *Aedes* (*Finlaya*) *khazani* Edwards, 1922

Ind Journ Med Res x, p 265 (♂ & ♀) TYPE-LOC Pudupadi,
Malabar Coast, xi. 1915 (*Khazan Chand*) TYPE ♂ & ♀ in
Brit Mus

ADULT * (Pl I, figs 9, 10, and Pl II, fig. 2) —A black and white species of moderate size, recognized by very long projecting scales on venter, as shown in Pl II, fig 2, and by absence of scales from *ppn*. The only other known Indian species which has very similar long projecting scales on venter is *A. (F.) prominens*, but in this there is a patch of white scales on *ppn*. The following characters are present in *khazani* —Scutellar scales in ♀ flat and dark brown, in ♂ white, on all lobes. Hind tarsi with a wide white ring over joint between segments 1 and 2 ♂ *hypopygium* see fig. 35, c

LARVA (one damaged skin available) —*Frontal hair* *A* quite small, with 4-5 fine branches, *B* single and placed some distance anterior to *A*, *C* slightly posterior to level of *A*, and almost in line behind *B*, actual hair missing; *d* anterior and slightly internal to *C*, with 4-5 small fine branches, *e* with 2-3 fairly long fine branches. Antennal shaft smooth, with single hair very near tip, shaft about 10 times length of width at base. Mouth-brush hairs simple. Lateral hairs of *thorax* well developed. No large spines, but some small hairs with 6-8 branches. *Abdomen*: I and II with fairly long 4-5-branched hairs, following segments with moderately long 2-branched hairs. Comb of fairly numerous teeth, each ending in a number of fine points. *Siphon* brown, with black ring at base, length about 1 mm.; pecten of 15 fairly short strong teeth, each with one large lateral denticle and one or two smaller, tuft missing, its base close to most distal pecten-tooth. *Anal segment* enclosed, except ventrally, within a chitinous saddle with some short strong spines on posterior border towards dorsum; *isc* of about 8 fairly long, subequal branches, a little more than

* *Barraud* 1923 *g*, p 488; *Borel* 1928 *b*, p. 57.

twice length of saddle, *lh* of 3 fine branches. Fan moderately developed, about 10 hairs arising from plate, each split into a number of branches. Papillæ apparently shorter than saddle, ends bluntly pointed.

HABITAT.—Tree-holes.

DISTRIBUTION.—MALABAR COAST * (as above) NORTH KANARA · Karwar *, ix 1921 (*Barraud*). EASTERN BENGAL : Rangamatti * (Chittagong Hill Tracts), ix 1922 (*Barraud*). NORTH BENGAL Sukna *, Darjeeling dist., x 1922 (*Barraud*). Recorded also from COCHIN CHINA (*Borel*).

74 *Aedes* (Finlaya) *prominens* Barraud, 1923.

Ind Journ Med Res xi, p 228 (*Finlaya*) (♂ & ♀). TYPE-LOC : Sukna, Darjeeling dist., ix 1922 (*Barraud*). TYPE ♂ & ♀ in Brit. Mus.

ADULT ♀ †.—Very similar to *khazani*, having tufts of very long outstanding scales on venter of abdomen and roughened scales on dorsum, but differs in having a patch of white scales on *ppn* and some white scales on lateral lobes of scutellum. On hind leg there is white scaling over tibio-tarsal joint, over joint between tarsal segments 1 and 2 (usually narrower than in *khazani*), and at base of 3.

♂ —*Hypopygium* appendage of style rather more than $\frac{1}{2}$ length of latter, blade of harpago much more expanded in middle than in *khazani*.

LARVA.—Has not been isolated in India, but has been described and figured by Borel (1928 *b*, p. 59). It is apparently very similar to that of *assamensis*, but the pecten-teeth, of which there are 12–18, are rather shorter. Siphon rather less than twice length of diameter at base. Comb of about 45 teeth. Arrangement and character of the clypeal hairs not stated.

HABITAT.—Tree-holes and bamboo-stumps.

DISTRIBUTION.—NORTH BENGAL · Sukna * (as above), Tindharia *, Darjeeling dist., ix 1922 (*Barraud*). ASSAM : Nongpoh *, Khasi Hills dist., vii 1922 (*Barraud*).

Recorded also from COCHIN CHINA (*Borel*).

75 *Aedes* (Finlaya) *inquinatus* Edwards, 1922.

Ind. Journ Med. Res. x, p. 267 (♂ & ♀). TYPE-LOC.: Simla, vii. 1915 (*Christophers*). TYPE ♂ and allotype ♀ in Brit. Mus.

ADULT † (Pl. II, fig. 4).—Similar to species of *gubernatoris* group, but mesonotum mainly ochreous-scaled. Wing about 4.5 mm.

† Barraud 1923 *g*, p 489; Borel 1928, p 57

† Barraud 1923 *g*, p. 490.

♀.—*Head*: dorsal surface mainly covered with flat yellowish scales, an indistinct darker patch on either side towards the front, flat white scales laterally and upright scales on nape. Antenna, clypeus, palpi, and proboscis dark brown. *Thorax*: mesonotum in front covered entirely with ochreous scales, posteriorly with dark brown scales, the latter, in some specimens, continued forwards as a pair of submedian broad lines to front margin, a patch of pure white scales in front of each wing-base. In one specimen the dark brown scales are more numerous, the ochreous scales being reduced to patches on front margin, a median line, and lateral curved lines over each wing-root. Fairly broad brown and yellow scales on mid-lobe of scutellum, narrower yellowish scales on lateral lobes. Broad white scales on *apn* and on larger part of *ppn*. Patches of broad, flat white scales on pleuræ. *Wings* dark scaled. *Legs*. fore femur dark anteriorly, mid-femur with a pale streak ventrally on anterior surface for rather more than basal $\frac{1}{2}$ and a pale marking near knee, both pairs pale on posterior surface on basal $\frac{1}{2}$, hind femur with a narrow dark ring near base and a wider sub-apical ring, remainder white. Fore tibia dark brown, faintly paler at apex, very small marking at base of segments 1 and 2 of fore tarsi, mid and hind tibiae dark brown, with pale spot near apex, a pale basal band to tarsal segment 1 and a pale ring over joint between 1 and 2, on hind leg there may be some pale scaling at joint between 2 and 3. *Abdomen*: dorsum brownish-black, lateral basal white patches to tergites. Sternites broadly white basally, with black apical bands.

♂.—Scales on dorsum of head flat and nearly all pale yellow. Palpi and proboscis brownish-black, the former about $\frac{2}{3}$ length of latter. Mesonotal scales as in ♀, but ochreous or pale scales predominating. *Hypopygium*. very similar to that of *A. (F) gubernatoris*, but blade of harpago only slightly expanded in middle.

LARVA.—Unknown

HABITAT.—Tree holes

DISTRIBUTION.—WESTERN HIMALAYAS Simla * (as above).
NILGIRI HILLS *, x 1915 (*Khazan Chand*).

Not known from elsewhere.

76. *Aedes (Finlaya) uncinatus* Edwards, 1922.

Ind Journ Med. Res x, p 266 (♂). TYPE-LOC.: Simla, viii 1915 (*Christophers*). TYPE ♂ in Brit Mus

ADULT † (Pl. I, figs. 13, 14).—Distinguished by the single white ring on hind tarsi, dark proboscis, and ornamentation of mesonotum. Wing about 4 mm

† Barraud 1923 g, p 490

♀.—*Head*: dorsal surface dark except for a median pale line on vertex and pale margin to eyes, some pale upright scales on nape. Antenna, clypeus, and proboscis black, palpi about $\frac{1}{2}$ length of proboscis. *Thorax* mesonotum black, marked with pale yellowish scales (Pl. II, fig 4). Scutellar scales mostly narrow and yellowish, some dark ones on mid-lobe. Broad white scales in irregular patches on pleuræ, a large patch on *ppn*, and a line on *apn*. *Wings* dark scaled. *Legs*. black, with white markings, fore femur white on posterior surface on basal $\frac{1}{2}$, mid-femur narrowly pale on anterior surface, ventrally, on basal $\frac{1}{2}$, a pale streak for whole length behind, hind femur white, except for narrow basal dark ring and much wider preapical dark ring. Tibiæ black except for white dorsal mark at tip on fore leg. Tarsi black, except for basal white ring to first segment on mid and hind legs. *Abdomen* dorsum usually black, with small lateral basal white patches, sternites with narrow basal pale bands.

♂.—Palpi a little shorter than proboscis, fairly slender, black, with few outstanding hairs on terminal segments. Head mainly covered with flat dull white scales. Mesonotum uniformly covered with pale ochreous scales, sometimes an indistinct darker area on either side towards front. Abdomen usually marked with narrow pale basal bands on dorsum. All tibiæ with small apical white mark in some specimens. *Hypopygium* (fig. 35, d) coxite with small lobe at base of dorsal border bearing a dense collection of long bristles which project at right-angles to coxite. Style rather short, with terminal appendage about $\frac{2}{3}$ length of style, blade of harpago expanded in middle, much longer than stem. Phallosome more strongly chitinated than usual, not divided into lateral plates, but with several teeth, or serrations, at apex on each side.

LARVA * (fig 38, b).—*Frontal hair A* of 2 rather short branches, *B* single, at about level of base of antenna, *d* very small, close to *B*, with several fine branches, *C* posterior to *A* and *B*, single and fairly long, *e* single. Preclypeal spines unusually stout. Median hairs of mouth-brushes with fairly large teeth. *Antenna* rather short, about 10 times length of greatest width, shaft smooth, a single hair further from base than apex. Lateral hairs of *thorax* and *abdomen* moderately developed, no stellate hairs or special structures on dorsum. Comb of about 30 scale-like fringed teeth in a triangular patch. *Siphon* dark brown, lighter on about apical $\frac{1}{2}$, as in *A. (F.) oreophilus*, about $2\frac{1}{2}$ –3 times length of diameter at base. Pecten of 10–12 unusually broad, short

* Barraud 1923 h, p. 499

teeth, hair-tuft of usually 4 branches, the base at about middle of tube. Chitinated saddle of *anal segment* enclosing rather more than $\frac{1}{2}$ the segment in side view, with strong spines on posterior margin, *lh* single and fine, *isc* 3-branched. Fan fairly well developed, each hair split into 2-3 branches. In addition to hairs attached to fan-plate, there are several small hairs arising from membrane towards base of segment. Papillæ a little longer than segment, ends pointed.

HABITAT—Tree-holes

DISTRIBUTION—W HIMALAYAS Simla *, viii 1915 (*Christophers*), Krol Mt. *, near Solan (Kalka-Simla road), ix 1924 (*Barraud*). E HIMALAYAS Kurseong *, ix. 1922 (*Barraud*)

Not recorded from elsewhere

77. *Aedes (Finlaya) alboeinctus* Barraud, 1924.

Ind Journ Med Res xi, p 1002 (♂ & ♀) TYPE-LOC Solan, Western Himalayas (Kalka-Simla road), 4,000', viii 1923 (*Barraud*) TYPE ♂ & ♀ in Brit Mus

ADULT.—Rather small species, with mesonotum largely white scaled and three hind tarsal segments basally banded. Wing about 3 mm

♀—*Head* dorsal surface covered with flat creamy scales, a darker area on either side of middle towards front, some pale upright scales on nape. Antennæ, palpi, clypeus, and proboscis dark brown or black, palpi about $\frac{1}{2}$ length of proboscis, latter about length of fore femur. *Thorax*. mesonotum mainly covered with a large undivided area of narrow white scales, continued back from the front, where they have a faint yellow tint, nearly to level of wing-roots, and extending posteriorly in faint lines to scutellum, the space between these lines dark brownish-black. Scutellar scales pale and narrow on all lobes. Some rather broad pale scales on *apn*, narrower yellowish scales on *ppn*. Pleuræ with irregular patches of broad white scales. *Wings*: dark scaled. *Legs*. all femora pale at base, fore femur dark anteriorly, broadly pale from base posteriorly, mid-femur dark, with a narrow pale area ventrally on more than basal $\frac{1}{2}$ on anterior surface, more extensively pale posteriorly; hind femur pale on both surfaces for more than basal $\frac{1}{2}$, no knee-spots. Tibiæ and tarsi brownish-black, former narrowly pale at base, narrow basal pale ring to segments 1 and 2 of fore and mid-legs, and to first three segments of hind legs. *Abdomen*: dorsum almost black, with very narrow pale basal bands, widening out at sides (in many specimens dorsum appears entirely dark, owing to slight shrinkage); tergite VIII and sternite VII pale.

♂—Palpi very slightly shorter than proboscis, thin and black, terminal segments hardly perceptibly thickened, turned downwards and bearing sparse hair-tufts. Other markings as in ♀, but fore and mid-tibiae more or less striped with pale scaling for whole length posteriorly. Distinguished from ♂ of *gilli* by absence of pale scales at base of apical segment of palpi, and by more extensive white area on hind femur, rather more than basal $\frac{2}{3}$ being pale. *Hypopygium* does not show any marked modifications.

LARVA (fig. 38, a).—This has several unusual characters, notably in the development of the frontal hairs and in the form of the pecten-teeth, the latter resembling those of *Uranotænia* larvæ. *Frontal hairs* A, B, C, and d all well developed and all about same size, with 9 or 10 rather long subplumose branches, hair e long, with few lateral branches. *Preclypeal spines* fairly long and stout. *Antennæ* long and curved, about 0.5 mm long, a few small appressed spinelets on shaft, hair-tuft of 3–5 branches attached at about $\frac{5}{8}$ from base on outer side. *Lateral hairs of thorax* long and divided into 6–8 subplumose branches, arising from chitinated tubercles, several pairs of stellate hairs on dorsum with long fine branches. *Lateral hairs of first two abdominal segments* 4-branched, those of following segments 2-branched. *Stellate hairs on dorsum* similar to thorax. *Comb* of about 30 long, pointed, fringed teeth in a patch. *Siphon* dark brown, almost straight-sided, slightly tapering, from 3–4 times length of diameter at base, *acus* small, *pecten* of about 20 close-set scale-like teeth fringed on both sides, usually 6–8 smaller and more pointed teeth irregularly arranged at base, *tuft* of 4–10 finely subplumose branches arising at about middle of tube and slightly beyond most distal pecten-tooth. *Anal segment* with a dark brown chitinated saddle covering rather more than dorsal $\frac{1}{2}$, both saddle and membranous part of segment covered with minute spines set in small groups; *lh* of 4–5 moderately long branches with lateral fraying. *Fan* of about 12 hairs arising from fan-plate, each hair dividing some distance from base into about 6 branches. *Dorsal papillæ* about length of segment, ventral pair shorter, *isc* of 3–5 branches, two of which are very long.

HABITAT —Tree-holes

DISTRIBUTION —W. HIMALAYAS: Solan* and Koti* (Kalka-Simla road), 2–4,000', viii. 1923 and vi. 1930 (*Barraud*), Kakar Hatti*, near Sabathu, viii. 1924 (*Barraud*).

Not recorded from elsewhere.

78. *Aedes (Finlaya) stvensoni* Barraud, 1923.

Ind Journ Med Res xi, p 219 (♂ & ♀) TYPE-LOC. Bombay
Deccan, Nagargah, viii 1921 (*Barraud*) TYPE. ♂ & ♀ in Brit.
Mus

ADULT † —Ornamentation shown in photographs (Pl III, fig 15, ♀, and Pl IV, fig 1, ♂) Differs from *albotæniatus* and other black and white species in having a distinct pale band on proboscis, three rather *small* white spots or patches on mesonotum, and basal white bands on first *three* segments of hind tarsi All the known specimens have the scutellum somewhat denuded, but the scales remaining appear to be black in ♀ and partly white in ♂

♂ —*Hypopygium* (fig 37, a) style comparatively short, with long terminal appendage, harpago with broad leaf-like blade

LARVA —Unknown

HABITAT —Bamboos.

DISTRIBUTION —NORTH KANARA. Kadra *, ix 1921 (*Barraud*), and type-locality * as above

Not known from elsewhere

79. *Aedes (Finlaya) albotæniatus* (Leicester in Theobald), 1904.

Entom xxxvii, p. 111 (♂ & ♀) (*Damelsia*) TYPE-LOC Kuala Lumpur, Malay Penn, from bamboos (*Leicester*) TYPE. ♂ & ♀ in Brit Mus.

Finlaya lepchana Barraud, 1923, Ind. Journ. Med Res xi, p 217 (♂) TYPE LOC Tindharia, Darjeeling dist., x. 1922 (*Barraud*). TYPE ♂ in Brit. Mus.

ADULT ♂ † —Ornamentation shown in photographs (Pl I, fig 15, and Pl II, fig. 6). *Palpi* and proboscis about same length, latter entirely dark brown, former with or without an ill-defined white ring in middle§ and with hair-tufts on terminal segments *Wings*: dark scaled, except for a short white line on anterior edge of oosta at base. *Legs*: fore femur dark above, lined with pale scaling beneath for whole length, mid-femur dark anteriorly, with a pale ventral line on basal ½, pale posteriorly, except for a dark ring near base and a dorsal dark streak, hind femur dark along dorsal edge, a dark ring near base and a subapical dark ring, otherwise white Tibiæ dark brown, pale at bases and more narrowly at tips, fore and mid-pair pale beneath for whole length. Tarsi dark brown, with narrow white markings or rings


† Barraud 1924 a, p 849

‡ Leicester 1908, p. 117, Barraud 1923 g, p 491

§ [Indicated in type ♂ of *lepchana*, but absent in some Indian specimens and absent in type ♂ of *albotæniatus*.]

at base of first two segments on fore and mid-legs, wider white rings at bases of first four segments on hind leg, and a few white scales at tip of segment 1. *Hypopygium*: coxite with long scales and bristles on the internal (ventral) border, dorsal border with a collection of very long scales

Fig 37



♂ hypopygial details (style, harpago, etc.) of *Aedes*, subgenus *Finlaya*: a, *stevensoni*; b, *albotarsatus*; c, *pseudotenebratus*; d, *elatis*; e, *shoriki*; f, *saxicola*; g, *chrysolineatus*; h, *oreophilus*; i, *formosensis*; j, *harveyi*; k, *greeni*; l, *christophersi*. Lettering as on p. 4

or flattened hairs near base. Harpago with leaf-like blade (fig. 37, b)

♀ †.—Ornamentation in Malayan specimens much as in ♂ (with large white patch on middle of mesonotum on anterior $\frac{1}{2}$), but proboscis with a distinct white ring in middle. On tergites VI–VII there are 7-shaped basal lateral white marks

[LARVA.—Not isolated in India. A drawing made by Mr A J E Terzi in 1915 from a specimen obtained at Ginting Simpah, F M S, by Dr A T Stanton shows the following features.—*Comb* of about 14 slender sharp-pointed teeth in a patch, uppermost and lowermost tufts of VIII remarkably large, each with about 15 branches, median tuft of only about 5 branches. Siphon very long for an *Aedes*, fully 6 times as long as its diameter at base, tapered on distal $\frac{1}{3}$ only; pecten of only about 10 almost simple teeth, with one widely detached from the rest, tuft 4-branched, well beyond the detached pecten-tooth; *osc* 4-branched; *lh* long and single. No drawing was made of the head, and the specimen appears to be lost.]

HABITAT.—Bamboos

DISTRIBUTION.—BENGAL: Tindharia * (as above), Sukna *, Darjeeling dist, viii 1928 (*Sobha Ram*). ASSAM: Nongpoh *, Khasi Hills dist, viii 1922 and iii 1927 (*Barraud*)

Known also from MALAY PENINSULA and DUTCH EAST INDIES

[var. *mikranus* Edwards, 1922

Ind. Journ. Med. Res. x, p. 269 (*A* (*F*) *mikranus*) (♀) TYPE-
LOC.—Mikir Hills, Sibsagar dist., Assam, i 1911 (*Christophers*).
TYPE ♀ in Brit Mus

ADULT ♀ †.—(Pl. I, fig. 16, and Pl. IV, fig. 5).—Closely resembles *albotæniatus* ♀ in all respects except that the mesonotum is brown above, lacking the large anterior median white patch. As typical ♂♂ of *albotæniatus* have been found in the same localities with these ♀♀ it is almost certain that *mikranus* is merely a local variety of *albotæniatus*.

LARVA.—Unknown

HABITAT.—Bamboos.]

DISTRIBUTION.—ASSAM. Mikir Hills * (type-locality), Nongpoh *, Khasi Hills dist., xi. 1921 (*Christophers*), and vii 1922 (*Barraud*); Haflong *, Cachar Hills, viii. 1922 (*Barraud*). BENGAL. Tindharia *, Darjeeling dist., ix. 1922 (*Barraud*).

Not recorded from elsewhere.

† See below, under var. *mikranus*.

‡ Barraud 1923 g, p. 491

80. *Aedes* (Finlaya) *maedougalli* Edwards, 1922

Ind Journ Med Res x, p 271 (♂ & ♀) TYPE-LOC. Diyatalawa, Ceylon, in. 1910 (*MacDougall*) TYPE. ♂ & ♀ in Brit Mus

ADULT* (Pl III, fig 14)—Resembles *A* (F.) *elsiae* and *pseudotæniatus*, but differs from both as indicated in key Wing about 4 mm

♀—*Head*. a median line of golden scales from nape to front of vertex and a narrow pale border to eyes, a dark area on either side of middle line towards front, numerous dark upright scales on nape, alternate patches of yellowish and dark flat scales at sides. Antenna and clypeus dark brown, palpi brownish-black, with pale scales in middle and at apices. Proboscis dark at base, mainly pale on about $\frac{2}{3}$ of length both below and above, but usually scattered dark scales along upper surface, apical $\frac{1}{3}$ dark brown. *Thorax* mesonotum brownish-black, marked with narrow lines of golden scales as shown in Pl III, fig 14. Scutellar scales all narrow and golden. Narrow golden scales along upper and posterior margins of *ppn*, patches of broad pale scales on sternopleura and mesepimeron, some smaller pale scales on postspiracular area. *Wings*: a short pale line on outer side of costa at base, otherwise scales are dark. *Legs*: dark brown, with longitudinal lines of yellow scaling on both surfaces for whole length. Fore and mid-tibiae similar, but lines not always well marked anteriorly, hind pair with pale line beneath at base only. Basal and apical pale rings over tibio-tarsal joint, over first one or two joints between tarsi on fore and mid-legs, and over first three on hind legs; all fairly narrow and about same width. *Abdomen*: dorsum black, with silvery basal bands, sternites yellowish, with white basal bands.

♂—Palpi about length of proboscis, dark brown, with a pale ring on long segment and white scaling at base of terminal segment, last two segments with hair-tufts. Proboscis dark brown, with a rather wide pale ring in middle. Other markings as in ♀. *Hypopygium*: style less swollen at base than in *A*. (F.) *pseudotæniatus* and more evenly tapering.

LARVA †.—Very similar to *pseudotæniatus*, differing chiefly in presence of some stellate hairs on abdominal segments. Comb-teeth usually more numerous (50–60), some comparatively short and broad, ending in a number of points. Spines towards dorsum of anal segment strong and with several lateral points (not usually the case in *pseudotæniatus*).

* Barraud 1924, p. 858.

† Senior-White 1927, p. 68.

HABITAT.—Bamboos and rock-pools (*Brug*), filthy rock-pool in ravine (*Senior-White*)

DISTRIBUTION—S INDIA: Nilgiri Hills *, Coonoor (*Patton*). CEYLON. type-locality, as above, and Suduganga *, Matale dist., ix. 1919 (*Senior-White*)

Recorded from SUMATRA (*Brug*).

81. *Aedes (Finlaya) pseudotaeniatatus* (Giles), 1901

Journ Bomb Nat Hist Soc xiii, p 607 (*Culex*) (♂ & ♀) TYPE-
LOC Naini Tal, U P, 7,000' (*Giles*) TYPE ♂ & ♀ in Brit Mus

ADULT † (Pl. III, fig 10, and Pl. V, fig 4) —Recognized by ornamentation of mesonotum, combined with the mainly dark proboscis, striped femora, and apical and basal white rings on hind tarsi

♀.—**Head** a median pale line from nape to front of vertex, a large dark patch on each side, a pale border to eyes, numerous dark upright scales on nape extending forwards some distance. Antennae, clypeus, and proboscis dark brown or black, the last sometimes with some pale scaling on upper surface on basal $\frac{1}{4}$, palpi about $\frac{1}{4}$ length of proboscis, dark, with pale scaling at base and apex. **Thorax** pale lines white or creamy, ground-colour black. Scutellar scales lanceolate, mainly pale, but a few dark. A line of pale lanceolate scales on *apn*, irregular patches of broad white scales on pleurae, scales on *ppn* black, except for a line of pale ones along upper margin, and a patch on lower posterior margin. **Wings**: a short line of white scales on outer margin of costa near base, otherwise dark. **Legs**. dark brown or black, femora and tibiae of fore and mid-legs with narrow longitudinal lines of pale scaling for whole length on both sides; hind femur similarly marked, but tibia with pale line only on about basal $\frac{1}{4}$. Pale rings over tibio-tarsal joints and over joints between tarsal segments 1-2, sometimes 2-3, and occasionally 3-4, on fore and mid-legs, similar but wider rings over joints on hind tarsi except 4-5, apical $\frac{1}{4}$ of 4 and whole of 5 being dark, though 5 may appear light in certain positions. **Abdomen**. dorsum black, with narrow basal pale bands on II-VII, sternites dark, with faint apical yellowish markings in some specimens.


♂.—Palpi slightly shorter than proboscis, last two segments turned downwards and bearing small hair-tufts, white scaling at base of last segment and less distinct pale markings in

† Theobald 1901 a, p 312 (*Stegomyia*), 1907, p. 222 (*Euleoosteomyia*); Barraud 1924 a, p 859. This is most probably the species referred to by some earlier writers as *notoscriptus* Skuse, an Australian species not occurring in India.

middle of long segment, base of penultimate, and tip of last. Other markings as in ♀ *Hypopygium* (fig 37, c): blade of harpago curved and longer than stem; style of moderate length, with a shorter appendage than in most allied species

LARVA* (fig 38, e)—*Frontal hairs* B, C, d placed well forward, anterior to level of bases of antennae and almost in transverse line near middle of clypeus, A of 5 moderately long branches, B and C of 7-9 short branches, d very small,

Fig 38



Larval structures of *Aedes*, subgenus *Finlaya* a, *albocinctus* (head, half mentum, comb-tooth, siphon, pecten-tooth), b, *uncinatus* (head and siphon); c, *pseudocinctus* (head and siphon).

4-5 branched, e usually single and fairly long. Preclypeal spines slender. *Antenna* with smooth shaft and a single hair at about middle, shaft about 10 times length of greatest width. Median hairs of mouth-brushes with small teeth. Lateral hairs of *thorax* and *abdomen* well developed. No stellate

* Barraud 1923 h, p. 502; Senior-White 1927, p. 68.

hairs or special structures on dorsum. Comb of 40-45 rather small fringed teeth in a triangular patch. *Siphon* 2-3 times length of diameter at base, acus fairly large, pecten extending along basal $\frac{2}{3}$ or more, of 21-28 long pointed teeth, all with lateral basal denticles, tuft of 4-6 branches, at $\frac{2}{3}$ from base, and usually between the pectens. Dorsal saddle of anal segment covering about $\frac{1}{2}$ segment in side view, *lh* single and long, *sc* of 2 branches. Papillæ long and pointed, 2-3 times length of segment. Fan of about 12 hairs, each of 2 branches, attached to fan-plate.

HABITAT—Tree-holes, rock-pools, cement sinks, drains, iron cisterns, etc.

DISTRIBUTION—One of the commonest species of the subgenus both in the hills and plains and extending from the NORTH-WEST FRONTIER and BALUCHISTAN to ASSAM and BURMA and throughout the Indian region to CEYLON. It occurs from sea-level up to 7,000' or more.

It has not been recorded from beyond the Indian region *

82. *Aedes (Finlaya) elsæ* Barraud, 1923

Bull. Ent. Res. xii, p. 406 (♂ & ♀). TYPE-LOC Shillong, Khasi Hills, Assam, vi 1922 (Barraud). TYPE ♂ & ♀ in Brit. Mus. *Aedes (Finlaya) simulatus* Barraud, 1931, Ind. Journ. Med. Res. xix, p. 611 (♀). TYPE-LOC Haflong, Cachar Hills, Assam, viii 1922 (Barraud). TYPE ♀ in M.S.I. collection, Kasauli.

ADULT † (Pl. III, fig. 1).—Distinguished from other species with similarly striped legs by double median yellowish line on thorax ‡. Wing about 4 mm.

♀.—**Head** dorsal surface mainly dark, a pale border to eyes and pale scales on nape seen amongst numerous upright dark scales. Antennæ and clypeus dark brown. Palpi with white scaling apically. Proboscis dark brown or black on upper surface, with pale scaling at sides in middle, pale beneath for whole length except for a small dark interruption some distance from apex. **Thorax** mesonotum deep brown, marked with narrow lines of yellowish scales, a double median line from front continued back and dividing either

* [Represented in the Philippines by the closely-allied *A. (F.) banksi* Edwards, 1922, which differs chiefly in having the 5th hind tarsal segment white above (black beneath). Probably *A. banksi* is not more than a varietal form of *pseudotenuatus*.]

† Barraud 1924 a, p. 857; Borel 1928 b, p. 51.

‡ [*A. (F.) macfarlanei* Edw. of Hong Kong (recorded also from Sumatra by Brug, and from Cochin China by Borel) is rather nearly allied to *elsæ*, but has proboscis black beneath on distal $\frac{1}{2}$, most of the abdominal tergites with a pair of creamy spots in middle, white line on anterior surface of hind femur not interrupted, and segment 5 of hind tarsi white.]

side of antescutellar space, a pair of submedian lines ending in front of level of wing-roots, and a pair of curved lines from sides in front, continued over wing-roots. Broad pale scales at base of mid-lobe of scutellum, black at apex, narrow pale scales on lateral lobes. Larger median area of *ppn* with black scales, upper margin with narrow yellow and broad white scales, a patch of broad white scales on lower border. Similar scales on *apn*. *Wings*. a short line of white scales on outer side of costa at base, otherwise dark scaled. *Legs* femora and tibiae of fore and mid-legs dark, with narrow pale longitudinal lines running the whole length both in front and behind, hind femur with similar but broader stripes, anterior pale line often narrowly interrupted at some distance from tip, tibia broadly pale beneath only at base. White rings over joint between tibia and 1st tarsal segment on fore and mid-legs, and over the following two or three joints between tarsal segments. Hind leg with similar, but more pronounced, white rings over joints between tarsi, except 4-5, segment 1 with white ring at base, very few pale scales at apex of tibia. *Abdomen* dorsum dark brown, very narrow basal pale bands on II-VII, wider at sides, not always visible on dorsum owing to shrinkage. Venter mainly yellowish.

♂ —Palpi slightly shorter than proboscis, last two segments slightly curved, with hair-tufts beneath, some white scales at base of apical segment. Other markings as in ♀, but tarsal rings on fore and mid-legs often very small. *Hypopygium* (fig. 37, d): style with rather long bristles along apical $\frac{2}{3}$, both stem and blade of harpago of only moderate length and subequal.


LARVA *† (fig. 39, a, c, d) —This has several peculiar characters, *e.g.*, the presence of two pairs of strong spines on dorsum of thorax, and a strong spinulose bristle representing the siphonal hair-tuft. *Frontal hairs* B, C, d in a transverse line towards front at about level of bases of antennae, A slightly posterior, the last with 5-7 branches, B 6, and C 8. Hair d very small and usually single; e also single, and fairly long. Pre-clypeal spines slender. Median hairs of mouth-brushes with small teeth. *Antenna* of moderate length, about 9 times length of greatest width, some small spines

* Barraud 1923 b, p. 502; Borel 1928 b, p. 53.

† [The larva of *A. (F.) macfarlanei* Edw. (recently obtained by Dr R. B. Jackson in Hong Kong) resembles *claus* and *shorti* in having strong spines arising from conspicuous chitinous plates on dorsum of thorax, but differs in having mesothoracic spine double (the two branches subequal) and the more slender metathoracic spine with a short basal barb. Siphonal tuft of 4-5 quite slender branches; see 4-5-branched, *lh* short and single.]

on shaft; hair-tuft at about middle and of 3-5 branches. Hairs on ventral surface of head unusually long and stout. Lateral hairs on *thorax** and *abdomen* well developed and arising from chitinised tubercles, those on abdominal segments mostly 2-branched. Two pairs of long and strong angle spines

Fig 39



Larval structures of *Aedes*, subgenus *Finlaya*. a, c, d, *elenae* (head, thorax, and siphon), b, e, *shortii* (head and siphon).

on thorax, the larger pair on mesothorax, smaller on metathorax, one smaller pair on abdominal segment I, all these have large, chitinised, tubercular bases. Comb of 60-70 fringed

* [The thoracic chaetotaxy of this species is shown in fig. 39, c, and, apart from the extraordinary transformation of hair 1 of meso- and metathorax into a strong spine, may be taken as typical of the genus *Aedes*. All the hairs can easily be homologized with those of *Aedes* (*Ochlerotatus*) as figured by Martini (1923, p. 550), the notation here adopted differs slightly from Martini's, and agrees with that of Puri (1929, 1931), the pleural hairs of each division of the thorax being numbered 9-12, on the mesothorax Martini's hair 9 is numbered 8, and Martini's hair 8 becomes 14. The following points are noteworthy.—*Prothorax*: hairs 1-3 are included in a small plate, 5-6 are very close together, their tubercles confluent; 7 separate, on lateral

teeth in a triangular patch. Subsiphonal tuft of 6-8 frayed branches. *Siphon* 2-3 times length of diameter at base, pecten of 14-19 long sharp teeth, all with quite small lateral denticles, tuft represented by a strong single or bifid spinulose bristle, arising rather more than half-way along tube from base, and usually lying between pectens. Chitinated saddle of *anal segment* covering about $\frac{1}{2}$ the segment, a number of sharp spines on posterior border, *lh* single and rather short, *isc* of 3-4 moderately long branches. Membranous part of segment covered with small hairs. Papillæ very long and tapering to a point, nearly 4 times length of segment. Fan of about 12 hairs, each divided into 2-10 fine branches arising from heavily chitinated fan-plate.

HABITAT —Rock-pools

DISTRIBUTION —ASSAM type-localities, as given above

It has been recorded from COCHIN CHINA (*Borel*) [and a male and female agreeing in all respects with Indian specimens have been examined at the British Museum from Wuch'ang, HUPBEI, CHINA (*Lan-chou Feng*)]

83. *Aëdes* (Finlaya) *shortti* Barraud, 1923*

Bull. Ent. Res. xii, p. 405 (♂ & ♀) TYPE-LOC Shillong, Khasi Hills, Assam, vi 1922 (*Barraud*) TYPE — ♂ & ♀ in Brit. Mus.

ADULT † (Pl. V, fig. 2) —Ornamentation very similar to *A. (F.) elsæ*, differing as follows:—

♀ —Proboscis broadly pale only in middle beneath and at sides. Anterior surface of fore and mid-femora speckled with pale scales tending to form broken lines, posteriorly lined with pale scaling for whole length, hind femur with conspicuous white marking near knee, and a line of pale scaling on basal $\frac{1}{2}$ on both surfaces. Tibiæ dark, narrowly

margin, 8 not visible dorsally; 9-12 definitely ventral in position, only one of this group moderately long, the rest short. *Mesothorax*: hair 1 forms the conspicuous spine, 6 and 7 arise from a small, common plate, 6 forming a large tuft, 7 single, 8 forms a large tuft, 9-12 in a rather small pleural tubercle, 9 forming a tuft, 10 and 11 long and single, 12 minute, 13 small, ventral; 14 latero-ventral (shown dotted in figure). *Mesothorax*: hair 1 forming the strong spine, 2-6 all small and separate; 7 fairly large; 8 small; pleural tubercle lateral in position, with a strong spine, and apparently with only 3 hairs (hair 11 missing?). Compare this figure and description with that of *Tripteroides* (fig. 9, p. 36) and *Culex* (fig. 102, p. 423).]

* [A somewhat damaged ♀ in the British Museum from Gangtek, Sikkim (*Lt.-Col. F. M. Bailey*) apparently represented an undescribed species related to *A. (F.) shortti*, differing in markings of hind femora, the median white patch on the anterior surface, and the white patch on under surface at tip being much more extensive; median yellowish line of mesonotum apparently single.]

† Barraud 1924 a, p. 856.

pale at base and apex, fore pair with a thin pale line posteriorly in some specimens. Fore and mid-tarsi with white markings at base of first three or four segments and sometimes a few pale scales at apices of segments also. Hind tarsi with four white rings at bases of segments 1-4, a few white scales at apices of segments also, not forming definite apical and basal banding.

♂—Palpi about $\frac{3}{4}$ length of proboscis, terminal segments with distinct hair-tufts. Proboscis with a small pale marking in middle, not usually forming a complete ring on upper surface. *Hypopygium* (fig 37, e). style very similar to that of *elsæ*, but with bristles only on apical $\frac{1}{2}$.

LARVA (fig 39, b, e)—Resembles that of *elsæ* in the presence of large strong spines on dorsum of meso- and metathorax and on abdominal segment I, and in most other details of structure, but metathoracic spine more slender and apical tuft of 3-4 strong subplumose or barbed branches.

HABITAT—Rock-pools.

DISTRIBUTION—ASSAM: Shillong*, vi 1922 (*Barraud*), Cherrapunji, x 1920 (*Semor-White*, Brit Mus). KASHMIR: Gulmarg*, x 1923 (*Sinton*), and Aru*, Lidder Valley, ix 1930 (*Barraud*).

Recorded from SUMATRA (*Brug*)

84 *Aedes* (*Finlaya*) *greeni* (Theobald), 1903.

MC III, p. 289 (*Howardina*) (♀) TYPE-LOC Peradeniya, Ceylon (*Green*) TYPE ♀ in Brit Mus

var *kanaranus* Barraud, 1924, Ind. Journ. Med Res xi, p. 850

(♂ & ♀) TYPE-LOC Karwar, N Kanara, ix. 1921 (*Barraud*)
TYPE ♂ & ♀ in Brit Mus

ADULT (Pl. III, figs 5, 6) †.—Distinguished by bright golden scales of mesonotum, white rings extending over tarsal joints, legs not conspicuously lined. Wing 3.5-4 mm.

♀—*Head* dorsal surface with a median band of golden scales from nape to vertex, continued along eye-margins and between eyes, a number of yellow upright scales, chiefly on nape, a dark patch on either side of median line, and white flat scales at sides. Tori pale brown, flagellum of antenna, clypeus, palpi, and proboscis brownish-black. *Thorax* mesonotum ornamented with bright golden and dark brown scales, variable in arrangement. In some specimens the front part is covered with golden scales, and posteriorly there is a pair of submedian dark lines and a dark area over each wing-root, in others there is a fairly large dark area on either side towards the front, and the dark areas posteriorly may be wider, leaving narrow golden lines between, or there may

† Barraud 1924 a, p. 849.

be a pair of submedian dark lines widening out posteriorly, continued to the scutellum. Scutellar scales and those on *ppn* narrow, golden. Postnotum and pleurae black, latter with irregular patches of broad silvery scales, white lanceolate scales on *apn*. *Wings* dark scaled. *Legs* brownish-black, fore and mid-femora broadly pale posteriorly or basal $\frac{1}{2}$, the pale scaling continued as a line to the knee, hind femur white on both surfaces on more than basal $\frac{1}{2}$, except dorsally, small white knee-spots to all femora, those on fore legs minute or even sometimes absent. Tibiae pale beneath at base, fore pair lined with pale scaling for whole length beneath. Fore tarsi dark or with pale scaling at base of segment 1 and a pale streak beneath, mid-tarsi with white ring at base of segment 1, usually a similar ring over joint between 2 and 3, hind tarsi with white rings at base of segment 1 and over joints between following segments, except at apex of 4, 5 mainly pale dorsally. *Abdomen* black, with basal silvery bands, wider on sternites than on tergites.

♂—Palpi $\frac{3}{4}$ length of proboscis, thin and black, with few outstanding hairs. Hind femur in some specimens entirely white on nearly basal $\frac{1}{2}$. White markings on tarsi usually narrower than in ♀, segment 5 of hind tarsi sometimes only very faintly pale. *Hypopygium* (fig 37, k) coxite has a basal lobe, as well as harpago, both connected with the harpagnal fold. The style is shown separately.

var. *kanaranus*—This differs in the markings of the mesonotum, the golden scaling being reduced to narrow lines on a dark ground.

LARVA (fig 40, a) *—Very similar to *christophersi* and *gills* (vide remarks under the former, p 195).

HABITAT—Tree-holes and bamboo-stumps.

DISTRIBUTION.—Common in the forested regions of the SOUTH-WEST of the Indian Peninsula, in the EASTERN HIMALAYAS, and ASSAM HILLS. KNOWN also from CENTRAL PROVINCES and CEYLON.

It has been recorded from JAVA and SUMATRA.

85. *Aedes* (Finlaya) *chrysolineatus* (Theobald), 1907.

MC IV, p 218 (*Howardina*) (♀) TYPE-LOC. Pundalucya, Ceylon (Green). TYPE. ♀ in Brit Mus.

Hulecoetomyia trilineata Leicester, in Theobald, 1904, Entom xxxvii, p. 163 (proce, Theobald, 1901) (♂ & ♀) TYPE-LOC. Kuala Lumpur, Malay Penn (Leicester) TYPE. ♂ & ♀ in Brit. Mus.

ADULT † (Pl. III, fig. 8, Pl. IV, fig. 3, and Pl. V, fig 1).—

* Barraud 1923 h, p. 500; Senior-White 1927, p. 68.

† Theobald 1907, p. 220, Leicester 1908, p. 107; Edwards 1913 b, p 226, and 1922 d, p 466 (syn.); Barraud 1924 a, p. 852; Borel 1928 b, p 46 (mostly as *trilineata*).

Belongs to a group of closely allied species characterized by golden lines on mesonotum and broad basal white rings on first three segments of hind tarsi * Wing about 4 mm

♀—*Head*: pale creamy scaling in middle of vertex, along eye-margins, and at sides, two dark areas either side of middle line towards front, many upright scales on nape extending forwards to vertex. Tori brown, flagellum of antenna dark brown, palpi brownish-black, white-tipped. Proboscis pale beneath from near base for more than $\frac{1}{2}$ the length (but amount of pale scaling variable), the pale scaling produced on to upper side to form a wide medial ring, otherwise brownish-black. *Thorax*: mesonotum deep brown or black, marked with sharply defined narrow lines of golden scales, a median line extending from the front back to scutellum, forking in front of antescutellar bare space, a pair of submedian lines which nearly meet a pair of lines curving from sides, and continued to lateral lobes of scutellum, another line of golden scales from wing-root, continued forwards a short distance. Mid-lobe of scutellum with narrow golden scales in centre, flat dark scales on each side, lateral lobes with narrower dark scales. Pleuræ with irregular patches of flat silvery scales, a line of similar scales on *apn*, many similar scales on *ppn*. *Wings* usually a short line of white scaling on outer or underside of costa at base, otherwise scales are dark. *Legs*. fore femur dark anteriorly except along ventral border, mid-pair with medial pale line along about basal $\frac{1}{2}$, both pairs pale posteriorly for whole length, hind femur dark along dorsal border for whole length, a dark ring near base, broader than in several allied species, and a subapical dark ring, a large white area on each side near middle, anterior white area usually separated from posterior by a narrow ventral line of dark scales, another white streak on outer side ventrally to knee. Fore tibiae indistinctly pale beneath for whole length, mid- and hind pair narrowly pale at bases only. Tarsi of fore and mid-legs brownish-black, with basal white rings to first two or three segments, hind tarsi with three white rings, that on segment 3 widest and occupying about basal $\frac{1}{2}$ of segment. *Abdomen*: almost black, with yellowish basal bands (sometimes reduced


* [Oriental species belonging to this group which have not yet been identified in the Indian area are —(1) *A. (F.) jugraensis* Leio (Malaya), with median golden line of mesonotum running back almost to scutellum without forking, sublateral lines tending to be obsolete, hind femur with comparatively small pale area in middle, (2) *A. (F.) rizali* Banks (Philippines), with median line of mesonotum running back almost to scutellum without forking, hind femur with basal $\frac{1}{2}$ nearly all whitish; (3) *A. (F.) japonicus* Theo (Japan and China), with mesonotal lines broader, less golden, median line tending to be double, hind femur conspicuously white all round at tip, hind tibia dark at base beneath]

or absent), and large basal lateral silvery spots. White basal bands on sternites, sternite VIII yellowish.

♂.—Palpi slender, upturned, without hair-tufts, a little shorter than proboscis, with three faint pale rings. Proboscis pale for whole length beneath, a narrow pale ring in middle *Hypopygium* (fig 37, g). blade of harpago curved, tapering, a little longer than stem. Style with terminal appendage a little less than $\frac{1}{3}$ of its length.

LARVA *† (fig 40, c)—*Frontal hairs* B, C, d almost in a transverse line slightly anterior to level of bases of antennæ, A and B with 6-7 branches of only moderate length, C 5-6-branched, d very small, and usually divided into 2 branches,

Fig 40



Larval structures (siphon and antenna) of *Aedes*, subgenus *Finlaya*: a, *greeni*, b, *saxicola*, c, *chrysolineatus* (with enlarged comb-scale)

c single and fine. Shaft of antenna with very few minute spicules and a 2-branched hair at about middle. Hairs on thorax and abdomen well developed, many on dorsum

* Barraud 1923 h, p. 500, Senior-White 1927, p. 68, Borel 1928, p. 47

† [The larva of *A. (F.) japonicus* Theo. (recently obtained by the isolation method by Dr. R. B. Jackson in Hong Kong) resembles *chrysolineatus* in many respects, having 6-8 long, fine branches to the dorso-lateral metathoracic tuft. It differs conspicuously in having a siphon more like *saxicola*, with 2-4 extremely large pecten-teeth beyond the tuft, tip of last tooth almost reaching apex of tube. On the other hand the larva of *A. (F.) koreicus* Edw., according to specimens obtained by Drs. H. E. Meleney and C. U. Lee in the neighbourhood of Peking in 1926-8, is much more like that of *A. (F.) pulchripes*, having all the pecten-teeth alike, more beyond the tuft, and none of the distal ones simple or obviously separated from the rank. These striking differences clearly show how plastic are the larval characters in this group; the adult of *koreicus* is so similar to *japonicus* that it was placed by Edwards, 1932, as a variety of that species.]

being of stellate form, no large spines on thorax, dorso-lateral hairs of metathorax with 6-8 branches, arising from moderately large tubercle. Comb of 26-30 fringed teeth in a triangular patch. *Siphon* 2-2½ times length of diameter at base. Pecten of 16-19 teeth, with two or three lateral denticles, except the two or three most distal teeth, which are larger, simple, and placed beyond the tuft, though not nearly reaching tip of tube. Hair-tuft of 5-6 long branches, arising at about ¾ length of tube from base. Chitinous saddle of *anal segment* covering rather more than ½ the segment, posterior border with numerous small spines, *asc* of 3-4 long branches, *lh* 3-4-branched and a little longer than the saddle, 10-12 fan-hairs attached to fan-plate, each hair divided into 3-8 fine branches. Papillæ apparently moderately long and pointed.

HABITAT—Tree-holes, bamboos, rock-pools. Also roof-gutter and broken chattu (*Senior-White*).

DISTRIBUTION—BOMBAY: Deccan, Nagargali *, viii 1921 (*Barraud*); North Kanara, Kadra *, ix. 1921 (*Barraud*); Yellapur *, x 1921 (*Barraud*). MALABAR: Pudukudi *, x 1915 (*Khazam Chand*). E HIMALAYAS: Kurseong, ix 1909 (*Annandale*). CEYLON: Pundaluoya and Peradeniya (*Green*); Suduganga, Matale dist, 1919 (*Senior-White*).

Known also from MALAY PENINSULA †, COCHIN CHINA, and DUTCH E INDIES.

86 *Aedes* (*Finlaya*) *harveyi* Barraud, 1923.

Bull Ent Res. xii, p 407 (♂ & ♀). TYPE-LOC: Kurseong, Darjeeling dist, ix 1922 (*Barraud*). TYPE: ♂ & ♀ in Brit. Mus. var. *nigrorhynchus* Brug, 1913, Arch Hydrobiol. Supp. Bd. xi, Bd 11, p 28 (see under "Larva").

ADULT ‡ (Pl IV, fig 2).—Closely resembles *A. (F.) chryso-lineatus*, differing chiefly as follows.—Proboscis (in type-form) indistinctly pale beneath in middle only, sides and upper surface dark (in other specimens the pale scaling is more distinct and extends further towards base, in var. *nigrorhynchus* as described by Brug the proboscis is entirely dark), *ppn* less scaly, flat white scales confined to lower posterior corner, scales on upper part narrow and yellowish. Fore and mid-femora with few or no pale scales on anterior surface. Hind femur at tip and hind tibia at base less

† [Malayan specimens differ from those from India and Ceylon in having few or no white scales on anterior surfaces of front and middle femur, and apparently no basal bands to abdominal tergites in ♀. In other respects (e. g., scaling of proboscis and *ppn*) they are closely similar to Indian specimens.]

‡ Barraud 1924 a, p. 855; Brug 1931 b, p. 26

extensively white beneath. Dorsum of abdomen in ♀ entirely dark, with no trace of pale bands, though tergites have large lateral basal silvery spots (white basal bands present in ♂) ♂ style and harpago as in fig 37, g

LARVA.—This has not been isolated in India. [The Javan larva described by Brug (1931, p 27) as *A. (F.) harveyi* var *nigrorhynchus* may belong here, though this is not quite certain† According to Brug's description and figures this larva is similar to the one described below as *A. (F.) formosensis*, but has a rather longer siphon, with no pecten-teeth beyond the tuft, *isc* 3-branched, *lh* 2-branched, etc.]

HABITAT (in Java).—Tree-holes (*Brug*)

DISTRIBUTION.—EASTERN HIMALAYAS Kurseong*, Mung-poo*, Sureil*, Tindharia*, Sukna*, all in Darjeeling dist., ix & x. 1922 (*Barraud*), MADRAS Coonoor, ix 1920, on bull (*Senior-White*) One ♀ from Mercara, Coorg, SOUTH INDIA, vi 1927 (*J. D. Baily*), appears to be this species

Recorded from DUTCH EAST INDIES (*Brug*).

87. *Aedes (Finlaya) formosensis* Yamada, 1921.

Annot Zool Jap x, p 67 (♀) TYPE-LOC Kakubanzan, Formosa, v 1921 (*Hayama*) TYPE. 2 ♀♀ co-types, location unknown

Finlaya khamana Barraud, 1923, Bull Ent Res xii, p 407 (♂ & ♀).

TYPE-LOC Shillong, Khasi Hills, Assam, vi 1922 (*Barraud*)

TYPE ♂ & ♀ in Brit Mus

ADULT ♀‡.—Closely resembles *A. (F.) chrysolineatus*, differing chiefly as follows.—Proboscis with the pale scaling extensive but confined to ventral and lateral surfaces, and scarcely visible from above. Mesonotum with the golden lines not quite so sharply defined. Scales on *ppm* all narrow and pale yellowish. Whitish lines on anterior surfaces of fore and mid-femora sometimes more definite, but apparently variable. Hind femur in types (but apparently not always) with a few dark scales at extreme tip beneath, beyond the white mark, dark ring at base narrower, only about twice diameter of femur. Abdominal tergites with median basal pale spots separate from the lateral spots, latter smaller and less silvery white; sternites more extensively dark.

♂.—Palpi slender, upturned, without hair-tufts, brownish-black, with a definite white ring on long segment and smaller rings at bases of last two segments. Proboscis with a pale area beneath and at sides rather beyond middle, no other pale scaling. Other markings as in ♀. *Hypopygium*. does not show any marked modifications: harpago as in fig 37 f

† See fo

‡ Barra

[LARVA.—Not isolated in India. Specimens from Bali † show the following characters.—*Head* dark, including antennae, which are rather long and almost smooth. *Frontal hairs* well forward, *A* with about 5 branches, *B* 6, *C* 8, *d* 3, *B* and *C* almost side by side, *d* small. *Thorax* without conspicuous structures, dorso-lateral tuft of metathorax with 6–7 very long branches, arising from small inconspicuous plate. *Comb* of 25–40 fringed scales, variable in shape. *Siphon* very short, black, pecten of about 10 teeth with strong basal denticles, one simple tooth (not specially large) beyond hair-tuft (according to Brug a second simple tooth may be present at extreme tip of tube), tuft at $\frac{2}{3}$, 3-branched. *Anal segment* with strongly spinulose hind margin of saddle, *isc* of 2 long branches, *lh* single. *Papillae* about twice length of saddle.]

HABITAT—Bamboo-stumps (*Barraud*), leaf-axils of *Colocasia* and other plants (*Thenemann*, *Brug*)

DISTRIBUTION—ASSAM Shillong*, vi 1922 (*Barraud*), and viii 1928 (*T. B. Fletcher*), Golaghat*, xii 1924 (*Barraud*), Nongpoh*, Khasi Hills dist., xi 1921 (*Christophers*). E HIMALAYAS Surel*, x 1922 (*Barraud*), Kurseong*, viii 1928 (*Sobha Ram*)

Originally described from FORMOSA, subsequently recorded from DUTCH E INDIES (*Brug*).

88 *Aedes (Finlaya) pallirostris* Edwards, 1922

Ind Journ Med Res x, p 270 (♀) TYPE-LOC Assam, Golaghat, Sibsagar dist., ii 1911 (*Christophers*) TYPE ♀ in Brit Mus.

[ADULT ♀ †.—Closely resembles *A. (F.) formosensis*, of which it may be only a variation, but proboscis mainly pale on basal $\frac{2}{3}$, with very few dark scales even on upper surface, mesonotal golden lines as clearly defined as in *chrysolineatus*, the sublateral lines definitely broken in middle, and some pale scaling towards hind margins of sternites as well as at base. Hind femur in type with some black scales

† [Briefly redescribed above from two isolated skins in British Museum from the series described by Brug (1931, p. 26) as *A. (F.) harveyi*. The corresponding adults, however, are not *harveyi*, but apparently *formosensis* (*khasiana*). In reporting this error of determination to Col Brug I unfortunately referred to the specimens in question as being from Sumatra, whereas they are from Bali. Perhaps misled by my error regarding the locality, Brug has published a note (1932, p. 390) stating that of the material previously determined by him as *A. (F.) harveyi*, specimens from Sumatra are *khasiana*, those from Java and Bali *harveyi*. It is, perhaps, more likely that those from leaf-axils (Sumatra and Bali) are *formosensis*, those from tree-holes (Java) are *harveyi*, but the matter requires re-investigation. There are no Sumatran or Javan specimens in the British Museum.—F. W. EDWARDS.]

‡ *Barraud* 1928 a, p. 855.

at extreme tip beneath and with narrow black basal ring as in types of *khasiana*]

♂ and LARVA —Unknown

DISTRIBUTION —Known only with certainty from the type-locality.

89. *Aedes* (Finlaya) *saxicola* Edwards, 1922.

Ind Journ Med Res x, p 466, nom nov for *H. fluviatilis* Leio
Hulcesteomyia fluviatilis Leicester, 1908 (neo Lutz), Cul Malaya,
 p 111 (♂ & ♀) TYPE-LOC Ulu Gombak, Malay Penin
 (Leicester) TYPES non-existent
Finlaya greigi Barraud, 1923, Bull Ent. Res xii, p 406 (♂ & ♀)
 TYPE-LOC Haflong, Cachar Hills, Assam, viii 1922 (Barraud)
 TYPE ♂ & ♀ in Brit Mus

ADULT * (Pl III, fig 2) —Differs from *chrysolineatus* as follows —Scales in middle of vertex more golden. Proboscis black both above and below. Palpi much more narrowly white-tipped. Mesonotum with the sublateral golden lines on anterior $\frac{1}{2}$ shorter and much less definite, median line rather deeper golden, and broader than in *chrysolineatus*. Scales on *ppn* fewer and confined to posterior margin, narrow and golden above, broader and white below. Sternite VIII of ♀ blackish. Front and mid-femora all dark on anterior surface, or with only indistinct traces of a whitish line towards base, hind femora much less extensively white beneath at tip, hind tibiae also rather less extensively white beneath at base.

LARVA †† (fig 40, b) —Frontal hairs and antenna much as in *chrysolineatus*. Lateral hairs of thorax and abdomen very strong and subplumose and arising from chitinised tubercles, a dorso-lateral pair of very strong 4-branched spinulose bristles [representing hair 7] on metathorax, [arising from large chitinised plate, hair 1 on mesothorax fairly slender, bifid, and almost simple, arising from small chitinised tubercle], a number of strong, spinulose, single bristles on abdomen. Comb of about 60 fringed teeth in a triangular patch. Siphon 2.5–3 times length of diameter at base. Acus fairly large. Pecten of about 20 teeth, those between base of siphon and hair-tuft with lateral denticles,

* Edwards 1923, p. 3, Barraud 1924 a, p 851

† Barraud 1923 h, p 501, Edwards & Given 1928, p. 344.

†† [The larva of *A. (F.) jugraensis* (found in fallen leaves in forest in Malaya) resembles *saxicola* in having hair 7 of metathorax composed of 3–4 remarkably strong spine-like and barbed branches, and arising from a large chitinous plate, but differs in having no simple pecten-teeth beyond hair-tuft on siphon, hair 1 of mesothorax in *jugraensis* is single and stouter than in *saxicola*.]

the 5-6 teeth between hair-tuft and apex very large, strong, and without denticles †, point of last tooth reaching nearly to apex of tube, hair-tuft of 5-6 subplumose branches, arising at about middle of tube. Anal segment very similar to that of *A. (F.) elisæ*.

HABITAT —Rock-pools

DISTRIBUTION —ASSAM Haflong *, Cachar Hills, viii 1922 (*Barraud*) EASTERN HIMALAYAS Kurseong *, Darjeeling dist, c 5,000', ix 1922 (*Barraud*)

Also known from MALAY PENINSULA (*Leicester*, *Given*, *Hacker*) and Mt Salak, JAVA (*R. W. Paine*, 1929)

90 *Aedes (Finlaya) oreophilus* (Edwards), 1916.

Bull Ent Res vi, p 357 (*Ochlerotatus*) (♀) TYPE-LOC Gharua, Murree Hills, W Himalayas TYPE ♀ in Brit Mus (from Agric Res Inst, Pusa)

ADULT ‡ (Pl. III, fig 9, and Pl V, fig 3) —Distinguished by dark tarsi and other characters given in key. Ornamentation of mesonotum of ♀ and ♂ shown in above-mentioned plates. Both sexes may apparently be distinguished from *A. (F.) suffusus* by the *ppn* scales being nearly all broad and flat instead of mostly narrow.

♂.—Differs from ♀ in having dorsal surface of head mainly white scaled, without lateral dark patches, mesonotum sparsely covered with pale yellow scales not arranged in lines, palpi a little shorter than proboscis, black, with small tufts on terminal segments. *Hypopygium* (fig. 37, h) coxite with dense collection of hairs on about middle of internal border, harpago elbowed at about middle, apical part tapering to tip, style of moderate length, with long terminal appendage.

LARVA (fig. 41, a) —*Frontal hairs* A, B, and d in a transverse row, slightly posterior to bases of antennæ, all branched; A with 4-6 branches, B 3-5, d quite small, usually 3-branched, C some distance posterior to and directly behind B, fairly long and simple (very rarely split into two), e usually with 2 short fine branches. Preclypeal spines strong, pointed, moderately long. *Antenna* with shaft straight, smooth, a very few minute spinelets visible under high magnification, about 9 times length of greatest width, tuft represented by single hair (rarely split), attached rather nearer apex than base of shaft. Median hairs of mouth-brushes with moderate-

† [In larvae from Mt Salak, Java, the last pecten-teeth are not quite so large as in Indian and Malayan specimens, and there are only 3-4 beyond the tuft, in other respects the larvae are similar.]

‡ *Barraud* 1924 a, p. 360.

sized teeth. Mentum broadly triangular, with 9-12 teeth of equal size on either side of central one. Lateral hairs of *thorax* and *abdomen* moderately developed, no special structures on dorsum. Comb of 22-25 comparatively large fringed teeth in three or four rows in triangular patch. *Siphon* $2\frac{1}{2}$ -3 times length of diameter at base, slightly tapering,

Fig 41



Larval structures (head, half mentum, siphon, enlarged pecten-tooth and comb-scales) of *Aedes*, subgenus *Finlaya* a, *orocophilus* (with enlarged elements of mouth-brush), b, *suffusus*.

dark brown, apical $\frac{1}{2}$ pale. Acus small. Pecten of 13-21 strong pointed teeth, with one or two large denticles towards base and a few hair-like points; tuft of 3-4 subplumose branches arising at about $\frac{2}{3}$ of length from base, and just beyond most distal pecten-tooth. *Anal segment* nearly enclosed by chitinated saddle; *th* 2-branched and about length of segment; *sc* of 3-4 branches of varying lengths,

osc single and long, as usual Fan-hairs strong and black, about 12 arising from fan-plate, each divided into 2-4 branches

HABITAT.—Tree-holes

DISTRIBUTION.—One of the commoner tree-hole breeding species of the subgenus in W HIMALAYAS Found also in E. HIMALAYAS and in Nilgiri Hills, SOUTH INDIA.

Not recorded from beyond the Indian region

91 *Aedes (Finlaya) suffusus* Edwards, 1922.

Ind Journ Med Res x, p 270 (♀) TYPE-LOC. Simla, viii 1915
(*Chrostophers*) TYPE ♀ in Brit. Mus

ADULT ♀* (Pl. III, fig 11) —Resembles *A (F.) oreophilus* very closely, differing as follows.—Dorsal surface of head covered with lanceolate yellowish-white scales, no definite dark areas, numerous dark upright scales on nape Narrow scales on upper part of *ppn*, instead of all scales on this part being broad and flat. Mesonotum covered with yellow and brownish scales, not arranged in definite lines, a pair of indistinct submedian dark lines

♂—Differs from that sex of *oreophilus* in scaling of *ppn*, as above *Hypopygium* does not show any very marked differences from allied species

LARVA (fig 41, b) —*Frontal hairs* *A*, *B*, *C*, and *d* all well developed and branched, *A*, *B*, *d* in transverse row slightly posterior to level of bases of antennæ, *C* slightly internal and posterior to *B*, *A* with 8-11 branches, *B* 5, *C* and *d* 8-10, *e* 2-branched *Antenna* comparatively long and curved, shaft about 14 times length of greatest width; scattered small spinelets along its length; hair-tuft of 5-6 branches at about middle. Mouth-brush hairs simple. Mentum with 9-11 teeth either side of larger median tooth, lateral teeth larger near base than those near apex. Lateral hairs of *thorax* and *abdomen* long and fine, stellate hairs present on dorsum of thorax and abdominal segments. Comb of 8-14 comparatively large, sharply pointed teeth in an irregular row *Siphon* dark brown, about 3 times length of diameter at base; pecten 16-21 teeth along basal $\frac{1}{2}$ of tube, each tooth, with one large and several smaller denticles. Siphonal tuft usually of 3 branches, at about $\frac{1}{3}$ from base. Acus moderately developed. Saddle covering about $\frac{1}{2}$ anal segment in side-view, $\frac{1}{2}$ of 3 fine branches, *isc* 5-6, 3 or 4 times length of saddle, fan of few long branches, fan-plate rudimentary Ventral papillæ about length of saddle, dorsal pair a little longer, both pairs bluntly pointed. The larva is remarkably distinct from that of *oreophilus*.

HABITAT.—Tree-holes.

* Barraud 1924 a, p. 861.

DISTRIBUTION —W HIMALAYAS only Simla *, Krol Mt.*, near Solan (Kalka-Simla road), Fagu * (Hindustan-Tibet road), vii-viii 1923-1930 (Barraud), all 7-8,000'.

92 *Aedes* (Finlaya) *christophersi* Edwards, 1922.

Ind Journ Med. Res x, p 267 (♀). TYPE-LOC Simla, viii, 1915 (*Christophers*). TYPE ♀ in Brit. Mus

ADULT † (Pl III, fig 12).—Characterized by thoracic adornment and three basally-ringed hind tarsal segments. Wing about 4.5 mm.

♀.—*Head*: dorsal surface covered with narrow pale ochreous scales and scattered upright scales, latter chiefly towards nape. Tori brown, flagellum of antenna, clypeus, palpi, and proboscis black, palpi about $\frac{1}{2}$ length of proboscis. *Thorax*: a median line of yellowish scales, continued back from the front and dividing either side of antescutellar space, bordered on either side by a broader band of brownish-black scales; laterally on sides in front scales are all yellowish, each of these areas divided posteriorly into two lines of pale scales, one running to lateral lobe of scutellum and one to wing-root, space between brownish-black. Scutellar scales narrow and yellow. Postnotum and integument of pleurae dark brown or black. Some fairly broad white scales on *apn* and on lower part of *ppn*, narrow yellow scales on upper part of latter, irregular patches of broad silvery scales on pleurae. *Wings*: dark scaled. *Legs*: bluish-black, with white markings, all femora with white knee-spots and white scaling from base, chiefly posteriorly and ventrally, hind femur white on both sides for basal $\frac{1}{2}$ or rather more. Tibiae dark, hind pair narrowly pale beneath at base. Tarsi with narrow basal white markings to first two segments on fore and mid-legs, similar but wider rings to first three segments on hind legs. *Abdomen*: black, with basal white bands to tergites and sternites, latter with some apical white scaling also.

♂ —*Palpi* black, little shorter than proboscis, the terminal segments not perceptibly swollen and with few outstanding hairs. Mesonotum almost entirely covered with ochreous scales, the dark stripes present in ♀ being indistinct or absent. Other markings as in ♀. *Hypopygium*: style and harpago as shown in fig 37, 1

LARVA (fig. 42, b).—Very similar to *A. (F.) gilli* and *greeni*, differing from former in having only one pair of frontal hairs (*B*) single and very long, longer than whole head. Preclypeal spines fairly long and very slender. Antennal

† Barraud 1924 a, p. 861.

hair of 2 branches (each branch again occasionally split into two). Comb of 40-50 teeth. Pecten usually of 19-22 teeth, but occasionally more. Siphonal tuft of 3-5 fairly long fine branches, *usc* of only 2 branches. Anal papillæ with rounded ends, dorsal pair longer than segment.

From *A. (F.) greeni* this larva appears to differ only in the shape of the anal papillæ, the dorsal pair of which in *greeni* are only about the length of the anal segment, the ventral pair shorter, both with pointed ends. The siphonal hair-tuft has usually 5-6 branches in *greeni*.

HABITAT—Tree-holes

DISTRIBUTION—KASHMIR. Gulmarg, 1913 (*Wyville-Thomson*). W. HIMALAYAS. Simla *, viii 1915 (*Christophers*), viii 1923, viii & ix 1924, and ix 1930 (*Barraud*), Fagu * (Hindustan-Tibet road), viii 1923 (*Barraud*), Krol Mt., near Solan (Kalka-Simla road), ix 1924, vii & viii 1930 (*Barraud*), Murree *, 1922 (*Gill*).

Not known from elsewhere

93 *Aedes (Finlaya) gilli* Barraud, 1924.


Ind Journ Med Res xi, p. 1000 (♂ & ♀). TYPE-LOC. Krol Mt. near Solan (Kalka-Simla road), 7,000', viii. 1923 (*Barraud*).
TYPE: ♂ & ♀ in Brit. Mus.

ADULT—Distinguished by ochreous-scaled thorax and basal white rings on first three hind tarsal segments. Wing about 4.5 mm.

♀—*Head*. dorsal surface covered with flat pale yellow scales and some upright scales of same colour, latter chiefly confined to nape, a patch of pale yellowish narrow scales on nape, and a narrow stripe of similar scales extending forwards to eyes, dark flat scales at sides of head. Antennæ, clypeus, palpi, and proboscis brownish-black, palpi about $\frac{1}{2}$ length of proboscis. *Thorax*. anterior $\frac{1}{2}$, or rather more, of mesonotum entirely covered with pale golden or ochreous narrow scales; this area ends irregularly in front of level of wing-roots, posterior part covered with dark brown and yellowish scales, the latter arranged chiefly in three lines to scutellum, a small area of white scales in front of each wing-root. Scutellar scales narrow and yellow. Postnotum brown, integument of pleuræ darker, scales on *apn* rather broad and white, *ppn* almost covered with broad, flat, pale yellowish scales, a few narrower ones on upper margin. *Wings*. dark scaled. *Legs*: bronzy black, marked with dull white, fore femur dark, except for a narrow white ring at base and on basal $\frac{1}{2}$ posteriorly; mid-femur with similar narrow basal white ring and with a ventral pale streak on anterior surface, broadly pale on basal $\frac{1}{2}$.

posteriorly, hind femur pale on both surfaces on about basal $\frac{1}{2}$; all femora sometimes with faint knee-spots, chiefly on posterior aspect Tibiæ dark brown, pale at base, and streaked with pale scaling posteriorly, hind tibia rather

Fig 42.



Larval structures of *Aedes*, subgenus *Finlaya*: a, gills, b, *christophersi*; c, *pulchrior* (with two enlarged pecten-teeth and one comb-scale)

broadly pale at base beneath. Tarsi with pale scaling or pale rings at bases of segments 1 and 2 on fore and mid-legs; wider pale rings on first three segments of hind legs. *Abdomen*: brownish-black, with narrow basal pale bands on segments 2-6 (sometimes invisible in bred specimens owing to shrinkage)

and with basal lateral silvery patches, largest on VI and VII. Sternites pale, with apical dark bands

♂.—*Palpi* a little shorter than proboscis, apical segments slightly swollen, turned downwards, and hairy, some pale scales at base of apical segment dorsally. Scales on head, mesonotum, and scutellum mainly white, not yellow. Other markings as in ♀ *Hypopygium* does not show any marked modifications

LARVA (fig 42, a).—Most nearly resembles *christophersi* and *greeni*. *Frontal hair* A of 2-3 fairly long fine branches, B and C both single and very long, as long as, or longer than, the whole head, which is itself comparatively large. *Maxillæ* large and prominent, mouth-brush hairs simple. *Preclypeal* spines slender and rather short. *Antenna* fairly long and slender, about 12 times length of width, shaft with few fine spicules and a single hair attached at nearly $\frac{2}{3}$ of length from base. *Lateral hairs* on *thorax* and on first two abdominal segments fairly well developed, no special structures on *thorax*. *Abdomen*. hairs on III-VIII small and fine. Comb of about 70 fringed teeth in a patch. *Siphon* deep brown or black, 4-5 times length of diameter at base, pecten of about 24-28 long teeth, with lateral denticles on basal $\frac{1}{3}$, one or two of these large, others much smaller, tuft attached some distance beyond pecten, of 6-9 fine branches, often difficult to see against the dark siphon (in *christophersi* the tuft is larger). Dorsal $\frac{1}{2}$ of *anal segment* covered with dark brown saddle, *lh* of 3-5 fine short branches, *isc* of 6-8 branches of varying lengths, outer single and long as usual. Fan of moderate size, about 10 hairs, each divided into a number of branches, several hairs arising from membranous part of segment. Dorsal papillæ long and pointed, about length of longest fan-hairs, ventral pair about $\frac{1}{2}$ this length

HABITAT.—Tree-holes

DISTRIBUTION.—W HIMALAYAS Krol Mt *, near Solan; Simla *, and Fagu * (Hindustan-Tibet road), vii.-viii. 1923-1930 (*Barraud*), Bhowali *, Naini Tal dist., vii. 1925 (*S Sundar Rao*).

Not recorded from elsewhere

94. *Aedes* (Finlaya) *simlensis* Edwards, 1922.

Ind. Journ. Med. Res. x, p 269 (♀). TYPE-LOC. Simla, vii.-ix. 1925 (*Christophers*). TYPE ♀ in Brit Mus.

ADULT ♀ †.—Differs from *A* (*F*) *gilli* chiefly as follows:—Fewer narrow scales on nape and none extending forwards

† Barraud 1924 a, p. 862.

in middle line to vertex. Scales on *apn* all rather narrow and curved. Hind femur white on rather more than basal $\frac{1}{2}$ on outer side, rather less so on inner side, all knees dark. Fore tibia blackish all round, except narrowly at base. Hind tibia more narrowly pale at base beneath.

♂ and LARVA —Unknown.

HABITAT —Tree-holes

DISTRIBUTION —W HIMALAYAS only, not common. Simla* (*Christophers*), Lansdowne*, ix 1923 (Ind Stn Hosp.), Bhowali*, vi 1925 (*S Sundar Rao*), Muktesar*, Kumaon, ix 1922 (*T B Fletcher*), all 7-8,000'

95 *Aedes* (*Finlaya*) *pulehriventer* (Giles), 1901.

Journ Bomb Nat Hist Soc xii, p 608 (*Culex*) (♂ & ♀). TYPE-LOC Naini Tal, W Himalayas (*Giles*) TYPE ♂ & ♀ in Brit Mus

Howardina himalayana Giles, 1904, Journ Trop. Med vii, p 384
TYPE-LOC Naini Tal, W Himalayas TYPE ♂ in Brit Mus

ADULT † —Recognized by dark tarsi and conspicuous orange patches on abdominal sternites. Wing about 5 mm

♀ —*Head* dorsal surface covered with narrow golden scales and numerous dark upright scales, latter chiefly on nape, indefinite dark patches on either side in front and lower down at sides. Tori light brown, flagellum of antenna, clypeus, palpi, and proboscis black, clypeus and palpi with a few golden scales in most specimens, palpi about $\frac{1}{2}$ length of proboscis. *Thorax*. mesonotum covered with golden and brownish-black scales, former arranged chiefly in a wide median area from front, narrowing posteriorly and dividing either side of antescutellar space, and in a curved line each side over wing-roots. Scutellar scales narrow and golden. Postnotum brown, pleurae almost black, with irregular patches of silvery flat scales, pale, fairly broad scales on *apn*, narrow golden scales on upper and posterior margins of *ppn*. *Wings* dark scaled. *Legs* brownish-black, all femora with pale knee-spots, fore femur dark anteriorly, narrowly pale posteriorly for nearly whole length; mid-femur with some golden scales on anterior surface forming an indistinct line on a dark ground, pale posteriorly on basal $\frac{1}{2}$; hind femur pale on both surfaces on basal $\frac{1}{2}$ except dorsally. *Abdomen*. dorsum black, with narrow basal silvery bands on II-VII, some orange scales on lateral margins of tergites, sternites with narrow white basal bands, narrow black medial bands, remainder covered with orange scales.

♂. —Palpi about $\frac{3}{4}$ length of proboscis, last two segments

† Theobald 1901 b, p. 48 (*Culex*), 1910 b, p 221; Edwards 1913 b, p. 228 (*Oncholorotatus*, syn.); Barraud 1924 a, p. 863.

with small hair-tufts. Ornamentation as in ♀. *Hypopygium* (fig 43, a): form of 9t, coxite, and harpago characteristic.

LARVA (fig. 42, c) — *Frontal hairs* A, B, C in a convex row towards front of clypeus, d slightly posterior and internal to C, A with 3-6 branches, B with 2-4, C with 3-6, d small, with 3 fine branches, e almost always single and long. Pre-clypeal spines fairly long and slender. Median hairs of mouth-brushes with moderately large teeth. *Antenna* comparatively short, 8-10 times length of greatest width, shaft with small spines, hair usually of 2 branches (sometimes 3 or 4), attached rather nearer apex than base. Mentum very similar to that of *oreophilus*, 12-15 teeth either side of median one. Lateral hairs of *thorax* well developed, arising from chitinated tubercles, stellate hairs of 3-5 short, strong branches on thorax and *abdomen*. Comb of 30 or more scale-like teeth, with fringed ends. Subsiphonal tuft of 8-12 branches. *Siphon* dark brown, $2\frac{1}{2}$ -3 times length of diameter at base, pecten usually of 20-22 teeth (variation 18-28), extending along more than basal $\frac{1}{2}$ of siphon, sometimes beyond base of hair-tuft, latter with about 4 branches. Pecten-teeth strong, with 4-6 basal denticles. Acus of moderate size. Saddle covering rather less than $\frac{1}{3}$ of *anal segment* in side view, posterior margin with small, strong spines, remainder of saddle, as well as siphon, covered with small ridges, many of which carry small points, *lh* long and single, about 3 times length of saddle; *usc* usually 4-branched, and 3-4 times length of saddle. Both pairs of anal papillae long and pointed, nearly twice length of saddle. Fan of 12-14 hairs, each dividing some distance from base into branches.

HABITAT.—Small pools and pot-holes in stream-beds, and tree-holes.

DISTRIBUTION—Common along the HIMALAYAS from Dungagali*, Hazara dist (*T. B. Fletcher*), Abbotabad*, KASHMIR (Gulmarg, etc, *Sinton*), to Sikkim border (Yatung*, Tibet, *F. M. Bailey*), and found in ASSAM HILLS (Cherrapunji, *Semor-White*). It is one of the commoner species breeding in both rock-pools and tree-holes in the Simla Hills.

Not recorded from outside the Indian area.

96 *Aedes (Finlaya) sintoni* Barraud, 1924


Ind Journ Med Res II, p 967 (♂ & ♀) TYPE-LOC.: Ferozepore Nullah, near Tangmarg, Kashmir, over 7,000', ix. 1923 (*Sinton*).
TYPE. ♂ & ♀ in Brit Mus

ADULT—Recognized by the largely dark thorax and presence of pale rings over some tarsal joints. Wing rather over 5 mm.

♀—*Head* a triangular area of creamy narrow scales on vertex, a border of similar scales to eyes, a black area on either

side towards front, a patch of broad white scales low down at each side, dark upright scales on nape. Palpi black, with a few pale scales at apices, about $\frac{1}{2}$ length of proboscis, latter entirely dark and noticeably long, longer than fore femur. *Thorax*: mesonotum mainly covered with brownish-black scales, creamy narrow scales on anterior margin arranged in three small patches, the lateral patches continued as a pale scaled line on each side to wing-root, a line of similar scales from front on each side curving over wing-root and continued

Fig 43



♂ hypopygial details of *Aedes*, subgenus *Finlaya*: a, *pulchreventer*, b, *sintoni*, c, *dissimilis*, d, *subsimilis*

to lateral lobe of scutellum; antescutellar bare space bordered with pale scales. Lanceolate white scales on mid-lobe of scutellum, narrower pale scales on lateral lobes. Broad pale scales on anterior and posterior margins of *ppn* and on *apn*. Patches of broad white flat scales on sternopleura and upper part of mesepimeron. *Wings*, dark scaled. *Legs*, black, a pale knee-spot on each femur, a few pale scales on anterior surface of fore and mid-pairs, chiefly along ventral border, posteriorly more extensively pale, hind

femur pale on both sides on basal $\frac{1}{2}$, the pale scaling continued in a line towards knee. Tibiæ black, except for small pale markings at base and apex, not forming definite rings. Tarsi with narrow pale rings at base and apex of segment 1 and base of 2, a few pale scales over joints between succeeding segments in some specimens. *Abdomen*. tergites black, with basal silvery bands, sternites with lateral basal and median apical patches of pale scales.

♂—Palpi a little less than $\frac{3}{4}$ length of proboscis, black, with indication of a pale ring on long segment, last two segments turned downwards and bearing moderate tufts of brown hairs. Other markings as in ♀ *Hypopygium*. in several respects resembles *A. (F.) pulchriventer*, but style and harpago (fig 43, b) of different form, and characteristic of this species.

LARVA—Unknown.

HABITAT—Rock-pools in stream-beds.

DISTRIBUTION—KASHMIR* (as above) W. HIMALAYAS. Murree*, 1922 (*Gill*).

97 *Aedes (Finlaya) auronitens* Edwards, 1922

Ind Journ Med Res x, p 268 (♂ & ♀) TYPE-LOC. Simla, viii-ix 1915 (*Christophers*). TYPE ♂ & ♀ in Brit. Mus.

ADULT ♀†.—Resembles *A. (F.) pulchriventer* in having orange patches on abdominal sternites, but differs as follows.—Dorsal surface of head covered with narrow yellow and upright scales without any dark areas, tibiæ pale at base, most pronounced on hind legs, fore tarsi with basal pale markings to first two or three segments, mid-tarsi with similar markings on first three segments, hind tarsi with narrow basal pale rings on segments 1-4, that on 1 being the widest.

♂—Palpi very slightly shorter than proboscis, slender, brownish-black, last two segments with few long hairs, not forming tufts. *Hypopygium*. resembles that of *A. (F.) christophersi*.

LARVA †.—Resembles the larvæ of *A. (F.) christophersi* and *greeni* very closely. Only three incomplete larval skins are available, one or more pairs of clypeal hairs being missing in each. The position of frontal hairs agrees with *christophersi*, and *A* and *O* both have a number of fairly long branches. Anal papillæ longer than in *greeni*, siphonal tuft of more branches (5-6) than in *christophersi*. Further material required before a more complete description can be given.

HABITAT—Tree-holes.

† Barraud 1924 a, p 864.

DISTRIBUTION —W. HIMATAYAS only. Simla *, 8,000', 1915 (*Christophers*), viii 1923, viii & ix. 1924 (*Barraud*), Krol Mt *, near Solan (Kalka-Simla road), 7,000', viii 1923, viii. & ix 1924, viii 1930 (*Barraud*), Muktesar *, Kumaon, 7,000', iv.-v. 1923 (*T. B. Fletcher*).

98 *Aedes (Finlaya) subsimilis* Barraud, 1927.

Ind Journ Med Res xiv, p 552 (♂) TYPE-LOC. Sukna, Darjeeling dist., 500', ix 1922 (*Barraud*) TYPE ♂ in Brit Mus

ADULT —Resembles *A. (F) dissimilis*, but differs in presence of white markings on palpi and tarsi, and in other details. Wing about 2.8 mm

♀ —Unknown.

♂ —*Head* vertex covered with flat black scales, a patch of silvery-white scales on each side, a very few dark upright scales on nape, orbital bristles black. Tori dark brown, some silvery scales in space between them. Antennal shaft and hairs brown. Clypeus dark. Palpi about length of proboscis, slender, without hair-tufts, mainly black scaled, but with a white ring nearer base than apex and some white scaling on upper and inner sides at base. Proboscis dark, about length of fore femur. *Thorax* mesonotum covered with dark brown scales, bristles brown. Scutellar scales brownish-black, flat on all lobes. Postnotum dark brown. Silvery-white flat scales on *apn*, also on upper and lower parts of sternopleura and upper part of mesepimeron, *ppn* without scales, 4-5 *ppn* bristles. *Wings* dark scaled. *Legs* brownish-black, femora yellowish ventrally, small white knee-spots dorsally on all femora. Tibiae dark, with very few pale scales at tips. Fore tarsi with small basal dorsal white markings on segments 1 and 2, not forming complete rings, similar markings on first three segments of mid-tarsi; hind tarsi with narrow basal white rings to segments 1-3, 4 entirely white except at tip, 5 dark. *Abdomen* tergites dark brown, faintly paler at base on I-V, lateral silvery patches commencing near base of each and not continued to the hind borders. *Hypopygium* (fig 43, d) coxite about $3\frac{1}{2}$ times length of greatest width in dorsal or ventral view. Style moderately long, widened towards tip, with short terminal appendage. Harpago represented by a single spine, somewhat similar to that of *A. (F) similis*.

LARVA. —Unknown.

HABITAT. —Hollow bamboos.

DISTRIBUTION —Known only from the type-locality

99 *Aedes* (Finlaya) *dissimilis* (Leicester), 1908

Cul Malaya, p. 91 (*Stegomyia*) (♂ & ♀) TYPE-LOC. Ampang, Malay Penin (Leicester) TYPE ♂ & ♀ in Brit Mus

var *karwari* Barraud, 1924, p. 865 (♀) TYPE-LOC Karwar, N. Kanara, ix 1921 (Barraud) TYPE ♀ in Brit Mus.

ADULT * (Pl III, fig. 16, and Pl IV, fig. 6) —Recognized by dark tarsi and silvery flat scales on *ppn*. Wing about 3 mm

♀—*Head*. dorsal surface mainly covered with flat black scales, a triangular area of flat white scales in middle in front, continued as a narrow border to eye-margins, a few yellow upright scales on nape. Antennæ, clypeus, palpi, and proboscis black, the last slender, and longer than fore femur, palpi about $\frac{1}{3}$ length of proboscis. *Thorax*. mesonotum almost entirely covered with black scales, usually a patch of golden scales a little distance in front of wing-roots and towards middle of mesonotum, more numerous in some specimens than in others, sometimes also a more or less pronounced median golden line. A patch of flat silvery scales laterally in front of each wing-root, similar scales completely covering *ppn* and on *apn* and larger part of pleuræ. *Wings*. black scaled, membrane with pronounced green, blue, and purple reflections. *Legs* fore legs almost entirely black, mid-femur with a silvery streak in middle on anterior surface, and another ventrally at knee both in front and behind, hind femur silvery-white, with a dark dorsal mark on apical $\frac{1}{3}$, not usually forming a complete ring. Tibiæ and tarsi black. *Abdomen* black, with large silvery basal patches; sternites with silvery basal bands.

♂—Palpi a little shorter than proboscis, black, with small terminal hair-tufts. Mesonotum in some specimens entirely covered with golden scales, in others the sides of mesonotum are dark. *Hypopygium* (fig. 43, c): differs from other species in the subgenus (except *subsimilis*) in shape of harpago, which is straight and resembles a bristle. Style short, with a terminal appendage about $\frac{1}{2}$ its length.

var *karwari*, ♀—Differs from type-form in having mesonotum almost entirely covered with golden scales, as in ♂.

LARVA.—*Frontal hairs* A, B, C, d all posterior to the level of bases of antennæ, A with 7–8 branches, B and C single and long, standing one behind the other, d small, with 4–6 branches, placed internal to B; e usually with 2 fine branches. Pre-clypeal spines slender. *Antenna* with long smooth shaft, about 14 times length of greatest width, and a single hair at about middle. Hairs of mouth-brushes simple. Mentum with usually 14 close-set teeth on either side of a large central one. Lateral hairs of *thorax* and

* Barraud 1924 a, p. 864

abdomen moderately developed, no stellate hairs or special structures on dorsum. Comb of 60-70 close-set teeth in a triangular patch, each tooth ending in a number of fine points. *Siphon* dark brown, about 3 times length of diameter at base. *Acus* small. *Pecten* of 22-25 sharp teeth, each with 2-3 lateral denticles at base. Hair-tuft of about 4 fine branches, at about middle of tube, its base just beyond most distal pecten-tooth, some long fine hairs on valves. Saddle covering rather more than $\frac{1}{2}$ *anal segment*, without spines on border, *lh* of 4-5 fine branches, *wsc* of 5-7 rather fine fairly long branches. Papillae long and pointed, dorsal pair about length of *wsc*. Fan of about 10 hairs, several of which are attached to membrane between fan-plate and base of segment.

HABITAT —Tree-holes

DISTRIBUTION —A fairly common forest species in S.W. and N.E. INDIA, where there is a heavy rainfall. No specimens have been seen from Burma, but it no doubt occurs there.

Known also from MALAY PENINSULA

100 *Aedes* (Finlaya) *albolateralis* (Theobald), 1908

Rec. Ind. Mus. II, p. 289 (*Stegomyia*) (♀). TYPE-LOC Syllhet, Assam, ix 1905 (Hall), and Lungleh, Lushai Hills, Assam, vii 1904. TYPE: ♀♀ in Ind. Mus.

Until recently this species has been considered synonymous with *A. (F.) niveus* (Ludl.), but Brug (1931) found that several forms or species have been confused under the last name. One of these is almost certainly *A. (F.) albolateralis*, described by Theobald from Assam. Unfortunately, so far as is known at present, this, and other forms, can only be distinguished from one another with certainty on details of structure of the ♂ hypopygia, or on the morphology of the larva. From examination of Indian specimens it appears that the commonest and most widely distributed is the one now considered to be *albolateralis*. The type-form of *A. (F.) niveus*, as interpreted by Brug, has been found in the Andamans and South Bengal, and there are three others, differing in ♂ hypopygial structure, one of which appears to be the form described by Brug as var. *B*. This form is known also from Indo-China, Sumatra, and Java. These three forms are here treated as distinct species, although it is possible that one or more of them may eventually be shown to be varieties of *A. (F.) niveus*.*

The following description has been made from specimens of *A. (F.) albolateralis* from Kasauli, Western Himalayas, where other forms resembling this species do not appear to occur.

ADULT †.—Resembles other species of the group in having flat scales on head and scutellum, large silvery patch on front

* [In addition to the forms here discussed, *A. (F.) pseudoniveus* Theo (= *subniveus* Edw.), of Borneo, Singapore, and Sumatra, belongs to this group. It differs from the five Indian forms in the presence of a silvery spot in middle of anterior surface of middle femora. ♂ and larva unknown.]

† Edwards 1921 a, p. 318 (*niveus*), Barraud 1923 g, p. 480 (*nivea*); Brug 1931 b, p. 24 (*niveus*, var. *A*).

of mesonotum (often divided in ♀), and dark legs, with basal $\frac{3}{4}$ of hind femur white. Wing about 3 mm.


♀.—*Head* a narrow border of silvery scales to eye-margins, rather wider in middle on front of vertex, remainder covered with flat black scales, faintly paler towards sides in some specimens, a few dark upright scales on nape. Antennæ, clypeus, palpi, and proboscis black, proboscis about length of fore femur, palpi short, only about $\frac{1}{4}$ or $\frac{1}{2}$ length of proboscis. *Thorax*: mesonotum with a large silvery-white area on either side in front, usually some silvery scales on front margin in middle, or more extensive silvery-white scaling tending to unite the lateral patches in one large area, remainder brownish-black, but usually white scales on either side of the antescutellar bare space. Scutellar scales flat on all lobes, white, brownish, or dark brown, variable in proportions. Integument of pleuræ almost black, patches of silvery flat scales, one covering larger part of mesepimeron, another large area in front of this, and another below, also smaller patches on coxæ, propleura, and *apn*, *ppn* bare. *Wings* dark scaled. *Legs*: fore femur black anteriorly, a pale streak on basal $\frac{1}{2}$ beneath; mid-femur similar, but more extensively pale posteriorly, and usually with some silvery scaling on ventral border on anterior surface, hind femur entirely white on basal $\frac{3}{4}$ or more. Tibiæ and tarsi entirely brownish-black. *Abdomen* black, with large white lateral, basal, wedge-shaped markings, not produced far on to dorsum on II–IV, but forming narrow basal bands on V–VII. Sternites mainly dark, V–VII usually somewhat projecting, VIII fairly large, without scales, cerci scarcely visible.

♂.—*Head* scales mostly creamy or light brown, but appearing darker in some lights, scales at sides lighter, in some specimens the dorsal surface of head appears much lighter than in others. Palpi black, a little shorter than proboscis, terminal segments with a moderate number of long hairs. Antennal plumes fairly dense, brown. *Thorax*: white patch on mesonotum usually complete and covering anterior $\frac{3}{4}$, but few or no white scales above wing-roots, and not many bordering bare antescutellar space. Other markings very similar to those of ♀. *Hypopygium* (fig 44, g, h, l). 9t narrow, with a pair of small submedian lobes close together, each with 2–4 strong flattened bristles. Coxite with a group of very large scales on ventral border near apex; dorsal border with a row of hairs, and a collection of long lanceolate scales at base (this last character distinguishing this species from allied forms). Style with terminal appendage nearly $\frac{1}{2}$ its length.

LARVA (fig 45, c–f)—*Frontal hairs* A, B, and d in a transverse row slightly posterior to level of bases of

antennæ, *C* almost directly behind *B*, *A* with 6-8 branches, *B* with 4-10, more usually 7, *C* with 5-11, *d* with 7-10, this hair being comparatively well developed, *e* single or split into 2. Preclypeal spines long and slender. Mouth-brush hairs simple. Antenna rather long and stout, some spines on curved shaft, hair-tuft at about middle, with 6-7 branches. Central tooth of mentum large, with 9-10 teeth on either side. Lateral hairs of thorax moderately

Fig 44



♂ hypopygial details (base of coxite from inside, middle of 9th tergite) of *Aedes* (Finlaya) *niveus* and related species: a, b, *albomiveus*; c, d, *novoniveus*; e, f, *niveoides*; g, h, i, l, *albolateralis*; j, k, *niveus* (h, whole coxite, showing scales; l, harpago.)

developed, pleural hairs on meso- and metathorax arise from tubercles furnished with a curved spine and a few smaller straight spines. A number of hairs with few short, fine branches on thorax and abdomen, not developed as obvious stellate hairs, as in *A. (F.) suffusus*. Larger lateral hairs on abdomen 2-3-branched. Comb of 8-12 large pointed teeth in a single row. Subsiphonal tuft with 6-8 branches.

Siphon slightly wider in middle than at either end, $3-3\frac{1}{2}$ times length of diameter at base. *Acus* very small and apparently detached, and lying separately in some specimens. *Pecten* of 15-24 rather small teeth, with one large and one or two small lateral denticles; *pecten* confined to basal $\frac{1}{2}$ of tube. *Hair-tuft* at about middle of tube, with 4-6 branches. *Saddle* almost enclosing *anal segment*, some spines on posterior margin towards dorsum; *lh* usually of 2 long branches, occasionally single, more rarely of 3-6 branches, *usc* of 3-4 branches. *Fan* of about 8 hairs, two or three of which arise from membrane towards base of segment, each hair divided into 2-4 branches, *fan-plate* small. *Dorsal papillæ* about twice length of ventral pair and about length of longer *fan-hairs*.

HABITAT.—Tree-holes and bamboo-stumps.

DISTRIBUTION *.—**ASSAM**: Golaghat, Sibsagar dist, v 1925 (*Barraud*), Hafiong, Cachar Hills, viii 1922 (*Barraud*); Nongpoh, Khasi Hills dist, vii 1922 (*Barraud*). **E BENGAL**: Rangamati, Chittagong Hill Tracts, ix 1922 (*Barraud*). **N BENGAL**: Sukna, Darjeeling dist, ix 1922 (*Barraud*), Kurseong, Darjeeling dist, viii. 1928 (*Sobha Ram*), Meen-glas, near Jalpaiguri, Duars, v 1923 (*M O T Iyengar*). **W. HIMALAYAS**, 4-8,000' Simla, ix 1924 (*Barraud*), Solan, viii 1923 (*Barraud*), Krol Mt., near Solan, vii 1930 (*Barraud*), Kasauli, vii-viii 1924 (*Barraud*), Koti, near Kalka, viii 1923 (*Barraud*). **N KANARA**: Karwar, ix. 1921 (*Barraud*).

101 *Aedes (Finlaya) niveus* (Ludlow), 1903 †.

Journ NY Ent Soc xi, p 139 b (*Stegomyia*) (♀) TYPE-LOC: Oras, Samar, Philippine Is. TYPE ♀ in Brit Mus; other ♀♀ (paratypes or cotypes) in US Nat. Mus.

Stegomyia pseudonivea Theobald, 1910 (♂ only), MC v, p 176. TYPE-LOC. Andaman Is (*Lewis*). ALLOTYPE ♂ in Brit Mus [Type ♀ from Singapore is a different species].

ADULT ♀—Appears to differ slightly from *albolateralis* in having a rather larger white area on front of mesonotum, this white area solid, as in the ♂, its posterior border not or scarcely emarginate. Scutellar scales all black.

♂—[Scales on dorsum of head black, with a narrow silvery margin to eyes (in all 6 ♂♂ examined). Scutellar scales black. Scales on posterior $\frac{1}{2}$ of mesonotum mostly black, no


* As checked by Author by examination of ♂♂

† As restricted by Brug 1931 b, p. 25, it is not known with certainty whether this was the form described by Ludlow. See also Edwards 1913 b, p. 227, and 1921 c, p. 318, Borel, 1928 b, pp 43-46

white ones over wing-base and few or none surrounding antescutellar bare space Abdominal tergites II-IV unbanded] *Hypopygium* (fig 44, j, k): ooxite without a group of scales at base of dorsal border, but with some long hairs in this position 9t with a pair of prominent submedian processes (close together), each with 4-5 strong hairs Style with appendage rather more than $\frac{1}{2}$ its length

LARVA—Thus has not been isolated in India Dr Brug has kindly furnished me with a translation from the German

Fig 45



Larval structures of *Aedes*, subgenus *Finlaya* a, b, *drassimilis* (head and siphon), c-f, *albolateralis* (head, siphon, comb, and pecten-teeth), g, h, *novoniveus* (pecten and comb-teeth enlarged and whole pecten); i, j, *alboniveus* (ditto).

of his remarks on this larva *, as found by him in the Dutch East Indies —“ Under the name *F. nivea*, Borel (1928) has described the larva of var. *B*. That of the type is distinguished from the latter in the following respects. —Scales of the comb on the 8th segment more in number and more hairy, the number of scales being 16-19, as against 8-10 in the var *B*

* Brug 1931 b, p. 25.

larva; the siphon is much shorter (index $2\frac{1}{2}$, in var. *B* larva 5), the teeth of the pecten are sharper, and, but for the tips, darker than in var. *B*."

HABITAT.—Tree-holes and bamboo-stumps

DISTRIBUTION.—SOUTH BENGAL, Cassiabad *, Sundarbans (*M. O. T. Iyengar*) ANDAMAN ISLANDS *, vii. 1926 (*Covell*)

Brug has recorded the type-form from SUMATRA, BALI, SINGAPORE, ANDAMAN IS., ? PHILIPPINE IS., and JAPAN.

102. *Aedes* (*Finlaya*) *alboniveus*, sp. n

TYPE-LOC Kuraseong, Darjeeling dist., E Himalayas, ix 1922 (*Barraud*) TYPE ♂ in Brit. Mus

ADULT ♀.—Not distinguished from *A. (F.) niveus* White patch on mesonotum complete, as in ♂

♂.—[Scales of head creamy. Most of the scales above wing-base, as well as a large number round antescutellar bare space, white. Scutellar scales creamy or pale brown. Abdominal tergites II–IV without complete basal pale bands.] *Hypopygium* (fig. 44, a, b). ♂t with a pair of small submedian lobes, much smaller than in the type-form, and rather more separated than in *albolateralis*, each lobe with 1–3 hairs. Some long hairs at base of coxite, and a ridge carrying an elbowed process and a stout hair (this character distinguishes this form from all others). Appendage of style long, about $\frac{2}{3}$ length of style.

LARVA (fig. 45, 1, 1).—One incomplete skin, from which a ♀ resulted, shows the following characters:—Antenna long and tapering, shaft with very strong spicules, tuft well before middle. Comb of about 12 long teeth, with lateral fringes for about $\frac{1}{2}$ the length (in *niveus* the teeth are fringed from near base to tip, the number of teeth being usually 16–19, in *niveoides* they are delicately fringed at base only, the number being usually 8–10). Siphon about 1 mm. long, not including valves (in *albolateralis* it is usually about 0.7–0.8 mm.). Pecten of 21 teeth, individual teeth about $\frac{2}{3}$ length of comb-teeth, basal teeth small; each tooth with one lateral denticle about half-way along its length. Thorax and abdomen with a number of hairs with short branches of stellate form, and long lateral subplumose hairs.

Further material is required before a complete description can be given.

HABITAT.—Tree-holes and bamboos.

DISTRIBUTION.—Known only from the type-locality.

* ♂♂ examined by author.

103 *Aedes* (Finlaya) *niveoides*, sp. n.†

TYPE-LOC. Nagargah, Bombay Deccan, viii. 1921 (Barraud)

TYPE. ♂ in Brit. Mus.

ADULT —Does not show any external difference from *A. (F.) niveus*♂—*Hypopygium* (fig. 44, a, f). 9t with a pair of submedian, rather flat lobes, each bearing only 2-3 hairs. A small boss-like lobe at base of coxite from which a collection of very long hairs arises, also a ridge carrying 6-7 strong hairs, but no processes as in *alboniveus*. Style with terminal appendage approximately $\frac{2}{3}$ its length.LARVA —Has not been isolated in India. Borel has described a larva from Cochim China which Brug has determined as that of his var. *B*. This var. appears to me to be probably identical with the form here described as *niveoides*. Some notes on the larva are given above under *A. (F.) niveus*.

HABITAT —Bamboos

DISTRIBUTION —INDIA. Nagargah (type-locality as above), one ♂ which corresponds closely in hypopygial structure, from Meenglas*, Jalpaiguri, Bengal Duars, ix. 1926 (*B. M. Khan*, per *M. O. T. Iyengar*).Brug records his var. *B* from INDO-CHINA, SUMATRA, and JAVA.104. *Aedes* (Finlaya) *novoniveus*, sp. n.

TYPE-LOC. Mungpoo, Darjeeling dist., E. Himalayas, x. 1922 (Barraud). TYPE. ♂ in Brit. Mus.

ADULT ♀ —Not distinguished from *A. (F.) niveus*. White area on mesonotum complete, as in ♂.♂ —[Scales of head and scutellum all creamy. Many white scales above wing-base and around bare antescutellar space. Complete basal white bands on tergites II-IV evident except in shrunken specimens †.] *Hypopygium* (fig. 44, c, d). 9t with moderate-sized, rounded, submedian lobes, each carrying 4-5 hairs. Coxite without a conspicuous collection of long hairs or scales at base, but a ridge carrying some fairly fine hairs, a few of these rather long. Harpago expanded fairly near the base, not gradually widening to the middle, as in other forms. Style with terminal appendage slightly more than $\frac{1}{2}$ its length.† Apparently *A. (F.) niveus* var. *B* of Brug (1931 b, p. 25). See also Borel 1928, pp. 43-45.‡ [2 ♂♂ in British Museum from Malaya (*Leicester & Stanton*) agree with the type ♂ in these respects, as well as in hypopygium.]

LARVA (fig. 45, g, h).—One isolated larval skin, from which a ♀ resulted, is thought to be most probably of this form, as it was obtained from Mungpoo, Darjeeling dist., at the same time as the type ♂. It differs from all other described larvæ of this group in having the 4-5 pecten-teeth nearest base of siphon without lateral denticles, and larger than the more distal teeth, which have denticles. Antennæ shorter than in *albolateralis* or *alboniveus*, spicules on shaft inconspicuous. In other respects the larva resembles *albolateralis* fairly closely. Comb of 8 large sharp teeth in a single row, *etc.* of 4 fairly long branches †.

HABITAT—Bamboo-stumps and tree-holes.

DISTRIBUTION—E HIMALAYAS. Mungpoo* (type-locality, as above), Sureil*, near Mungpoo, Darjeeling dist. x. 1922 (*Barraud*), Kurseong*, Darjeeling dist., ix. 1922 (*Barraud*) ASSAM. Nongpoh*, Khasi Hills dist., vii. 1922 (*Barraud*).

Subgenus **CHRISTOPHERSIOMYIA** Barraud, 1923.

Ind. Journ. Med. Res. x, p. 786. Genotype, *Stegomyia thomsoni* Theo.

ADULT ‡.—Black and white ornate mosquitoes, resembling *Stegomyia*, but having a pale band on proboscis and palpi about $\frac{1}{4}$ length of proboscis in both sexes. Scales on vertex of head and on scutellum all flat, but those on *apn* narrow. Plumose hairs on antennæ of ♂ projecting mainly dorsally and ventrally, not in regular whorls. Phallosome of ♂ hypopygium scoop-shaped, not divided into lateral plates, and without teeth. Coxite with small lobe on inner surface. Style simple, slightly tapering, with terminal appendage. Paraproct without lateral arm, no teeth or small hairs at crown.

LARVA (4th stage).—Very similar to *Stegomyia*, but antenna longer and with spicular shaft, and with branched hair at about middle.

DISTRIBUTION and BIONOMICS—The three known species are confined to India and Ceylon, only one of these being at all common. The larvæ are found in tree-holes during the monsoon periods.

† [In this specimen the antennal hair-tuft is about at middle of shaft and the pecten has about 12 teeth. A drawing made by Mr. A. J. E. Terzi of a larval skin (from which a ♀ hatched), obtained in Malaya by Dr. A. T. Stanton in 1915, shows large, simple, basal pecten-teeth as in the above specimen, but the antennal tuft is far *beyond* the middle and only 8 pecten-teeth are present.]


* ♂♂ examined by author.

‡ Edwards 1932, p. 159

Key to Adults.

- 1 Hind tibia with a white ring on basal $\frac{1}{4}$. . . *thomsoni*, p 213
 Hind tibia without a white ring on basal $\frac{1}{4}$ 2
 2 Fore femur dark, except for a white mark
 near base on posterior surface (beneath) . . . *annulirostris*, p. 215
 Fore femur with a white mark on apical
 $\frac{1}{4}$ of anterior surface, connected with a
 white streak on posterior surface . . . *abis*, p 215

Fig 46



♂ hypopygial structure of *Aedes*, subgenus *Christophersiomyia*.
 a, *thomsoni*, b, *annulirostris*.

105. *Aedes* (*Christophersiomyia*) *thomsoni* (Theobald), 1905 *

Gen. Ins., Fam. Culicidae, p 18 (*Stegomyia*) (♀) TYPE-LOC :
 N.W. Provinces, India (*Wynville-Thomson*) TYPE : ♀ in Brit.
 Mus.


ADULT (Pl VI, figs. 12, 16, and Pl. VIII, fig 1) —Wing 2.5–3.5 mm Distinguished from the two other species on characters given in key Mesonotum ornamented with silvery-white and pale brown scaling, and with a pair of brownish eye-like spots in front. Amount of pale brown scaling on posterior part of mesonotum variable All lobes of scutellum with flat white scales. Palpi about $\frac{1}{4}$ length of proboscis in both sexes, latter with a fairly wide white ring.

* Theobald 1907, p 174; Barraud 1923 a, p. 787.

♂.—*Hypopygium* (fig 46, a) see remarks under subgenus above.

LARVA * (fig. 47) —Body creamy or white, contrasting with dark head and siphon. *Frontal hairs* A, B, and C all fairly long and branched. A with about 10 branches, B usually with 2, one of these thicker and longer than the other, C with about 8. d small, with few fine branches, and internal to other three pairs. All four pairs are a little posterior to level of bases of antennæ. Preclypeal spines long, slender, tapering to a fine point. Hairs of mouth-

Fig 47



Larval structures of *Aedes* (*Christophersomyia*) *thomsoni*: a, tail-end, b, base of metathoracic pleural hairs, c, head: d, mentum; e, f, pecten and comb-breath

brushes simple. Antenna fairly long, shaft with small spicules; tuft of 2-3 hairs at about middle. Mentum rather pointed, triangular, about 11 teeth either side of central one, basal teeth large, four or five nearest apex much smaller. Hairs on *thorax* and *abdomen* moderately developed, no special structures present. Comb of about 14 strong, pointed teeth in one close-set rank; teeth with delicate fringe from base, almost to tip. *Siphon* $3\frac{1}{2}$ -4 times length of diameter at base, widest in middle. Aous very small or absent. Pecten.

* Barraud 1923 h, p. 503

of about 16 small teeth, each with 2-3 basal lateral denticles; hair-tuft of 6-8 branches, at about middle of tube *Anal segment* almost enclosed in chitination, some very small spines scattered over surface towards dorsum, *osc* of 2 branches, *osc* single and long, *lh* of 4-5 fine branches. Dorsal papillæ about 3 times length of segment, ventral pair shorter. Fan small, longest hairs only about $\frac{1}{2}$ length of dorsal papillæ.

HABITAT —Tree-holes

DISTRIBUTION —A common monsoon species occurring from N W FRONTIER as far east as BIHAR (Pusa) and southwards to MADRAS. So far as is known it is absent from Assam, Burma, and Ceylon, and it has not been recorded from beyond limits of Indian region.

106 *Aedes (Christophersiomyia) annulirostris* (Theobald), 1905.

Journ Bomb Nat Hist Soc xvi, p. 239 (*Stegomyia*) (♀) TYPE-LOC Peradeniya, Ceylon, 1 1902 (*Green*) TYPE ♀ in Brit Mus

ADULT † (Pl VI, fig. 11, and Pl VIII, fig. 1) —Very similar *A (C) thomsoni* and *ibis*, but differs as indicated in key.

♂ —*Hypopygium* (fig. 46, b) does not show any marked difference from that of *thomsoni*.

LARVA —Known only from one incomplete larval skin, very similar to *thomsoni*, but further material is required before a description can be given.

HABITAT —Tree-holes and water-butts.

DISTRIBUTION —BOMBAY DECCAN Belgaum* and Nagar-gali*, viii 1921 (*Barraud*) N KANARA Karwar*, bred ii. 1931 (from tree-hole material supplied by *T R Bell*) CEYLON type-locality, as above, Labugama, 1 1933 (*Henry*).

107. *Aedes (Christophersiomyia) ibis* Barraud, 1931.

Ind Journ Med Res xix, p. 610 (♀) TYPE-LOC Sukma, Dar-jeeling dist., 500', viii. 1928 (*Sobha Ram*) TYPE ♀ in Brit Mus

ADULT ♀ —In general appearance and character of scaling resembles *A (C) annulirostris*, but differs as indicated in key, also in having no lateral dark areas on mesonotum in front of wing-roots.

♂ and LARVA. —Unknown

DISTRIBUTION —Known only from type-locality

† Theobald 1907, p. 173, Barraud 1923 a, p. 788.

Subgenus **RHINOSKUSEA** Edwards, 1929*

Bull. Ent. Res. xx, p. 342. Genotype, *Ficalbia longirostris* Loic
Ficalbia Leicester, 1908 (nec Theobald), Cul. Malaya, p. 228

The subgenus includes only one species, the characters of which are given below. The larvæ live in crab-holes and pools in mangrove-swamps. Nothing is known of the habits of the adult.

108. *Aedes* (*Rhinoskusea*) *longirostris* (Leicester), 1908

Cul. Malaya, p. 228 (*Ficalbia*) (♂ & ♀) TYPE-LOC. Kuala Klang, Malay Penin (Leicester) TYPE ♂ & ♀ in Brit. Mus.

ADULT †—A small, slender, brownish-black mosquito, without any ornamentation. Wing 2-2.7 mm.

♀—*Head* mainly flat-scaled. Palpi quite short, projecting only slightly in front of clypeus. Proboscis long and slender, as long as whole body. Scutellum flat-scaled. All tarsal claws simple. Legs unusually long; segment 1 of hind tarsi slightly longer than tibia (a character distinguishing this mosquito from all others of the genus *Aedes*). Cerci rather long.

♂—Resembles the ♀ in all details except structure of hypopygium and antennæ, latter only moderately plumose, hairs evenly spread round shaft. *Hypopygium* (fig. 48): remarkably distinct from that of any other species. Coxite rather narrow, with a hairy lobe near apex, and two large processes arising from inner surface. Style fairly long, hairy on apical $\frac{1}{3}$, and with a small double appendage. Paraprocts with about 5 small hairs near crown. Phallosome simple, not toothed nor divided into lateral plates.

LARVA.—Described by Edwards (1926 a, p. 120) from Singapore specimens as follows:—

“Head very broad and rather light in colour. Antennæ long, curved, and tapering, surface with numerous spicules, especially towards the base, tuft of 3 or 4 long hairs placed a little before the middle; a long fine hair near the tip. Frontal hairs placed close together and far back, the arrangement being practically the same as in *Aedes cinereus* Mg., the three main pairs of tufts very long, consisting respectively of 10-12, 2, and 6-7 hairs. Mentum with about 12-14 teeth on each side of the stronger median tooth. Abdominal hairs inconspicuous. Comb consisting of about 40 small narrow bare teeth in a patch. Anal segment with small saddle which is covered with small spicules; lateral hair single and rather long; about 8 tufts in the brush,

* Edwards 1932, p. 160.

† Edwards 1917, p. 224, and 1924, p. 391; Barraud 1928 b, p. 563.


none before it Siphon not very dark, index about 3.5, well-developed acus at base, pecten just beyond the middle, with about 12 spines, none detached, tuft of 3 very long hairs, placed far beyond the pecten and only a little before the tip of the siphon. Valves normal "

HABITAT.—Pools in mangrove-swamps; crab-holes.

DISTRIBUTION.—ANDAMAN Is. *, caught in crab-hole, Port Monat, ix. 1911 (*Christophers*), vii. 1926 (*Sobha Ram*).

Known also from MALAY PENINSULA and ARCHIPELAGO and NORTH AUSTRALIA.

Fig. 48.



♂ hypopygium of *Aedes (Rhinoskusea) longirostris* Leac, ventral and lateral view. Lettering as on p 4.

Subgenus *STEGOMYIA* Theobald, 1901.

M.C. i, p. 283. Genotype, *Oulex fasciatus* Fab.

Scutomyia Theobald, 1904, Entom xxxvii, p. 77. Genotype, *S. albolineata* Theo.

Quasistegomyia Theobald, 1906, 2nd Rept Wello Lab. p. 69. Genotype, *Q. unilineata* Theo.


Catantassomyia Dyar & Shannon, 1925, Insec Ins. Mens. xii, p. 71. Genotype, *C. meronephada* D. & S.

ADULT †.—Small or medium-sized black or dark mosquitoes, highly ornamented with patches, spots, or lines of snow-white scales. Two or more basal white bands on tarsi of at least one pair of legs, or with one or more tarsal segments completely white (*vide* Pls. VII & VIII). Tarsi in no case all completely dark nor with apical and basal banding

† Edwards 1932, p. 160.

Proboscis entirely dark (except in the aberrant species *A (S) vittatus*, in which there is some yellow scaling on that organ). Scales on vertex and scutellum broad and flat, a few upright scales on nape (more numerous in *A (S) vittatus*, where they extend forwards for some distance). Palpi of ♂ rather slender, terminal segments upturned and with only few hairs. Antennæ of ♂ with plume-hairs mainly projecting on two sides, not in regular whorls. ♂ *hypopygium* coxite with plaque or lobe on inner surface bearing hairs

Fig 40



♂ hypopygial structures of *Aedes (Siegomys) w-albus* Theob. a, hypopygium, from above, 9th tergite removed, b, coxite from beneath; c, lateral plate of phallosome from above and from side; d, 9th tergite, e, paraproct, from above and from side. Lettering as on p. 4

or processes of diagnostic importance, form of 9th tergite also characteristic in many species, style usually, though not always, rather long, with straight tube-like appendage near tip. Harpago absent. Paraprocts without teeth or small hairs at crown, and, except in *A. (S.) aegypti* and *desmotes*, without a ventral or basal arm. Phallosome divided into small lateral plates with numerous teeth. ♀ *hypopygium*: cerci quite short, segment VIII partially retractile, sternite of only moderate size. The above-mentioned characters

distinguish *Stegomyia* from certain species of the subgenus *Finlaya* and genus *Armigeres*, which have a rather similar black-and-white ornamentation. In the species last referred to the tarsi are either entirely dark or with apical and basal banding, or with only one basal band. In some of these species, also, there is a pale band or extensive pale scaling on the proboscis.

LARVA (4th stage).—Antennal shaft tooth, with usually a single hair in place of tuft. Frontal hairs *B* and *C* usually one in front of the other. Thorax without large spines on dorsum, but stellate hairs often present, as well as on abdomen. Comb-teeth in a single row, in some species few and arising from posterior margin of chitinated plate. Siphon 1-3 times length of diameter at base. No acus. Pecten well developed, teeth in regular rank, and all about same size, no detached teeth (except in *A. (S.) vittatus*, in which there is a detached tooth beyond the hair-tuft), siphon-tube with one pair of hair-tufts at about middle, or between this and apex. Chitinated saddle of anal segment usually rather small, anal fan often small also.

DISTRIBUTION and BIONOMICS.—The subgenus is represented in all parts of India, some species being very common. Many are day-flying insects, and readily attack man. *A. (S.) aegypti* (*Stegomyia fasciata*) is the well-known yellow-fever mosquito, and the Indian race has been proved capable of transmitting this disease. It is considered to be the carrier of dengue fever. It is almost entirely a house-frequenting species, breeding in artificial collections of water. The literature dealing with the bionomics of this and other species of the subgenus, in connection with disease, is extensive, and references will be found in the 'Tropical Diseases Bulletin'. Other species breed extensively in tree-holes and bamboo-stumps, some occurring up to 8,000' or more in the Himalayas. The eggs are laid singly and are capable of resisting desiccation over long periods.

About 44 species are known, of which 16 have been found in India, the remainder occurring either in Africa or the eastern part of the Oriental region, some extending to Australia and the Pacific islands. *A. (S.) aegypti* is found throughout the tropics and subtropics of the world.

Key to Adults

- | | |
|---|--------------------------|
| 1. All tibiae with white rings | 2 |
| Tibiae without white rings | 3 |
| 2. Mesonotum marked with 4-8 small white spots; femora with preapical white rings, proboscis with scattered yellow scaling. . . | <i>vittatus</i> , p 245. |
| Mesonotum otherwise marked; femora without preapical white rings, proboscis entirely dark | <i>desmotes</i> , p 225. |

- 3 Mesonotum marked with a pair of lateral curved white lines, and usually also with a pair of submedian yellowish lines, two dots of white scales on clypeus. *egypti*, p. 221
Mesonotum otherwise marked, clypeus without white scales 4
- 4 Palpi entirely dark; proboscis thin and longer than fore femur, mesonotum with a fairly narrow, silvery, median stripe in front, last two hind tarsal segments all dark *albolineatus*, p. 243.
Palpi with white scaling, proboscis normally thick and about length of fore femur 5
- 5 Mesonotum with a narrow median white line running nearly the whole length 6
Mesonotum with a broad white anterior stripe, or with a white patch or pair of patches in front 10
- 6 Mid-femur with a preapical white spot on anterior surface. *unilineatus*, p. 242.
Mid-femur without a white spot 7
- 7 White scales on pleurae arranged more or less in two lines; a line of white flat scales over wing-root continued nearly to lateral lobe of scutellum *scutellaris*, p. 240.
White scales on pleurae arranged in irregular patches, a patch of white scales in front of wing-root only 8
- 8 Pale markings on mesonotum, except for the median white stripe, of a yellowish colour *flavopictus*, p. 230.
Pale markings on mesonotum all silvery white 9
9. Abdomen without silvery bands on dorsum (? always) *subalbopictus*, ♂, [p. 238.
Abdomen with silvery basal bands on dorsum { *albopictus*, p. 233;
pseudalbopictus,
p. 235, *novalbopictus*, p. 237.
- 10 Mid-femur with a median white spot on anterior surface, mesonotum with several white patches *u-albus*, p. 232.
Mid-femur without a white spot 11.
- 11 Mesonotum with a broad median white stripe, narrowing posteriorly *medicopunctatus*, p. 230.
Mesonotum with a roundish white area in front 12.
12. Mesonotum with a small median white spot in front; all segments of hind tarsi white-ringed *edwardsi*, p. 232.
Mesonotum with a large median white patch in front, last one or two hind tarsal segments all dark *annandalei*, p. 227;
craggi, p. 229

109 *Aedes* (*Stegomyia*) *aegypti* (Linnæus), 1762

Hasselquist's Reise nach Palestina, p 470 (*Culex*) (♀ ?) TYPE-LOC Egypt TYPE non-existent

SYNONYMY (abridged) * —

Culex argenteus Poirët, 1787, Journ Phys xxx, p 245 (♀) TYPE-LOC Barbary TYPE non-existent

Culex fasciatus Fabricius, 1805, Syst Anthat p 36 (♂ ?) TYPE-LOC. West Indies ("Americas Insulis") TYPE non-existent (originally in Copenhagen Mus)

Culex calopus Meigen, 1818, Syst Besch 1, p 3 (♂♀) TYPE-LOC. Portugal TYPE non-existent (unless in Vienna Mus)

var *atrata* Edwards, 1920, Bull Ent Res x, p 129 (♂ & ♀). TYPE-LOC Accra, Gold Coast, W Africa, vi 1919 (*J W S Macfie*) TYPE ♂ & ♀ in Brit Mus

var *luciensis* Theobald, 1901, M C 1, p 297 (♂ & ♀) TYPE-LOC Georgetown, Demerara, South America, vi. 1899 (*Quelch*) TYPE ♂ & ♀ in Brit Mus

var *queenslandensis* Theobald, 1901, M C 1, p. 297 (♀) TYPE-LOC Burpengary, South Queensland, Australia, xi (*Bancroft*) TYPE ♀ in Brit Mus

ADULT—Recognized by characteristic ornamentation as shown in Pl VI, figs 1, 13, and Pl VII, fig 1 The following details will assist in identification in cases where the mesonotum has become denuded (as very frequently occurs)

♀—*Head*: silvery-white flat scales in middle of vertex, continued downwards between eyes, similar scales on tori, two small silvery-white dots on clypeus (white scales are not present in this position in any other species except *A. (S.) vittatus*); tips of palpi conspicuously white *Thorax* flat silvery-white scales on all lobes of scutellum. *Legs* mid-femur, when viewed from the front, with a white longitudinal line running from the base for nearly whole length, but not continued quite to knee, anterior surface of hind femur pale on basal $\frac{1}{2}$, or more, the pale scaling continued as a thin longitudinal line nearly to knee Tarsi of fore and mid-legs with narrow basal white bands to first two or three segments, those of hind legs with more conspicuous white basal rings on segments 1-4, that on 4 being usually the widest, segment 5 usually entirely white *Abdomen*: tergites brownish to black, with narrow dull white basal bands on II-VI, VII with two small silvery-white dots, lateral basal silvery-white patches on I-VII not usually visible in dorsal view

♂.—Ornamentation similar to that of ♀ Palpi (Pl VI, fig. 13) with two white rings on the long segment and with


* For further synonymy and refs., vide Edwards 1932, p. 162. As regards India, vide Barraud 1928 c, pp. 377-388. For account of work on this species in other parts of the world, vide reviews in Trop. Dis. Bull. and Rev. Appl. Ent., also Bonne-Wepster & Brug 1932, p 47.

white marks on undersides of last two segments at base *Hypopygium* (fig 51, a) differs from all other species in shape of 9t. The paraproct has a well-developed ventral arm, an unusual feature, absent in all other Oriental species except *A (S) desmotes*.

var *lucienensis* Theo —Differs from the type-form in having the apical part of segment 5 of hind tarsi black. Occasional specimens of this form are met with in India.

var *queenslandensis* Theo —In this form there are some dark purplish scales on mid-lobe of scutellum and, on the

Fig 50




♂ hypopygial structures of *Aedes (Stegomyia)*: a, *aegypti*; b, *mediotpunctatus*.

abdomen, both basal and apical bands of pale scales, and a broad, pale, median longitudinal line, or the whole dorsum may be almost entirely pale. Specimens resembling this var have been received from Bangalore, South India.

var. *atrifarsis* Edw.—Tarsi of fore and mid-legs almost entirely black, only 2–3 white scales at bases of first two segments, hind tarsi black, with very narrow white rings at bases of segments 1–3 and 5. White rings on ♂ palpi narrower than usual. No Indian specimens of this var. are known.

Other Variations—Connal (1927, p. 5) has described and figured some remarkable varieties of this species from Lagos, Nigeria, including the var *atrutarsis* referred to above. In some Indian specimens the scales on the middle of the mesonotum are much paler than usual, and the narrow submedian yellowish lines obliterated.

Fig 51




Larval structures of *Aedes (Stegomyia)*. a-l, *egypti*, g-k, *vittatus*.
b, l, comb-teeth, c, h, pecten-teeth, f, base of metathoracic
pleural hairs

LARVA (fig. 52, a-f)—Distinguished from those of other species by the form of the 8-12 comb-teeth, which have well-developed lateral denticles, the teeth do not arise from a chitinated plate, as is the case in *A. (S.) desmotes* and some other species. Antennal hair single and attached a little nearer apex than base of shaft. Lateral hairs of *thorax* moderately developed, thorn-like process at bases of pleural hairs on meso- and metathorax comparatively large and

ending in a single point *Siphon* usually rather more than twice length of diameter at base, tuft of 3-5 branches, a little beyond middle of tube from base Pecten of 12-20 teeth, with basal lateral denticles *Anal segment* almost enclosed in chitination, *th* single or 2-branched, *asc* of 2-4 branches *Papillæ* with rounded extremities and about

Fig 52.



Distribution in India of *Aedes (Stegomyia) aegypti*, as at present known

twice length of segment. Longest hairs of fan about twice length of papillæ.

HABITAT.—Artificial collections of water near, or within, human habitations, such as chattis, water-butts, anti-formicas, disused kitchen utensils, etc. It is unusual to find larvæ in tree-holes or bamboo-stumps, or in natural ponds or wells

DISTRIBUTION.—Generally distributed throughout the Indian region. The names of places where the species has been found are shown on the accompanying map

110 *Aêdes* (*Stegomyia*) *desmotes* Giles, 1904.

Journ Trop Med vii, p 387 (♀) TYPE-LOC · Philippine Is
TYPE ♀ in Brit Mus

Stegomyia gracilis Leicester, 1908, Cul Malaya, p 81 (♂ & ♀).
TYPE-LOC : Malay Penin (*Leicester*). TYPE ♂ & ♀ in Brit
Mus

Stegomyia albipes Theobald, 1910, Rec. Ind Mus iv, p 11 (♀)
TYPE-LOC · Maddathoray, Travancore, xi 1908 (*Annandale*)
TYPE ♀ in Ind Mus

Anisocheleomyia ? *albitarsis* Ludlow, 1905, Can Ent. xxxvii,
p 131 (♀) TYPE-LOC Pampanga, Luzon, Philippine Is TYPE.
♀ in US Nat Mus

ADULT * — Recognized by thoracic markings and white
rings on tibiae † Wing 2.5–3 mm


♀ — *Head* : a broad area of flat white scales on vertex
from back to front, bordered on either side with flat black
scales, this dark area divided from another lateral dark area
by a narrow band of white scales, lower down at side there
is another patch of white scales. A narrow line of white
scales between eyes, white scales on tori and at tips of palpi.
Clypeus and proboscis black. *Thorax* . black, with white
narrow scales on front margin of mesonotum, a pair of narrow
submedian lines of white scales from front, continued back
about half-way to level of wing-roots. A large patch of
white scales over wing-root, continued forwards more narrowly
along lateral margin of mesonotum. Three short longitudinal
white lines between wing-roots. Flat white scales on all
scutellar lobes. *Wings* . dark scaled. *Legs* fore femur
black, with a thin white line beneath, two white spots
on mid-femur, one oblong near base, one rounder nearer tip,
posterior surface black, but white at base and knee, hind
femur white on about basal $\frac{1}{2}$, white scaling continued
on outer side towards knee, a subapical black ring, wider
dorsally than ventrally, a white streak on outer side at apex,
including knee. Tibiae black, all with a white ring near
middle. Tarsi of fore and mid-legs black, with narrow basal
white rings to first two segments, wider basal white rings
to segments 1–3 of hind tarsi, 4 and 5 entirely white. *Abdomen* :
black, with narrow basal white bands on dorsum and
large lateral white patches on I–VI; VII with small lateral
white dot. Sternite black, with basal white bands.

* Edwards 1913 b, p 225, and 1922 d, p. 464 (syn.), Barraud 1923 a,
p. 778, and 1923 f, p. 224; Dyar & Shannon 1925, p. 74; Borel 1928 a,
p. 80; Bonne-Wepster & Brug 1932, p. 102

† [*A. (S.) chemulpoensis* Yamada, of Japan and China, has white
rings on tibiae, but thoracic markings quite different from *desmotes*
(more suggestive of *egypti*) and several silvery marks on anterior
surface of middle femora.]

♂.—Palpi about length of proboscis, with four white rings, the second from base being widest. Markings as in ♀ *Hypopygium* (fig. 53, a): 9t flat in middle, with small sub-

Fig. 53.



♂ hypopygial structures of *Aedes*, subgenus *Stegomyia*:
a, *dæmotes*; b, *edwardsi*; c, *raggi*; d, *annandalei*.

median hairy lobes; style with long hairs at apex but no distinct appendage; *bl* with two hairy lobes, one developed from *ehf*; paraproct with small ventral arm.

LARVA † —Not yet found in India, the following details are taken from Borel's account —Antennal hair short, single, arising rather nearer apex than base of shaft. Comb of 3 teeth, with lateral basal denticles (as in *ægypti*), these teeth originating from a semicircular chitinated plate. Siphon short, very little longer than diameter at base, tuft of 3 branches, arising rather nearer apex than base, pecten of only 2 small fringed teeth. Both *isc* and *osc* single, *lh* short, 2-branched, fan of about 6 hairs of varying lengths; papillæ about as long as fan-hairs, pointed at apices.

HABITAT —Bamboos

DISTRIBUTION.—ASSAM Golaghat *, Sibsagar dist, iv 1925 (*Barraud*). BIHAR. Pusa *, 1913 (*C S S*). BENGAL. Sukna *, Darjeeling dist, viii 1928 (*Sobha Ram*). SOUTH INDIA Ootacamund *, Nilgiri Hills, xi 1915 (*Khazan Chand*), Kadra *, N Kanara, ix 1921 (*Barraud*), Travancore, type-locality *, as above.

Known also from MALAYA, COCHIN CHINA, PHILIPPINE IS., BORNEO, and SUMBA.

111 *Aedes* (*Stegomyia*) *annandalei* Theobald, 1910.

Rec Ind. Mus iv p. 10 (♀) TYPE-LOC. Sukna, Darjeeling dist., 500', vii. 1908 (*Annandale*) TYPE ♀ in Ind Mus

var *quadricinctus* Barraud, 1923, Ind Journ Méd Res xi, p. 227 (♀). TYPE-LOC Nongpoh, Khasi Hills dist., Assam, vii. 1922 (*Barraud*) (*vide below*) TYPE ♀ in Brit Mus

ADULT ‡ —Ornamentation § shown in Pl. VI, fig 9, and Pl VIII, fig 5.

♀.—A large roundish patch of broad white scales in front of each wing-root. Scutellar scales white on lateral lobes, black on mid-lobe. *Legs*. fore femur black, with pale scaling beneath at base, mid-femur black, with white knee-spot (no white spot in middle), hind femur white on more than basal $\frac{1}{2}$ and with a white knee-spot. Tibiæ black. Tarsi of fore leg black, with narrow basal white ring to segment 1, sometimes some white scales at base of 2; white basal rings to segments 1 and 2 on mid-leg, wider basal white rings to segments 1 and 2 on hind leg, 3 and 5 black, 4 nearly all white. *Abdomen* dorsum black, with narrow basal white bands to II–VII, widening out considerably at sides. Sternites with basal white bands.

† Borel 1928 a, p. 82, Bonne-Wepster & Brug 1932, p. 103


‡ Theobald 1910 b, p. 139 (*Kingia*), Barraud 1923 a, p. 781, and 1923 f, p. 226; Borel 1928 a, p. 95; Brug 1931 b, p. 22; Bonne-Wepster & Brug 1932, p. 98

§ [*A. (S.) horseshensis* Yamada, described from ♀ only from Formosa, is extremely similar to *annandalei*, differing in having white bands of abdominal tergites subbasal instead of basal in position.]

♂.—Palpi slightly longer than proboscis, with four white markings, second from base widest and most conspicuous. Markings of hind tarsi variable* In 27 ♂♂ examined, segment 4 of hind tarsi (which is nearly all white in ♀) is white on basal $\frac{1}{2}$ or more, at least dorsally, in 19 specimens, in five others the base of this segment is only faintly or narrowly pale, and in three others it appears dark, as in *A. (S.) craggi*. *Hypopygium* (fig 53, d) style with a fairly long appendage attached some little distance from tip, *bl* with three long clubbed processes, and shorter thick hairs.

var *quadricinctus*—Differs from the type-form in having basal white markings to first four tarsal segments on all legs,

Fig 54



Larval and pupal structures of *Aedes (Stegomyia) annandalei*. a, tail-end, b, comb-plate; c, pecten-teeth, d, metathoracic pleural spines, e, antenna; f, mentum, g, paddle.

markings at base of segments 3 and 4 on fore and mid-legs small and not forming complete rings, complete white rings

* Brug (1931, p. 22) states that all ♂♂ of *A. (S.) annandalei*, reared by him in Dutch East Indies, have the 4th hind tarsal segment black, and suggests that *A. (S.) craggi*, which I described from the ♂ only, may be the ♂ of *annandalei*. This, however, is not the case, as the two species are quite distinct in the structure of the ♂ hypopygium. The ♀♀ of the two species are probably very similar, possibly indistinguishable. Brug does not describe or figure the ♂ hypopygium. Borel (1928, p. 97) describes the ♂ from Cochun China also as having the last three hind tarsal segments brown or black. In this case, from his figure of the ♂ hypopygium, it is evident that he had *annandalei* before him. The tarsal markings on the fore and hind legs are as in the ♀.

on tarsal segments 1-4 on hind legs, that on 4 occupying nearly whole segment.

LARVA (fig 54) *—*Antenna* shaft smooth, about 0.26 mm long, 6 times length of greatest width, hair resembling a short thick bristle arising nearer apex than base of shaft. Pre-elypeal spines fairly long and very slender. Lateral hairs of *thorax* of 1-3 fairly long branches. *Comb* of 5-6 large simple teeth arising in a row along posterior margin of a semicircular chitinised plate, bases of teeth finely fringed. *Siphon* about twice length of diameter at base. *Pecten* of 8 rather small simple teeth. Hair-tuft of 2-3 fine branches at about middle of tube near posterior border, *osc* long and single, *asc* 2-branched, *lh* fairly long, of 2 subplumose branches. Papillae and fan-hairs long, former rounded at ends.

HABITAT.—Tree-holes and bamboos

DISTRIBUTION.—Common in NORTH BIHAR, NORTH and EAST BENGAL, and ASSAM, and is found up to 5-6,000' in the hills. It is also present in the ANDAMANS and in SOUTH-WEST INDIA. No specimens have been seen from Burma, but it doubtless occurs there. Recorded from COCHIN CHINA and DUTCH EAST INDIES [A specimen from Huchow, Chekiang, CHINA (*Lan-Chou Feng*), differs in some respects from the Assam form, and may represent a distinct variety or species]

112. *Aedes* (*Stegomyia*) *craggi* Barraud, 1923

Ind Journ. Med. Res. xi, p. 227 (♂) TYPE-LOC Haflong, Cachar Hills, Assam, viii 1922 (*Barraud*) TYPE ♂ in Brit. Mus.

Aedes (*Stegomyia*) *purii* Barraud, 1931, Ind Journ. Med. Res. xix, p. 228 (♂) TYPE-LOC Marianbarrie Tea Estate, near Sukna, Darjeeling dist., 500', viii 1928 (*I. M. Puri*). TYPE. 2 co-type ♂♂, one in Brit. Mus., one in M S I coll., Kasauli.

ADULT.—*A. (S.) craggi* is at present known only from three ♂♂. These agree in markings with the form of ♂ of *annandalei*, in which segments 3-5 of hind tarsi are entirely dark. The ♀ is unknown, and is probably very similar to, possibly indistinguishable from, that sex of *annandalei*.

A. (S.) purii is known only from the two co-type ♂♂, which are in poor condition, but as far as can be seen they agree in markings with the form of ♂ (also the ♀) of *annandalei*, in which the 4th hind tarsal segment is mainly white, there being only a narrow black apical ring.

* Borel 1928 a, p. 97; Bonne-Wepster & Brug 1932, p. 100. Described and figured here from two damaged skins from Assam (*Barraud*) and one from Java (*Brug*), from all of which ♀♀ had been reared, presumably of this species rather than of *craggi*.

♂—*Hypopygium* (figs 53, c, 56, f) *: remarkably distinct from *annandalei* in form of *bl*, which is very large and hairy, and there are three harpago-like processes arising at the base of the plaque on the harpaginal fold

LARVA—Unknown

DISTRIBUTION.—ASSAM: Haslong, Cachar Hills, viii 1922 (2 ♂♂, including type) (*Barraud*), Nongpoh, Khasi Hills dist., vii. 1922 (*Barraud*). E HIMALAYAN foothills. type-locality of *A (S) puri*

113. *Aedes (Stegomyia) mediopunctatus* Theobald, 1905

Journ Bomb Nat. Hist Soc xvi, p 240 (♀) TYPE-LOC Para-deniya, Ceylon, xi 1901 (*Green*) TYPE ♀ in Brit Mus

var *perplexus* Leicester, 1908 (*Stegomyia perplexa*), Cul Malaya, p 83 (♂ & ♀) TYPE-LOC. Kuala Lumpur and The Gap, Selangor, Malay Penin (*Leicester*)

var *submediopunctatus* Barraud, 1923, Ind Journ. Med Res x, p 781 (♀) TYPE-LOC: Nagargah, Bombay, Doecan, viii 1921 (*Barraud*) TYPE ♀ in Brit. Mus

var. n *surelensis*. TYPE-LOC Surel, Darjeeling dist., x 1922 (*Barraud*) TYPE ♀ in Brit Mus (*vide* below).

ADULT †.—Ornamentation shown in Pl. VI, fig. 10, and Pl VIII, fig. 4. Wing about 2.5–3 mm

♀—Resembles *A (S) annandalei* fairly closely, but in that species the white area on front of mesonotum is rounder and not continued back towards the scutellum (*cf* Pl VI, figs 9 & 10), and the scutellar scales are black on mid-lobe, white on lateral lobe, whereas in the type-form of *mediopunctatus* they are white on mid-lobe, dark brown on lateral lobes. Specimens of *A (S) mediopunctatus* and *annandalei* in which the mesonotum and scutellum are denuded are difficult to separate, but this may be done on the markings of the outer side of hind femur. In the former this part is white from base to knee along the ventral border, except for a small dark interruption some distance from knee, in the latter there is a broad subapical black ring.

♂—Palpi about length of proboscis, with small white markings at base of last two segments and two broader white rings nearer base. Segment 4 of hind tarsi white, as in ♀. *Hypopygium* (fig 51, b) remarkably distinct in the form of the style, which is forked and bears a number of hairs and bristles Coxite comparatively wide, with an area at the apex carrying a group of long hairs.

* The ♂ hypopygium of *A. (S) puri* appears to be identical in structure with that of *craggi*. It seems probable, therefore, in view of what is known of variation in tarsal markings in this group of closely allied species, that *craggi* and *puri* are varieties of one species.

† Theobald 1907, p. 187; Barraud 1923 a, p 780, and 1923 f, p 226; Borel 1928 a, p. 87.

var *perplexus* Leic — Differs from type-form in having both the 4th and 5th hind tarsal segments entirely white. This form is not known in India at present, but intermediate forms occur in which there is some pale scaling at base of 5th hind tarsal segment.

var *submediopunctatus* Barraud — Differs slightly from the type-form in having only one white band on tarsi of fore and mid-legs (at base of 1st segment).

var n *surellensis*, ♀ — *Head*. as in type-form. *Thorax* a broad longitudinal median band of white scales from front margin, narrowing posteriorly, and running back to antescutellar space, where it divides, on either side of median line posteriorly, between wing-roots, are two narrow longitudinal lines of pale scales with a yellow tinge, lateral to these there are broad white scales over wing-roots, continued anteriorly along margin of mesonotum on to *ppm* and *apn*. Scutellum with flat broad white scales on all lobes. *Legs*. fore and mid-femora black, with a white line beneath for whole length, a white knee-spot on anterior aspect of mid-femur, hind femur white on basal $\frac{1}{2}$, and white scaling beneath continued to underside of knee, without any dark interruption, a white knee-spot on outer aspect, otherwise black. Tibiæ black, with white scaling beneath at base, most pronounced on hind leg, fore tibiæ, when viewed from behind, with brassy sheen for whole length. Fore and mid-tarsi black, with small white markings, chiefly on upper side, at base of segments 1 and 2. Hind tarsi with well-marked white rings to first two segments, segment 4 white except at tip, 3 and 5 black. *Abdomen* black, with narrow basal snow-white bands on IV-VII, a few pale scales at base of III, large basal lateral white patches on I-VI. Sternites hidden.

Known only from one ♀. It is very similar to the type-form, but differs in the scaling of the scutellum and in marking of hind femur. The specimen may be an unusual variation only, or there is a possibility that it may belong to a distinct species.

LARVA *.—Very similar to *A. (S.) annandalei*, according to Borel differs as follows.—Pecten-teeth with more conspicuous lateral denticles, siphonal tuft of 3 subapical branches, lateral hair of anal segment of 2 simple branches (not subplumose). In larvæ of both species the comb-teeth are 5 in number and arise from the posterior margin of a semi-circular chitinous plate.

HABITAT.—Bamboos

DISTRIBUTION (as checked by author from examination of specimens).—BOMBAY DECCAN: Nagargali, 13. viii. 1921

* Borel 1928 a, p 90

(Barraud) N KANARA Yellapur, x 1921 (Barraud)
 E HIMATAYAS. Mungpoo, Sureil, and Tindharia, all Darjeeling
 dist., x 1922 (Barraud). ASSAM. Nongpoh, Khasi Hills
 dist., vii 1922 (Barraud). Shillong, Khasi Hills dist., vi. 22
 (Barraud), Haflong, Cachar Hills, viii 1922 (Barraud),
 MALABAR COAST, x 1915 (Khazan Chand)

114 *Aedes (Stegomyia) edwardsi* Barraud, 1923

Ind Journ Med Res x, p 784 (♂ & ♀) TYPE-LOC Andaman Is.,
 ix 1911 (Christophers) TYPE ♂ & ♀ in Brit. Mus
 var *tulagiensis* Edwards, 1926, Bull Ent Res. xvii, p 101 (♀),
 TYPE-LOC near Tulagi Hospital, Santa Cruz Island, Solomon Is.,
 i 1926 (Carment) TYPE ♀ in Brit Mus (not known in Indian
 region)

ADULT †—Ornamentation shown in Pl VI, fig 5, and
 Pl VII, fig 3‡ Size and shape of small white spot on front of
 mesonotum subject to some variation Scutellar scales white
 on all lobes

♂—*Hypopygium* (fig. 53, b) 9t with prominent median
 lobe and small submedian lobes, coxite and style both rather
 long, bl small, with hairy rounded extremity

LARVA—Unknown

DISTRIBUTION—ANDAMAN Is *, ix 1911 (Christophers),
 viii 1926 (G Covell)

Recorded also from COCHIN CHINA, and the var. mentioned
 above from SOLOMON IS †

115 *Aedes (Stegomyia) w-albus* Theobald, 1905.

Ann Mus Nat. Hung. iii, p. 74 (*Stegomyia w-alba*) (♀). TYPE-LOC
 "India Orientalis," 1902 (Biró) TYPE ♀ in Nat Mus Hungary,
 Buda Pest

Stegomyia argenteomaculata Theobald, 1907, M.C. iv, p 184 (♀).
 TYPE-LOC. Narcondam Is., Bay of Bengal (G Rogers). TYPE:
 ♀ in Brit Mus

Stegomyia imitator Leicester, 1908, Cul Malaya, p 80 (♀). TYPE-
 LOC. Kuala Lumpur, Malay Penin (Leicester) TYPE ♀ in
 Brit. Mus

Stegomyia minutissima Theobald, 1910, Rec Ind. Mus. iv, p 9 (♀).
 TYPE-LOC Sukna, Darjeeling dist., 500', vii 1908 (Annandale)
 TYPE ♀ in Ind Mus

Aedes christianus Dyar, 1921, Insec Ins. Mons ix, p. 148
 (♂ & ♀). TYPE-LOC Canton and Honan, China (C. W. Howard).
 TYPES. 5 ♂♂ and 3 ♀♀ (in U.S. Nat Mus, coll. C. W. Howard.
 and coll C S Banks, Manila, P I)

ADULT ♀§.—Ornamentation shown in Pl VI, figs. 6-8,

† Borel 1928 a, p 98

‡ [A (*S*) *meronephada* (Dyar & Shannon) of Philippine Is has
 thoracic ornamentation similar to *edwardsi*, but last two hind tarsal
 segments entirely dark.]

§ Theobald 1907, p 180, and 1910, p 168 (*minutissima*); Edwards
 1913 b, p 225, and 1932, p 164 (syn.); Barraud 1923 a, p 782, 1923 f,
 p. 228, Bonne-Wepster & Brug 1932, p 107.

and Pl. VIII, fig 3, markings of thorax rather variable. Scutellar scales black at base of lobes, otherwise white, subject to some variation. Extent of white scaling on 4th hind tarsal segment shows some variation, as in other species in this group. The presence of a round white spot in middle of anterior surface of mid-femur should be noted.

♂—*Hypopygium* (fig 49) *bl* with rounded hairy tip, a few hairs proximal to this on harpaginal fold, style and its appendage both long.

LARVA—Unknown.

DISTRIBUTION—Widely spread from the PUNJAB to BIHAR and NORTH BENGAL, and through Peninsular India to CEYLON. Apparently uncommon or absent from Assam and Burma.

Known also from SIAM, SOUTH CHINA, FORMOSA, and MALAYA. Specimens sent to me from the Philippines as *A. (S.) gardneri* Ludl agreed in ♂ hypopygial structure with *w.-albus* *.

116. *Aedes (Stegomyia) albopictus* (Skuse), 1894 †

Ind Mus. Notes, III, p 20 (*Culex*) (♀). TYPE-LOC. Calcutta
TYPE ♀ in Sydney Mus

ADULT (Pl. VI, figs 2 & 14, and Pl VII, fig. 2).—Resembles the next six species in having a narrow silvery-white median line running nearly whole length of mesonotum, broad white rings on all segments of hind tarsi, etc.

♀.—Black, with snow-white markings. Scutellar scales flat and snow-white on all lobes. Markings not subject


* [*A. (S.) gardneri* Ludlow, of Philippine Is, is closely related to *w.-albus*, but differs as follows.—Subapical black ring of hind femur interrupted ventrally, white scales reaching from base to tip on under surface and on lower half of anterior surface; last hind tarsal segment mainly white, white sublateral patches of mesonotum always widely disconnected in front.]

† There seems little doubt that the above name is correctly restricted to the common Indian species which was previously known as *S. scutellaris* Theo (nec Walker), but the synonymy usually quoted is open to doubt, as it has been found quite recently that at least three other species have been confused under this name. Until an examination of the hypopygia of the type ♂♂ of *S. samarensis* Ludlow and *S. quasi-nigritha* Ludlow (which are in the U S Nat Mus) has been made, it is not possible to say whether these names should be regarded as synonyms of *A. (S.) albopictus* or not. *S. nigritha* Ludlow was described from the ♀ only, so the identity of this species must remain in doubt, as at present it has not been found possible to separate the ♀♀ of several species in this group.

The following references most probably concern *A. (S.) albopictus* as now restricted.—Theobald 1901 a, p 298 (*scutellaris*), Leicester 1908, p 86 (*scutellaris*), Edwards 1917, p 209; Barraud 1923 a, p 779, and 1931 b, p 222, Brug 1924 b, p. 41, Borel 1928 a, p 83, Martini 1930, p. 238, Bonne-Wepster & Brug 1932, p. 73.

to much variation, except width of white tarsal bands, especially that on 4th hind tarsal segment, as in some other species. A line of flat silvery scales on border of mesonotum in front of wing-root, but not continued over wing-root as in *A. (S.) scutellaris*, white scales on pleuræ in irregular

Fig 55



♂ hypopygial structures of *Aedes*, subgenus *Stegomyia* a, *flavopictus*; b, *scutellaris*, c, *albopictus*, d, *unilineatus*

patches, not in regular lines as in *A. (S.) scutellaris*; white transverse bands on abdomen at bases of segments.

♂.—Very similar to ♀. *Hypopygium* (figs. 55, c, 56, a): shape of 9t characteristic, apical border produced in middle as a blunt point.

LARVA * (fig 57, a-f) (skins definitely identified as this species from resulting ♂♂)—*Antenna*: shaft about 10 times as long as wide, smooth, with single hair arising rather more than half-way from base. *Frontal hairs* all placed far forward, all fine and inconspicuous, *A* near base of antenna, with 2-3 fine branches, *C* internal and about level with *A*, usually single, fairly long, *B* single but fairly frayed, some distance from, but directly in front of, *C*, *d* slightly anterior and internal to *B*, with a number of fine branches, *e* single, fine, and long. *Lateral hairs of thorax* of moderate length, of 1-3 branches, pleural tubercles with minute sharp thorn-like processes with several points (fig 57, f). Fairly long lateral single or 2-branched and smaller 3-branched hairs on *abdomen*. Comb of 8-12 large strong teeth, without lateral denticles, in a single row, not arising from a chitinised plate. *Siphon* rather more than twice length of diameter at base. Pecten of 7-14 small teeth, with basal lateral denticles. Tuft of 2-3 branches at about middle of siphon, usually just beyond most apical pecten-tooth. *Anal segment* nearly enclosed in chitinous ring, a few minute spines on hind margin towards dorsum, *lh* of 2 long branches. Both *isc* and *osc* single and long, or inner may be split into two. Fan-hairs fairly long but not numerous (8-10), arising from small fan-plate, papillae long, tips rounded.

HABITAT.—Tree-holes, bamboos, leaf-axils, only rarely in artificial receptacles or rock-pools.

DISTRIBUTION.—A very common species throughout India, including ASSAM and BURMA, also in the ANDAMANS and CEYLON. Occurs up to 5-6,000' in the hills.

The range probably extends throughout the Oriental region to Australia. Owing to the recent discovery of several closely allied species apparently differing only in the structure of the ♂ hypopygium, previous records—except those given by me (1931, p. 223), which were checked by examination of ♂♂—are not entirely reliable.

117. *Aëdes (Stegomyia) pseudalbopictus* Borel, 1928

Arch. des Inst. Pasteur d'Indochine, p. 85 (♂ & ♀). TYPE-LOC. Terres Rouges, Cochin China (Borel). TYPE location unknown.


ADULT †.—Apparently indistinguishable from *A. (S.) albopictus*, except in structure of ♂ terminalia. Palpi of ♂ longer than proboscis by rather more than $\frac{1}{2}$ length of terminal segment.

* Barraud 1923 *b*, p. 497; Senior-White 1927, p. 68; Borel 1928 *a*, p. 84; Martini 1930, p. 239; Bonne-Wepster & Brug 1932, p. 76.

† Barraud 1931 *b*, p. 223; Bonne-Wepster & Brug 1932, p. 81.

♂.—*Hypopygium* (fig 56, b) 9t nearly flat in middle, a pair of large hairy sublateral lobes. Coxite (when mounted flat) rather more than 3 times length of greatest width,

Fig 56



♂ hypopygial structures (style, basal plaque, and 9th tergite) of *Aedes*, subgenus *Stegomyia*. a, *albopictus*, b, *pseudalbopictus*; c, *novalbopictus*, d, *subalbopictus*; e, *flavopictus*; f, *craggi*.

bl unusually long and narrow, with one strong spine-like process and numerous hairs distal to it, several of which, at apex, are clubbed. Style comparatively long and slender, appendage stout, arising some distance from the pointed tip of style

LARVA †.—Very similar to *A. (S.) albopictus*, but (according to Borel) hair on antenna shorter and arising almost exactly at the middle point of the shaft, comb-teeth rather longer and more slender, with numerous very fine spinules at base, pecten of fewer teeth (6 noted, but this number is almost certainly subject to variation), teeth shorter and wider, less chitinated, and with lateral denticles along one side for nearly whole length Siphonal tuft of 2 branches

HABITAT.—Bamboo-stumps (Borel)

DISTRIBUTION.—NORTH BENGAL Marianbarrie Tea Estate*, and Sukna*, Darjeeling dist., viii. 1928 (Puri) BOMBAY DECCAN Nagargali*, viii. 1921 (Barraud).

Originally described from COCHIN CHINA, and since recorded from SUMATRA.

118 *Aedes (Stegomyia) novalbopictus* Barraud, 1931.

Ind Journ Med Res xix, p 224 (♂) TYPE-LOC Pusa, Bihar, vii. 1916 (S. K. S.). TYPE. ♂ in Brit Mus

ADULT.—Resembles *A. (S.) albopictus* very closely Palpi of ♂ about length of proboscis.

♂—*Hypopygium* (fig. 56, c). 9t with middle part forming a wide, slightly rounded lobe, with sometimes a very small rounded projection in the centre of apical margin, sublateral hairy lobes fairly well developed, *bl* with broad stem, numerous hairs of varying lengths, many with bent tips, arising from apex, and a row of slender flattened spines along sternal side, these spines often appear sharply pointed, but are actually slightly expanded below the tips Style moderately long and of even width throughout, appendage arising very near tip

LARVA (fig. 57, n-q) (three larval skins definitely identified as this species by examination of resulting ♂♂)—Very similar to *A. (S.) albopictus*, appears to be indistinguishable from *flavopictus*. Differs from *albopictus* in length of siphon, which is 3 times length of diameter at base, instead of only slightly more than twice. Antennal shaft 7-8 times length of greatest width.


HABITAT.—Tree-holes.

DISTRIBUTION.—♂♂ have been examined from the following places:—BIHAR: Pusa, various dates from 1908 to 1931.

† Borel 1928 a, p 87.

ORISSA: Ranchi, viii. 1922 (*T. B. Fletcher*) PUNJAB: Koti, between Kasauli and Kalka, iii 1931 (*Barraud*), bred out from tree-hole material. BOMBAY Harbour, Elephanta Island, and Trombay, vii 1921 (*Barraud*), Deccan, Belgaum, viii. 1921 (*Barraud*)

Fig. 57



Larval structures of *Aedes* (*Stegomyia*): a-i, *albopictus* (b, comb-teeth; f, base of metathoracic pleural hairs), g-j, *flavopictus*, k-m, *unilineatus*; n-q, *novalbopictus* (base of metathoracic pleural hairs, mentum, pecten- and comb-teeth)

119. *Aedes* (*Stegomyia*) *subalbopictus* Barraud, 1931.

Ind Journ. Med Res. xix, p. 225 (♂) TYPE-LOC.: Belgaum, Bombay Deccan, viii 1921 (*Barraud*). TYPE ♂ in Brit. Mus

ADULT.—Known only from the type-specimen, resembles *A. (S) albopictus* very closely in ornamentation of head, thorax, and legs, but differs in having dorsum of abdomen entirely dark, without any indication of basal white bands, though with small basal, lateral, silvery-white patches on tergites. Palpi are about same length as proboscis.

♂—*Hypopygium* (fig 56, d) 9t with middle part forming a wide curved lobe, sublateral lobes small, not projecting, each with about 4 hairs Coxite rather more than 3 times length of greatest width. *bl* with long, moderately broad stem, apex carrying a number of slightly curved blade-like processes and hairs, many of the latter with hooked tips, hairs and processes all about same length Style unusually long, tip pointed, appendage arising some little distance from it

LARVA.—Unknown

HABITAT.—Tree-holes

DISTRIBUTION.—Known only from the type-locality

120. *Aedes (Stegomyia) flavopictus* Yamada, 1921.

Ann Zool Jap x, p 52 (♂ & ♀) TYPE-LOC · Shuba, Tokyo, Japan, iv 1916 (*Yamada*) TYPE 2 ♀♀, 3 ♂♂, syntype, Govt. Inst Infect Dis, Tokio

ADULT*.—Very similar to *A (S) albopictus*, differing in mesonotal markings, which (except for the median silvery line) are usually of a yellowish colour. In some specimens the yellowish tinge is faint, or even absent, and in these cases it is very difficult, if not impossible, to separate the ♀♀ of the two species ♂ palpi only very slightly longer than proboscis

♂—*Hypopygium* (fig 56, e). 9t moderately wide, middle part produced into a rounded lobe, sublateral lobes not well developed and with few hairs, *bl*, as seen when mounted separately, with narrow stem and wide apical part carrying flattened, slightly clubbed processes, on sternal aspect, and hairs, both these and processes all almost same length Style very similar to *albopictus*, but both this and appendage rather longer

LARVA (fig. 57, g-j) (skins definitely identified as this species from resulting ♂♂) —Differs from *A (S) albopictus* in following details †.—Siphon rather more than 3 times length of diameter at base, tuft of 4-5 branches, usually arising some distance from most distal pecten-tooth; single hair towards tip of siphon on anterior border longer and more distinct, stellate hairs on thorax and abdomen with stouter branches, antenna about 8 times length of greatest breadth, head and siphon usually darker, *lh* often 3-branched

HABITAT.—Tree-holes, usually at 5,000' or more.

* Barraud 1923 f, p. 225, 1931 b, p 224.

† [In addition to the differences noted by the author, there appear to be small distinctions, perhaps of specific value, between the larvæ of the *albopictus* group in the branching of the metathoracic pleural hairs and in the form of the tubercle from which these hairs arise. These points are illustrated in fig. 57, f, g, k, n.]

DISTRIBUTION (as checked from ♂♂ by author) — ASSAM : Shillong, 5,000', vi. 1922 (*Barraud*) W HIMATAYAS Murree, 7,000', 1922 (*Gill*), Kasaui and Krol Mt, near Solan, 6-7,000', vii-ix. 1924-30 (*Barraud*) S W INDIA Coorg, Mercara, vi 1927 (*J D Baily*)

121 *Aedes (Stegomyia) scutellaris* (Walker), 1859.

- Proc Linn Soc. Lond iii, p 77 (*Culex*) (♀) TYPE-LOC · Aru Is (*Wallace*) TYPE ♀ in Brit Mus
Culex variegatus Doleschall (nec Schrank), 1858, Nat Tijd Ned.-Ind. xvii, p 77 (♀ ?) TYPE-LOC Amboma, East Indies TYPE. non-existent (formerly in Vienna Mus)
Culex zonatipes Walker, 1861, Proc. Linn Soc. Lond v, p. 229 (♀) TYPE-LOC. Dorey, New Guinea TYPE ♀ in Brit. Mus
 var *pseudoscutellaris* Theobald, 1901, Entom lxii, p 156 (♀) TYPE-LOC Suva, Fiji (*Jepson*) TYPE 3 co-type ♀♀ in Brit. Mus
 var *hebrideus* Edwards, 1926, Bull Ent Res xvii, p 102 (♂ & ♀) TYPE-LOC Espiritu Santo, Hog Harbour, New Hebrides, vi-viii 1925 (*Buxton*) TYPE ♂ & ♀ in Brit Mus
 var *tongae* Edwards, 1926, Bull Ent Res xvii, p. 103 (♂ & ♀) TYPE-LOC Haapai, Tonga Is., 26 ii 1925 (*Hopkins*) TYPE ♂ & ♀ in Brit Mus
 var *andrewsi* Edwards, 1926, Bull Ent Res xvii, p 103 (♂ & ♀) TYPE-LOC Christmas Island, S of Java, i. 1898 and x 1908 (*Andrews*) TYPE ♂ & ♀ in Brit Mus
 var *alorensis* Bonne-Wepster & Brug, 1932, Geneesk Tijd. Ned.-Ind Byblad ii, p 92 (♂) TYPE-LOC Alor, Lesser Sunda Is TYPE Geneesk Lab., Batavia, Java

ADULT* — Resembles *A. (F) albopictus* and allied species in general appearance, but differs as follows —

♂♀ — A line of flat silvery scales over each wing-root, continued nearly to lateral lobes of scutellum; two lines of similar scales across pleurae (instead of a number of irregular patches), white markings on dorsum of abdomen removed from base of each segment and curved towards middle of each tergite, sometimes forming complete bands; white scales on all scutellar lobes, but some black ones on apical border of mid-lobe

Specimens from the Andaman Islands appear to be intermediate in some respects between the type-form, as restricted by Edwards, and three of the varieties mentioned above, described by the same authority, but are distinct from var *andrewsi*, from Christmas Island, which is one of the two other localities within the Oriental region where this mosquito has been found. The following characters are present

* Edwards 1917, p. 209 (syn.); 1924, p 370, 1926 a, p 101, 1929 a, p. 4, 1932, p 165; Brug 1924 b, p 43; Barraud 1928 a, p. 654; Paine & Edwards 1929, p. 305, Bonne-Wepster & Brug 1932, p 83 (all as *variegatus* except Edwards 1932)

in specimens from the Andamans — Palpi of ♂ slightly shorter than proboscis, marked as in *A. (S.) albopictus*. White dorsal bands on abdomen complete, or only very slightly interrupted in middle, 3rd and following sternites with black apical bands. Hind femur with a large white spot at tip, white ring on 4th hind tarsal segment occupying basal $\frac{2}{3}$.

♂—*Hypopygium* middle part of 9t slightly convex, style moderately swollen at tip, 6l flat-topped, with hairs and some broader blade-like processes, all about same length, coxite comparatively long and narrow (fig. 55, b). The above-mentioned characters are variable in other forms of the species, and have been used to separate the different vars referred to above.

LARVA †—That of the Andamans form of this species is not known. Specimens of var *pseudoscutellaris*, which Dr P. A. Buxton very kindly sent to me from Samoa, show the following characters.—*Frontal hair* A 3-branched, B and C single, d anterior to all three and with a number of branches dividing off some little distance from base, B slightly internal to, and some distance in front of, C. Antenna about 6 times length of greatest width, a single hair near middle of shaft. Lateral hairs of *thorax* moderately developed, mesopleural hairs without any obvious thorn-like processes at base, but small points present at bases of metapleural hairs, no obvious stellate hairs on thorax or abdomen, but some small fine hairs with 3-4 branches. *Comb* of 10-12 strong teeth in a single row, delicately fringed at bases, not arising from a chitinated plate. *Siphon* 2-2½ times length of diameter at base. *Pecten* of 10-14 teeth, each with 2-4 denticles at base. *Hair-tuft* of 4-5 fine branches, base at about middle of tube or slightly nearer apex than base. *Anal segment* almost enclosed in chitination, some small spines on posterior margin towards dorsum, *asc* usually of 3 fairly long branches, *lh* of 2 long fine branches: fan and fan-plate moderately developed. Dorsal papillae about twice length of segment, ventral pair shorter, both pairs bluntly pointed or slightly rounded.

HABITAT—In Samoa, chiefly coconut husks and shells, tins, tree-holes, and concrete drains.

DISTRIBUTION.—In Indian region, ANDAMANS* only, vii 1926 (*Sobha Ram*).

Known also from CHRISTMAS ISLAND (S of Java), SUMATRA, DUTCH EAST INDIES (eastern islands), and PACIFIC ISLANDS.

† Buxton & Hopkins 1925, pp. 295, 297, 300; 1927, pp. 110-113; Bonne-Wepster & Brug 1932, p. 80.

122. *Aedes* (*Stegomyia*) *unilineatus* (Theobald), 1906.

2nd Rept. Welle Res Lab p 70 (*Quasistegomyia*) (♀) TYPE-
Loc.: Bahr-el-Ghazal, Sudan, ix 1905 (Bray) TYPE: ♀ in
Brit. Mus.

ADULT † (Pl VI, fig. 3) —Superficially resembles *A* (*S*) *albopictus*, but readily distinguished by presence of two round spots on posterior part of mesonotum and a round white spot on anterior surface of mid-femur

♂ —*Hypopygium* (fig. 55, d). 9t with a large median lobe, slightly hairy along border; coxite long and narrow; bl long, with rounded hairy extremity, a few small hairs on lower part near harpaginal fold

LARVA † (fig. 57, k-m) —*Antenna* comparatively short, about 6 times length of greatest breadth, hair-tuft represented by a single small strong hair, rather nearer apex than base. *Frontal hair* *A* of 2 fine barbed branches, *B* of 2 branches, each frequently widened on basal $\frac{1}{2}$ and tapering to extremity, *C* single, fairly long, lying some distance posterior to *B*. Lateral hairs of *thorax* and *abdomen* moderately developed, pleural hairs with unusually small thorn-like processes at base. Many hairs on abdominal segments of stellate form, having a small number of short branches, which are stronger and blacker in some specimens than in others. Comb of about 8 large teeth, each ending in a long sharp point, occasionally one tooth may be divided and end in two points, a few very small pointed denticles at base, teeth arranged in one row and not arising from a plate. *Siphon* a little more than twice length of diameter at base, slightly tapering. *Pecten* of 7–12 rather small pointed teeth, each with one or two lateral denticles at base. Hair-tuft of 2–4 branches, arising just beyond most distal pecten-tooth. *Anal segment* almost enclosed in a rather narrow chitinated plate, some small spines on posterior margin towards dorsum. Both *isc* and *osc* single and long, or *isc* may be divided into 2 branches, *lh* long, barbed, either single or 2-branched (apparently usually 5-branched in the African race of this species), arising from a round clear area on the chitinated plate. *Papillæ* long, about 4 times length of saddle, tips rounded. Fan with few long and some shorter hairs; no definite fan-plates.

HABITAT —Tree-holes.

DISTRIBUTION.—PUNJAB. Mahdipur*, Gurdaspur dist. (*Christophers*); Lahore*, ix. 1923 (*Sinton*); Koti*, between

† Theobald 1910, p 151 (*gelebsinensis*) and p 157 (*gelebsinensis*); Edwards 1912 a, p 13 (*Howardina*, syn.), Barraud 1923 a, p 778.

† Ingram & Macfie 1917 a, p. 81 (pupa), and 1917 b, p. 137; Kumm 1931, p. 69

Kasauli and Kalka, and Beja *, near Kasauli, Himalayan foothills, viii 1923 (*Barraud*), Karnal *, viii. 1928 (*Barraud*), Ambala *, vi 1930 (*Barraud*); Pinjaur *, Patiala State, viii. 1931 (*Jamna Dass*) DELHI PROV: Delhi *, vii 1927 (*Senior-White*) BOMBAY Harbour, Elephanta Island *, and Trombay *, vii. 1921 (*Barraud*)

Known also from TROPICAL AFRICA.

123 *Aedes* (*Stegomyia*) *albolineatus* (Theobald), 1904.

Entom xxxvii, p 77 (♀) (*Scutomyia*). TYPE-LOC. Kuala Lumpur, Malay Penin (*Leicester*) TYPE ♀ in Brit Mus.


ADULT ♀ †.—*Head*. covered with dark brown or black scales, except for a broad median silvery area on vertex, continued downwards between eyes. Clypeus, palpi, and proboscis black, last unusually long and slender *Thorax*. a median stripe of silvery scales on mesonotum, broader than in *A. (S.) albopictus*, and not continued back as far as level of wing-roots, no silvery scales over wing-roots, or upon lateral margins of mesonotum, or upon *ppn* or *apn* Scutellar scales silvery on mid-lobe, black on lateral lobes Pleuræ with large patch of silvery scales in middle *Legs*. fore and mid-femora black on upper or anterior surfaces, fore pair narrowly pale beneath for whole length, mid-pair broadly pale on basal $\frac{1}{2}$ posteriorly and with a silvery knee-spot Hind femur silvery on outer side for more than basal $\frac{1}{2}$, except for a narrow dark dorsal line from near base, a sub-apical black area and a silvery knee-spot, inner surface pale on basal $\frac{1}{2}$, with thin dark dorsal line, apical $\frac{1}{2}$ black. Tibiæ and tarsi of fore and mid-legs black with bronzy sheen, a few pale scales dorsally at base of tarsal segment 1 on mid-leg, hind tibiæ black, tarsi with rather narrow white rings at bases of segments 1 and 2, some white scales at bases of 3 not forming a complete ring. *Abdomen*: dorsum black, large lateral silvery patches on I-VII, in some specimens produced on dorsum to form narrow subbasal bands on VI and VII

♂.—Palpi black, much shorter than the rather long slender proboscis. Markings as in ♀, but tibiæ and tarsi of fore and mid-legs black, white scaling at base of segments 1 and 2 of hind tarsi only *Hypopygium* (fig 58, h). *bl* with 4 strong spines and some weaker bristles; appendage of style stout and arising nearer middle of style instead of near tip.

† Leicester 1908, p. 105 (*Scutomyia*); Edwards 1925 a, p. 258; Barraud 1927 a, p. 553; Borel 1928 a, p. 90, Paine & Edwards 1929, p. 314, Bonne-Wepster & Brug 1932, p. 94.

LARVA*.—Not isolated in India [*Antenna* moderately long, smooth Shaft-hair inserted slightly beyond middle, 1-3-branched *Frontal hairs*. *A* with several branches, *B* single and long, *C* many-branched, almost immediately behind and some distance from *B*, *d* many-branched, but rather small, internal to *B* *Thorax* and *abdomen* with some strong stellate hairs, and with dense microscopic pile on a large part of the integument (not always very evident in balsam mounts) Meso- and metapleural spines rather strong and dark Comb of 10-12 long sharp teeth in one regular row *Siphon* twice as long as broad, dark Pecten

Fig 58



♂ hypopygial characters of *Aedes*, subgenus *Stegomyia*
a, *vittatus*, b, *albolineatus* (style, basal lobe).

extending $\frac{2}{3}$ length of tube, composed of only 5-8 subequal and fairly regularly spaced teeth which have fine denticles on about their basal $\frac{1}{2}$. Tuft of 4-7 branches, just before middle of tube. *Anal segment* with numerous teeth on posterior border of saddle, *asc* forming a conspicuous tuft of 8-10 branches, *osc* single as usual, *th* 3-branched. Papillae very short]

HABITAT.—Tree-holes (*Barraud*). Tree-holes, tins, water-butts, axils of leaves of sago-palm, coconut-husks (*Paine & Edwards*, in Solomon Is.).

* Brug 1924 a, p. 433; Borel 1928 a, p. 92; Paine & Edwards 1929, p. 306; Bonne-Wepster & Brug 1932, p. 96. [Briefly redescribed above from four specimens in British Museum, from Solomon Is., agreeing essentially with Brug's description and figures.]

DISTRIBUTION.—ASSAM · Haflong, Cachar Hills, viii. 1922 (Barraud).

Known also from MALAY PENINSULA and ARCHIPELAGO, COCHIN CHINA, PHILIPPINES, BORNEO, and SOLOMON IS

124. *Aedes* (*Stegomyia*) *vittatus* (Bigot), 1861

Ann Soc Ent. France (4), 1, p. 227 (*Culex*) (♀) TYPE-LOC stated to be Corsica (perhaps in error). TYPE: ♀ in Brit. Mus

ADULT * —Distinct on ornamentation (Pl VI, figs 4 & 15, and Pl VII, fig 4) There are some small white spots on the mesonotum, preapical white rings on femora, and medial white rings on tibiae It is a somewhat aberrant species of *Stegomyia*, there are numerous upright scales on the head extending forward on to the vertex, and a varying amount of pale yellowish scaling on proboscis, characters not present in other species of the subgenus

♂ —*Hypopygium* (fig 58, a) distinct from all other species in the form of the style, which is much enlarged at the extremity and carries a preapical curved appendage

LARVA † (fig 52, g-k) —*Frontal hair* A with 4-6 branches, B, C, and d single, B slightly anterior to level of bases of antennae, C slightly internal, and some distance posterior to, B, d small, slightly internal and posterior to B, e single and fairly long Antennal shaft about 8 times as long as wide, smooth, tuft of 3 branches, rather nearer base than apex Median hairs of mouth-brushes minutely toothed apically Lateral hairs of thorax moderately developed and arising from chitinated tubercles Lateral hairs on abdominal segments II-III branched Comb of 6-9 large simple teeth, arranged in one row, but not always in a regular rank. Siphon about twice length of diameter at base Pecten of 20-34 long pointed teeth with basal lateral denticles, extending along basal $\frac{2}{3}$ of siphon, one, occasionally two, simple pecten-teeth detached from main rank and lying between hair-tuft and apex of siphon (a character distinguishing this from other known larvae of the subgenus), hair-tuft of 4-6 branches at about $\frac{1}{4}$ from base (nearer apex of tube than in most species. Anal segment with chitinated saddle, not enclosing segment,

* Theobald 1901, p. 300 (as *S. sugens* Wied.), 1905 a, p. 19, and 1907, p. 199 (*Scutomyia sugens*), 1907, p. 262 (*Reedomyia albopunctata*); Patton 1905, p. 634 (*sugens*), Neveu-Lemaire 1905, p. 8 (*S. brumpti*); Edwards 1912 a, p. 9 (*sugens*, syn.), 1917, p. 210 (syn.), 1921 c, p. 326; Bacot 1916, pp. 130, 142 (*sugens*); Barraud 1923 a, p. 777, Borel 1928 a, p. 77; Martin 1930, p. 262.

† Edwards 1912 b, p. 375 (*sugens*); Barraud 1923 h, p. 496; Senior-White 1927, pl. vii, Borel 1928 a, p. 79; Kumm 1931, p. 68.

lh single and short, *osc* single and long, *asc* of about 6 subequal branches. Papillae long and pointed, about twice length of segment. Fan moderately developed, hairs divided into branches, and arising from fan-plate.

HABITAT—Water-butts and other domestic collections of water, rock-pools, and tree-holes.

DISTRIBUTION—Widely spread from the NORTH-WEST FRONTIER to ASSAM and BURMA, and through PENINSULAR INDIA to CEYLON. At Pusa, North Bihar, it occurs throughout the year, but is commonest in the rains.

Known also from MEDITERRANEAN REGION, AFRICA, and COCHIN CHINA.

Subgenus **AEDIMORPHUS** Theobald, 1903

M. C. III, p. 290 Genotype, *Uranotaenia domestica* Theo.

Ecculex Felt, 1904, N Y State Mus Bull. p. 391 c Genotype.

Culex sylvestris Theo

Reedomyia Ludlow, 1905, Can Ent xxxvii, p. 94 Genotype.

R. pampangensis Ludl

Pecomyia Theobald, 1905, Journ Econ. Biol. 1, p. 24 Genotype.

Pec. maculata Theo

Pseudograhamsia Theobald, 1905, Journ. Bomb. Nat. Hist. Soc. xvi, p. 243 Genotype, *Ps. maculata* Theo.

Lepidotomyia Theobald, 1905, Ann Mus Nat Hung iii, p. 80

Genotype, *L. albocutellata* Theo

Geitonomyia Leicester, 1908, Cul. Malaya, p. 134. Genotype.

Culex caesus Theo.

Leslieomyia Christophers, 1911, Paludism, no. 2, p. 68 Genotype.


L. taeniorhynchoides Chr

ADULT *.—Medium-sized mosquitoes with various kinds of ornamentation and scaling. Structure of ♀ hypopygium similar to subgenus *Ochlerotatus*, but in the ♂ the phallosome is divided into lateral plates bearing teeth, there is no harpago, and the style is in many species highly modified. Proboscis usually longer than fore femur. Scales on *ppn* not broad and flat as in subgenus *Diceromyia*. Palpi of ♂ usually about length of proboscis (about $\frac{1}{2}$ as long as proboscis in *nummatus*), the last two segments more or less swollen, hairy, and turned downwards. Antennae of ♂ projecting mainly dorsally and ventrally. No lower mesepimeral bristles. Tarsal claws of fore and mid-legs of ♀ toothed, usually also larger claw on corresponding legs of ♂. Abdomen of ♀ tapering to tip, VIII completely retractile (except in *nummatus*), cerci usually long and, in dried specimens, projecting from segment VII.

* Edwards 1932, p. 165

LARVA—Antenna with spicules on shaft Siphon with the 2-3 most distal pecten teeth (except in *nummatus*) much more widely spaced than those nearer base (These characters may be present in larvæ of other subgenera)

Fig 59



♂ hypopygial structures (style, etc.) of *Aedes*, subgenera *Aedimorphus* and (k) *Banksinella* a, *trimaculatus*, b, *mediolineatus*, c, *jamesi*, d, *alboscuteclatus*, e, *pipersalatus*, f, *taeniorhynchoides*; g, *pallidostriatus*, h, i, *nummatus* (h, proctiger), j, *culicinus*, k, *lineatopennis*, l, *nigrostriatus*, m, n, o, p, *cæcus* (coxite, style, proctiger, lateral plate of phallosome).

Key to Adults.

- 1 Scutellum completely covered with flat silvery-white scales, contrasting with brown, or reddish-brown, colour of mesonotum 2.
- Scutellar scales not silvery, usually narrow and yellow (some rather broad white scales in *pipersalatus* and *cæcus*) 4
- 2 Hind tarsal segments with basal white rings only..... *jamesi*, p. 249.

- Hind tarsal segments with basal and apical pale rings *lounsi*, p 250.
Hind tarsi entirely dark 3
- 3 Abdominal tergites with pure white basal lateral spots, and sometimes with pale basal bands *alboscutellatus*, p 250
Abdominal tergites with a continuous yellow lateral stripe on each side *nivoscutellum*, p 251
- 4 Hind tarsal segments marked with distinct basal pale rings 5
Hind tarsi not ringed, or with only indistinct traces of rings at bases of first three segments 10
- 5 Wings speckled with light and dark scales 6
Wings not speckled 7
- 6 Mesonotum brown, with pale scales forming roundish spots, some rather broad white scales on mid-lobe of scutellum *perpersalatus*, p 258
Mesonotum with numerous pale scales evenly distributed, not forming spots; narrow pale scales on mid-lobe of scutellum [p 260.
taeniorhynchoides,
8.
- 7 Anterior surface of mid-femur speckled with pale scales
Anterior surface of mid-femur dark brown, without speckling of pale scales *cæcus*, p 257.
- 8 Pale scales on anterior surface of mid-femur evenly sprinkled, lobe on inner side of coxite of ♂ with numerous hooked hairs 9
Pale scales on anterior surface of mid-femur of ♀ arranged in definite spots, forming two longitudinal lines; lobe on inner surface of coxite of ♂ with one long hair and four long leaf-like processes *syntheticus*, p. 256.
- 9 Both claws of hind tarsi usually toothed in both sexes, proboscis of ♂ entirely dark brown; postnotum usually light brown or yellowish *stenætrus*, p 255.
Both claws of hind tarsi (in Indian specimens) usually simple in both sexes; proboscis of ♂ with pale area in middle; postnotum usually dark brown *vevans*, p. 253.
- 10 A large patch of white scales on front of mesonotum *nummatus*, p 265.
No patch of white scales on mesonotum 11.
- 11 Femora and tibiae marked with longitudinal pale stripes *pallidostriatus*, p. 261.
Femora and tibiae not striped 12.
- 12 Mesonotum with a pair of black stripes from the front to scutellum; a black spot in front of each wing-root *nygrostratus*, p. 262.
Mesonotum not marked with black stripes. 13
- 13 Integument of mesonotum with a median dark area, widest in front, and continued back nearly to scutellum; a shorter dark area each side posteriorly over each wing-base *trimaculatus*, p. 264
Mesonotum without such markings 14

- 14 Scales of vertex and mid-lobe of scutellum
all broad and flat, legs unusually short
and thick *pulverulentus*, p. 268.
Scales of vertex and scutellum narrow,
legs normal 15
- 15 Mesonotum with a conspicuous median
stripe of golden scales on a dark ground *ostentatio*, p. 267
Mesonotum without special adornment 16
- 16 Dorsum of abdomen dark brown, with
well-marked basal creamy bands *culcinus*, p. 252.
Dorsum of abdomen with the base and
sides of each segment ochreous, with a
triangular dark area in middle (♂), or
with an ochreous median stripe bordered
on each side with dark brown (♀) . . . *mediolineatus*, p. 263

125 *Aedes* (Aedimorphus) *jamesi* (Edwards), 1914.

Bull Ent Res v, p. 77 (*Ochlerotatus*) (♀) TYPE-LOC: Colombo,
Ceylon (*James*) TYPE ♀ in Brit. Mus.

ADULT*.—Belongs to the typical group of the subgenus,
with silvery-white scales on scutellum†. Readily distinguished
from its allies by basally ringed hind tarsal segments
Wing 3–4 mm.

♀—*Head*: narrow yellowish scales on vertex, alternating
patches of broad black and pale scales at sides, numerous
narrow upright scales on nape, extending forwards on to
vertex, a patch of narrow dark scales in middle in front
Tori, clypeus, and proboscis brown, flagellum of antenna
and palpi rather darker, the last about $\frac{1}{2}$ length of proboscis.
Thorax: mesonotal scales dark brown, with golden scales
forming indefinite spots in some specimens, especially on
front border. Scutellar scales flat and silvery. Pleuræ
dark brown, with three small patches of silvery flat scales,
a few narrow yellowish scales on *apn* and *ppn*. *Wings*:
dark scaled, usually with a small white dot at base of costa.
Legs: dark brown, silvery spots at tips of femora and tibiae;
hind tarsi with narrow basal white rings on segments 2–5
(none on 1), small white markings at bases of one or two
tarsal segments on fore and mid-legs. *Abdomen*: dark
brown, with narrow basal pale bands and lateral basal pale
patches.

♂.—Palpi dark brown, longer than proboscis by a little
more than length of apical segment; last two segments
with brown hair-tufts. *Hypopygium*: style (fig 59, c) very
similar to that of *A. (Aedim.) alboscuteellatus*, but shorter,

* Barraud 1928 a, p. 658

† [The group includes several African species and one Oriental
species (*A. orbatus* Edw.), in addition to the four occurring in India.]

with narrow basal portion shorter; coxite without any hair-tuft and without any regular row of long hairs

LARVA.—Unknown

HABITAT.—Pools in jungle (*Barraud*)

DISTRIBUTION.—A fairly common species in PENINSULAR INDIA and CEYLON, extends as far north and east as BIHAR (Pusa) There are no records from the Punjab or the north-west, nor from Assam or Burma

Not recorded from beyond limits of Indian region

126. *Aedes* (*Aëdimorphus*) *lowisi* Theobald, 1910.

M.C v, p 257 (*Reedomyia*) (♂ & ♀) TYPE-LOC Andaman Is (*Louis*) TYPE ♂ & ♀ in Brit Mus

ADULT *.—Very similar to *A jamesi*, but differing in leg-markings, as indicated in key Segment 5 of hind tarsi pale, 1 dark at base (as in *jamesi* †)

[♂—*Hypopygium*. style in type ♂ short and very broad, with narrow basal portion quite short, apparently indistinguishable from *jamesi* Coxite without special arrangement of hairs]

LARVA.—Unknown

DISTRIBUTION.—Known only from type-locality † Theobald gives Ceylon, but it was found later that the specimens to which he referred were *A jamesi*

127. *Aedes* (*Aëdimorphus*) *alboscuteatus* (Theobald), 1905.

Ann. Mus Nat Hung iii, p 80 (*Lepidotomyia*) (♀). TYPE-LOC Papua (*Bird*). TYPE ♀ in Nat Mus Hungary, Buda Pest.

Reedomyia pampangensis Ludlow, 1905, Can. Ent xxxvii, p. 94 (♀) TYPE-LOC. Pampanga, Philippine Is. (*Whitmore*). TYPE: ♀ in U S Nat Mus

Culex argentinosus Banks, 1909, Phil Journ. Sci iv, p 547 (♂ & ♀) TYPE-LOC. Rizal, Philippine Is. (*Banks*). TYPE: ♂ & ♀ in Bureau of Science, Manila

Aedes omurensis Yamada, 1921, Annot Zool Jap x, p. 73 (♂ & ♀). TYPE-LOC. Omura, Kiushu, Japan. TYPES: ♂♂ & ♀♀, co-types, in Govt. Inst. Infec Dis, Tokio.

* Barraud 1928 a, p. 658

† [The Malayan *A orbita* Edw (*taeniata* Leic.) resembles *lowisi* in having the hind tarsal rings extending over the joints, but differs in having these rings broader and whiter, segment 1 of hind tarsi also being broadly white at base. *A. orbita* is quite distinct in hypopygial structure from *lowisi*, but closely resembles *cæcus*]

[‡ Recorded by Edwards (1929 a, p. 5) from Calapan, Mindoro, Philippine Is, but comparison of the hypopygium of the ♂ from this locality with Theobald's type ♂ from the Andamans shows that it belongs to a different species, having basal $\frac{1}{2}$ or more of style slender, and a short finger-like process dorsally on distal swollen portion; coxite with row of hairs in *alboscuteatus*.]

ADULT †.—Resembles *Aedes jamesi* in ornamentation of head, thorax, femora and tibiae, but tarsi entirely dark. Abdomen with lateral basal white patches, not usually visible dorsally; narrow basal pale bands in some specimens, not in others. Wings with a white dot at base of costa as in *jamesi*. Palpi of ♂ longer than proboscis by nearly length of last two segments.

♂.—*Hypopygium* small, style as in fig 59, d. [Coxite with a rather regular row of long hairs extending almost whole length of upper (sternal) surface towards internal border, but no hair-tufts]

LAEVA.—Unknown

HABITAT.—Pools in jungle (*Barraud*)

DISTRIBUTION.—BOMBAY DECCAN Tavargatti*, Belgaum dist, viii 1921 (*Barraud*) N BENGAL Sukna*, Darjeeling dist, 500', viii 1928 (*Sobha Ram*) BIHAR Pusa*, vi 1913 (Agric Res Inst. Coll) ASSAM. Golaghat*, Sibsagar dist, xii 1924 and i 1925 (*Barraud*) BURMA Bhamo*, x 1931 (*Feegrade*)

The range extends through MALAY PENINSULA, SIAM, DUTCH EAST INDIES, and PHILIPPINE IS to AUSTRALIA and JAPAN

128. *Aedes* (Aedimorphus) *niveoscutellum* Theobald, 1905

Journ Econ Biol 1, p 22 (*Reedomyia niveoscutella*) (♀), MC iv, p 259, 1907 (♂ & ♀) TYPE-LOC India (*James & Christophers*).

ADULT ‡.—Resembles *alboscuteallatus* in its entirely dark tarsi and silvery scales on scutellum, but differs from that species, as well as from *jamesi* and *louisii*, in the following points.—*Head* with very numerous narrow scales all over vertex and nape, almost all pale, sides with flat yellowish scales, without black patches. Proboscis often largely yellowish, especially beneath. *Thorax* with the mesonotal scales uniformly light brownish, no golden scales round front margin, *ppn* more densely covered with narrow light brown scales. Scales on pleurae light brownish, concolorous with integument. *Wings* lacking the white dot at base of costa. *Legs* with silvery spots at tips of femora smaller, none at all at tips of tibiae, femora largely yellowish, except anterior surface of mid-femora; front and middle tibiae almost all yellowish, leaving only a narrow dark line (running whole

† Theobald 1907, p. 261 (*Reedomyia alboscuteallata*); Leicester 1908, p. 182 (*Lepidotomyia*), Edwards 1922 b, p. 101, 1924, p. 372 (syn)., Dyer & Shannon 1925, p. 70, Barraud 1928 a, p. 659, Martini 1930, p. 239.

‡ Barraud 1928 a, p. 659, Borel 1928 b, p. 01

length) on anterior surface of front tibia and on *under* surface of mid-tibia. *Abdomen* light or dark brown dorsally, sides of tergites yellowish, forming a stripe down each side for whole length.

[♂—*Hypopygium* †. larger than in the allied species, coxites swollen, with a rather dense patch of short hair on inner side before middle, and a large patch of very long and stout bristly hairs on outer surface towards base. Style somewhat as in *jamesi*, but rather different in shape and with a more conspicuous tuft of long hair on outer edge beyond middle.]

DISTRIBUTION.—BIHAR. Pusa*, viii 1913 (*Sharma*), vii 1927 (*Shaffi*), Kierpur*, Purneah dist, ix 1915 (*Pawa*)

Recorded also from Java and Cochin China ‡

129 *Aedes* (*Aedimorphus*) *culicinus* Edwards, 1922

Aedes (*Ecculex*) *culicinus*, Ind Journ Med Res x, p 271. TYPE-LOC Delhi, 1914 (*Christophers*) TYPE. ♂ and allotype ♀ in Brit Mus

ADULT§—Resembles *Culex fatigans* in general coloration and markings, but distinguished in ♀ by form of terminal segments of abdomen (cerci long, segment VIII retractile), in ♂ by form of palpi (only a little longer than proboscis, last two segments turned downwards and hairy), and structure of hypopygium. The ♂ style (fig. 59, j) somewhat resembles that of *Aedes alboscuteUellatus*

LARVA.—Unknown.

DISTRIBUTION.—PUNJAB AND DELHI PROVINCE only: Delhi*, as given above, Amritsar* 1911 (*Christophers*); Lahore*, ix. 1923 (*Sinton*), Karnal*, viii 28 (*Barraud*).

† [Theobald's type ♂ was badly damaged, but enough remains to show the above features on re-mounting the slide. A ♂ in the British Museum from Rizal, Luzon, Philippine Is (determined by Dyar as *niveoscutella*, and agreeing in ornamentation with ♀♀ from the same place and with those from India), differs somewhat in hypopygial structure: the long bristly hairs on the coxite are spread over the tergal surface and not aggregated towards the base, style differently shaped, its hair-tuft divided into two parts. The Philippine form would seem to be specifically distinct from the Indian, and if Ludlow's *pampangensis* is not *alboscuteUellatus* it may be this species.]

‡ [Ludlow's type of *R. pampangensis* from the Philippine Is. is said by Dyar and Shannon to be *niveoscutellum* rather than *alboscuteUellatus*, as supposed by Edwards, but the point requires confirmation; the converse is suggested by the original description.]

§ Barraud 1928 a, p 667.

130. *Aedes* (Aedimorphus) *vexans* (Meigen), 1830

Culex vexans, Syst. Bechr vi, p. 241 (♀) TYPE-LOC. near Berlin (Ruthe) TYPE 2 ♀♀ in Paris Museum

Culicada minuta Theobald, 1907, M C iv, p. 338 (♀) TYPE-LOC. : India (*Christophers*) (probably Lahore) TYPE ♀ in Brit Mus

? *Culicada eruthrosops* Theobald, 1910, M C v, p. 209 (♀) TYPE-LOC. Trincomalee, Ceylon (*Green*) TYPE ♀ in Brit Mus

var *nipponi* (*Culicada*) Theobald, 1907, M C iv, p. 337 (♀) TYPE-LOC. Kaniuzana, Japan (*Cornford*) TYPE ♀ in Brit Mus

ADULT * —Medium-sized species with little ornamentation, hind tarsal segments basally ringed. Wing about 4 mm

♀ —*Head*. narrow yellowish scales on vertex, bordered laterally by dark scales, a patch of broad flat pale scales at each side, numerous dark upright scales on nape and vertex. Palpi brown, with pale scales at tips, about $\frac{1}{4}$ length of proboscis, latter extensively pale beneath and at sides, upper side brown, with scattered pale scales. *Thorax*. integument of mesonotum almost black, covered with golden-brown narrow scales. Scutellar scales narrow and pale. Scales on *apn* and *ppn* similar to those on mesonotum. Pleuræ with several patches of broad creamy or white scales. *Wings* dark scaled. *Legs* femora brown, speckled with pale scales †, especially anterior surface of mid-pair, where the pale scales are usually numerous and evenly sprinkled. Tibiæ also speckled when viewed from front, mainly pale when seen from behind. Hind tibia without obvious pale ring at base. Tarsi dark brown, with narrow basal pale rings to first two or three segments on fore and mid-legs, and usually to all segments on hind leg. Tarsal claws of hind legs usually simple †. *Abdomen*. dorsum dark brown, tergites with pale basal bands, usually narrowed and sometimes divided in middle, also separated from the lateral pale spots. In some specimens there are apical median pale markings on tergites also.

♂ —Palpi longer than proboscis by about length of terminal segment $\frac{1}{2}$, a pale ring at about middle of long segment, white

* Edwards 1917, p. 218 (*Ochlerotatus*, syn.); 1921 c, p. 322 (*Eculex*, syn.), 1924, p. 372 (syn.), Lang 1920, p. 85, Séguy 1924, p. 86, 1925, p. 68; Brug 1924 b, p. 34, Barraud 1928 a, p. 660, Borel 1928 b, p. 68, Martini 1930, p. 259


† [In European examples of *A. vexans* there are few or no pale scales sprinkled over the dark parts of the femora, and the hind claws are usually toothed, at least in ♀, specimens from Persia resemble those of Europe, but in specimens from other regions, including those from North and South India, Central Siberia, China, Japan, and North America, the speckling is very pronounced, especially on anterior surface of mid-femora, as noted above, and the hind claws may be simple.]

‡ [In European ♂♂ the length of the palpi is subject to much variation, as shown by Peus.]

scales dorsally at bases of last two segments, which are turned downwards and bear tufts of yellowish hairs. Proboscis with indefinite pale area at about middle, especially beneath. Other markings as in ♀. Tarsal claws of hind legs usually simple. *Hypopygium* (fig 60, a) apex of style and of appendage both rather blunt. Plaque on inner surface of coxite with a number of hooked hairs.

LARVA *—[Typical European larvæ show the following characters.—*Antenna* $\frac{1}{2}$ as long as head, shaft with small

Fig 60



♂ hypopygial structures of *Aedes*, subgenus *Aedimorphus*: a, *vacans* (basal lobe, style, proctiger, and lateral plate of phallosome; also abnormal style, with double appendage); b, *stenocetrus*; c, *synthetous*.

spicules, tuft of about 7 branches, inserted at about $\frac{1}{2}$ length of shaft from base. Median hairs of mouth-brush with comb at tip. *Frontal hairs*: A with 7-12 branches; B 1-4 (usually 1 or 2), C 3-5, almost directly behind B; d 2-5, very small. *Comb* of about 10-14 sharp-pointed teeth in a patch or in one or two irregular rows. *Siphon* fully $2\frac{1}{2}$ times length of diameter at base, somewhat tapered on distal $\frac{1}{2}$.

* Brug 1824 a, p. 436, Buxton & Hopkins 1927, p. 61; Dyar 1928, p. 236; Martini 1930, p. 260.

Pecten reaching slightly beyond middle of tube, of about 20 teeth, with 1-3 strong basal denticles, last two teeth larger, simple, and widely spaced, tuft 5-7-branched, at $\frac{2}{3}$ length of tube. *Anal segment* nearly ringed by the saddle, posterior margin of which is without teeth or spines, *asc* 6-8 branched, *osc* single as usual, *lh* short and bifid, several ventral tufts before the fan proper. Papillæ nearly twice length of saddle, pointed.

No Indian larvæ are available for comparison, but specimens in the British Museum from Fiji and Samoa, as well as some (probably this species, but not isolated) from Nuwara Ehiya, Ceylon (*Major MacDougall*), differ from the European form in having fewer branches in the frontal hairs, *B* being single (in all specimens examined) and *C* single or double.]

DISTRIBUTION.—This mosquito has a very wide range, occurring in the PALÆARCTIC, NEARCTIC, and ORIENTAL REGIONS, and in several of the PACIFIC ISLANDS, occurring up to a height of 10,000 ft in TIBET (Yatung, *Lt.-Col Bailey*). In India it is fairly common from the NW FRONTIER, through NORTHERN and CENTRAL INDIA, to ASSAM, BURMA, and CEYLON. In the mountains of South India and in the higher parts of Ceylon it seems to be at least partly replaced by *Aedes stenæstrus*, described below.

131 *Aedes* (Aëdimorphus) *stenæstrus* (Theobald), 1907.

M.C. iv, p. 395 (*Culex*) (♀). TYPE-LOC Maskohya, Ceylon, 4,000' (*Green*). TYPE ♀ in Brit. Mus.
Culex pseudostenæstrus, Theobald, 1910, M.C. v, p. 343 (♀). TYPE-LOC. Hakgala, Ceylon, 8,500' (*Green*). TYPE ♀ in Brit. Mus.

ADULT †.—Very similar to *Aedes vexans*, as described above, but differs as follows:—Tarsal claws of hind legs toothed in both sexes (this character may be subject to occasional variation). Hind tibia with a more distinct narrow white ring at base. Postnotum usually pale brown or yellowish (in *A. vexans* usually dark brown). Proboscis of ♂ entirely dark brown. Palpi of ♂ longer than proboscis by only $\frac{1}{2}$ length of terminal segment. Style of ♂ *hypopygium* (fig. 60, b) with a longer and more pointed appendage, *bl* larger, and with more numerous hooked hairs.

LARVA.—Unknown.

DISTRIBUTION.—CEYLON: type-localities, as given above, Ohiya, iv 1928 (*Henry*). SOUTH INDIA. Nilgiri Hills*, 1915 (*Khazan Chand*); Coonoor* (*Patton*), Palni Hills, Kodaikanal* (*Patton*; *Fletcher*).

Not recorded from elsewhere.

† Barraud 1928 a, p. 661.

132 *Aedes* (*Aedimorphus*) *syntheticus* Barraud, 1928.


Ind Journ Med Res xvi, p 375, nom nov for *Aedes* (*Aedimorphus*) *fisheri* Barraud, 1928, Ind Journ Med Res xv, p 602 (♂ & ♀).

TYPE-LOC Bombay Deccan, Tavargatti, viii 1921 (*Barraud*),

TYPE. co-type ♂♂ and ♀♀ in Brit Mus

ADULT—closely resembles *Aedes vexans* and *A. stenocrus*, but may be distinguished as follows —Pale scales on anterior surface of mid-femur of ♀ arranged in definite small spots,

Fig 61



Larval structures of *Aedes*, subgenus *Aedimorphus*: a-d, *syntheticus*; e-h, *cecus* (a and b with details of structure of chitin of siphon and head respectively).

forming more or less two longitudinal rows (spots usually less definite in ♂) Palpi of ♂ longer than proboscis by nearly length of last segment Proboscis of ♂ with pale area in middle, the border of this area nearest the base sharply defined Apical $\frac{1}{2}$ of proboscis of ♀ blackish-brown on upper side, remainder dark brown, speckled with pale scales. Post-notum usually dark brown. Basal white abdominal bands

of fairly even width, or slightly narrowed in middle. Claws of hind tarsi simple in both sexes. Style of ♂ *hypopygium* (fig 60, c) very similar to that of *A. vexans*, but *bl* with one strong bristle and four leaf-like processes.

LARVA (fig 61, a-d).—Length about 7 mm, including siphon, which is slightly more than 1 mm long. Head yellowish-brown, dorsum covered with numerous dark spots. Antenna pale. Shaft with spinelets, most numerous on basal $\frac{1}{2}$, tuft of 5–6 branches arising near middle of shaft, two apical spines articulated a very little below tip. Preclypeal spines slender and fairly long. Frontal hair *A* with 8–10 branches, *B* single and stout, *C* standing almost directly behind *B*, with 3–5 branches, *d* very small, with about 4 fine branches, lying internal and almost level with *B*, *e* single, fine, and fairly long. Mentum with 15–16 teeth on either side of central one. Lateral hairs on *thorax* and *abdomen* moderately developed. Comb of about 10 large teeth arranged in an irregular row. Siphon brown, about 2–3 times length of diameter at base, surface covered with minute ridges; a narrow black ring at base. Acus present. Pecten of 13–19 teeth, majority with 3 lateral denticles, the two or three distal teeth larger, more widely spaced, and without lateral denticles. Siphonal tuft very small, 3–5 branches, at $\frac{2}{3}$ from base of tube. Anal segment very similar to that of *Aedes pipersalatus* (see fig 62). Papillæ about length of longest fan-hairs.

DISTRIBUTION.—BOMBAY DECCAN type-locality*, as given above. S INDIA Somwarpet*, Coorg, vi 1927 (*Barly*). MADRAS Anamali Hills*, viii 1928 (*Shaffi*).

Not known from elsewhere.

133 *Aedes* (Aedimorphus) *cæcus* (Theobald), 1901.

MC 1, p. 413 (*Culex*) (♀) TYPE-LOC Solangor, Malay Penin., x 1899 (*Butler*) TYPE ♀ in Brit Mus.

Culex *suknaensis* Theobald †, 1910, Rec Ind Mus iv, p. 21. TYPE-LOC Sukna, Darjeeling dist., 500', vii 1908 (*Annandale*) TYPE ♀ in Ind Mus.

ADULT†.—Resembles the *Aedes vexans* group of species, but differs in having dark femora, without any speckling of pale scales, the anterior surface of the mid-femur particularly being entirely dark brown. Scutellum with lanceolate

† [This was formerly quoted by Edwards as a synonym of *A. imprimens* Walk., but is, perhaps, more probably *A. cæcus*. The identity of *A. imprimens* is uncertain, as the ♂ has not been obtained from the type-locality (Ambouma), in ornamentation the type ♀ of *A. imprimens* resembles *A. cæcus* and also *A. (Bankswella) brugi* Edw. of New Guinea, the latter is possibly the ♂ of *A. imprimens*.]

‡ Theobald 1907, p. 268 (*Pecomys*), Leicester 1908, p. 135 (*Gertonomyia*), Barraud 1928 a, pp. 663–4 (*cæcus* and *imprimens*); Borel 1928 b, p. 66.

white scales, sometimes on all lobes, in other specimens mainly on lateral lobes, scales on mid-lobe being narrower. Proboscis dark brown, with usually an indefinite pale area in middle on underside and sides. Palpi of ♂ longer than proboscis by about length of last segment, a pale area at base, a pale ring in middle of long segment, white scales dorsally at bases of last two segments, these two segments with tufts of brown hairs. Tarsal segments with narrow basal pale rings, and abdomen with narrow, straight, basal pale bands.

♂—*Hypopygium* (fig. 59, o, p) style swollen at apex, with subterminal appendage and finger-like projection. Coxite with small hairy lobe on inner surface.

LAEVA † (fig. 61, e-h).—Resembles other known species of the subgenus in most details, but differs markedly in presence of a ring of spines round siphon near tip, and of a patch of similar spines at about middle of siphon on anterior surface.

HABITAT.—Natural pools in open jungle (*Barraud*)

DISTRIBUTION.—N. BENGAL: Sukma*, type-locality, as given above. E BENGAL: Rangamati*, Chittagong Hill Tracts, ix 1922 (*Barraud*). ASSAM: Dinapur* and Dibrugarh*, vii & viii 1922 (*Barraud*), Golaghat*, i 1925 (*Barraud*). S.W. INDIA: Malabar Coast*, x. 1915 (*Khasan Chand*). BURMA: Rangoon*, 1930 (*Feegrade*).

Known also from MALAY PENINSULA, DUTCH EAST INDIES, and COCHIN CHINA.

134. *Aedes* (*Aëdimorphus*) *pipersalatus* (Giles), 1901.

Giles, in Theobald, M.C. II, p. 316 (*Stegomyia*) (? only). TYPE-LOC.: Gonda, U.P. TYPE: ♀ in Brit. Mus.

Pseudograhamia maculata Theobald, 1905, Journ. Bomb. Nat. Hist. Soc. xvi, p. 243 (♂ & ♀). TYPE-LOC.: Galgamuwa, Ceylon (*Green*). TYPE: ♂ & ♀ in Brit. Mus.

ADULT ‡.—This and the following species may be distinguished from others in the subgenus by speckling of light and dark scales on wings, proboscis, and legs. Rather small species. Wing 3-3.5 mm.


♀.—*Head*: dorsum covered with narrow white and brown scales intermixed, and numerous dark upright scales; a small dark patch at each side and a larger patch of broad white scales lower down. Palpi brownish-black, with white-scaled tips. Proboscis dark brown sprinkled with pale scales. *Thorax*: mesonotum covered with dark brown scales, pale

† Brug 1924 a, p. 436, Borel 1928 b, p. 65.

‡ Edwards 1913 b, p. 227 (*Ochlerotatus*, syn.); Barraud 1928 a, p. 664.

scales forming 6-8 roundish white spots, much more distinct in some specimens than in others, some pale scales along front and lateral margins, variable in extent, and around antescutellar space. Scutellar scales white and fairly broad on all three lobes, *apn* with narrow pale scales, *ppn* with

Fig 62.



Larval structures of *Ædes*, subgenus *Aedimorphus*. a-c, *pipersalatus*; d-f, *pallidostratus*.

both pale and brown narrow scales, some patches of fairly broad pale scales on pleuræ. *Wings*: evenly sprinkled with brownish-black and white scales. *Legs*: brownish-black; femur, tibia, and 1st tarsal segment of all legs plentifully sprinkled with pale scales; white markings at tips of femora

and over tibio-tarsal joint Tarsi with narrow basal pale rings to first three or four segments on fore and mid-legs, and to all segments of hind legs. *Abdomen*: dorsum almost black, with white basal bands on segments II-VI and lateral basal pale patches, longest on segments VI and VII Sternites mainly pale scaled

♂—Palpi longer than proboscis by about length of last segment: dark brown, speckled with pale scales, a pale ring in middle of long segment and white scales at bases of last two segments Tip of long segment and last two segments with long outstanding brown hairs Antennal plumes dense, brown or yellowish according to direction of light Markings as in ♀, but white spots on mesonotum usually faintly indicated, and speckling of wings less marked *Hypopygium*: form of style as shown in fig. 59, e

LAERVA (fig. 62, a-c).—Length about 8 mm, including siphon, which is 1.25–1.75 mm long *Antenna* pale in colour, shaft with numerous spinelets, hair-tuft at about middle of shaft with 8–10 frayed branches Preclypeal spines long, moderately stout and dark. *Frontal hairs* placed as shown in fig. 62, c, all branched Mentum with 13–14 teeth either side of central one Lateral hairs of *thorax* and *abdomen* moderately developed Comb of 20–30 fringed scale-like teeth in a patch Subapical tuft small *Siphon* about 3 times length of diameter at base, surface covered with minute ridges *Acus* present *Pecten* of 14–19 teeth; the most distal two or three teeth widely spaced and without lateral denticles, the more basal teeth close together and with lateral denticles. Siphonal hair-tuft quite small, with about 5 fine branches, inserted usually about midway between most distal pecten-tooth and apex of tube *Anal segment* as in fig. 62, a.

HABITAT.—Ground-pools, water-filled ditches, etc

DISTRIBUTION.—Common from NORTH-WEST INDIA as far east as BENGAL, and through PENINSULAR INDIA to CEYLON. There are no records from Assam or Burma, or from outside the Indian region.

135. *Aedes* (*Aëdimorphus*) *tæniorhynchoides* (Christophers), 1911.

Paludism, no 2, p. 68 (*Leishnomyia*) (♂ & ♀). TYPE-LOC.: Amritsar, Punjab, ix 1910 (*Christophers*). TYPE: ♂ & ♀ in Brit. Mus.

Poomyia maculata Theobald, 1905 (name preoccupied), Journ. Econ. Biol. 1, p. 23 (♂ & ♀). TYPE-LOC.: India (*James & Christophers*). TYPE: ♂ & ♀ in Brit. Mus.

ADULT*.—Very similar to *Aedes pipersalatus*, but differs

* Edwards 1913 b, p. 227 (*Ochlerotatus*, syn.); Barraud 1928 a, p. 665.

as follows:—Mesonotal scales lighter in colour, owing to presence of more numerous pale scales, which do not form spots. Scales on mid-lobe of scutellum mainly narrow.

♂.—*Hypopygium* style (fig. 59, f) of slightly different form to that of *A. persalis*.

LARVA.—Unknown.

HABITAT.—Ground-pools, etc.

DISTRIBUTION.—PUNJAB and DELHI PROVINCE only.

136. *Aedes (Aedimorphus) pallidostriatus* (Theobald), 1907

MO IV, p. 410 (*Oulex*) (♂ & ♀) TYPE-LOC Peradeniya, Ceylon (Green), and India (*Christophers*) TYPE ♂ & ♀ in Brit Mus.

Oulex parascelos Theobald, 1910, Rec Ind Mus IV, p. 18 (♀) TYPE-LOC Madras Town, x 1908 (*Hodgart*) TYPE ♀ in Ind Mus.

ADULT*.—Belongs to a small group of medium-sized species with legs mainly yellow in posterior view, no rings on tarsi, and narrow scales on *ppn* and scutellum. Distinguished from other members of the group (described below) by striped femora and tibiae†. Wing about 4 mm.

♀.—*Head* scales nearly all yellow, numerous narrow and upright scales on vertex and nape, those at sides broader. Tori, olypeus, palpi, and proboscis yellow, apical $\frac{1}{2}$ of proboscis dark brown, also larger part of flagellum of antenna. *Thorax*. Integument brown, covered with narrow golden-brown and yellow scales, latter forming a pair of submedian lines from the front, running back to scutellum, and a lateral line on margin each side in front of wing-roots, sometimes, also, a faintly indicated median line. Scutellar scales narrow, golden. Some patches of mostly rather narrow creamy yellow scales on pleurae. *Wings* outer margin of costa yellow scaled from base to apex, otherwise scales are dark. *Legs*: femora, tibiae, and one or more tarsal segments conspicuously striped longitudinally with pale scales in front (on anterior aspect), most marked on mid- and hind legs, when viewed from behind, all legs are mainly yellow, no definite pale rings on tarsi. *Abdomen*. completely covered with bright ochre-yellow scales, without bands or spots.

♂.—Palpi mainly yellow, longer than proboscis by about length of terminal segment, last two segments turned downwards and with outstanding yellow hairs, similar hairs also at apex of long segment. Antennal plumes yellow. Other

* Edwards 1913 b, p. 228 (*Ochlerotatus*, syn.), Barraud 1928 a, p. 665.

† [The only other member of the group, apart from the four Indian species, is the African *A. ochraceus* Theo., which closely resembles *A. pallidostriatus* in both adult and larval stages, differing in the form of the ♂ style.]

markings as in ♀, but stripes on legs often less marked. Abdomen with conspicuous lateral yellow hairs, and usually some brown scaling towards apical margins of tergites. *Hypopygium* form of style as shown in fig 59, g.

LARVA (fig. 62, d-f).—Length 10–12 mm., including siphon, which is 2·5–3 mm long. *Head* comparatively large. *Antenna* moderately long, dark brown or nearly black, usually lighter at base and tip, shaft with numerous small strong spinelets, hair-tuft of about 20 subplumose or frayed branches, rather nearer base than apex of shaft. *Preclypeal spines* moderately stout and long, tapering to a fine point. *Frontal hairs* placed as shown in fig 62, f, all with a moderate number of branches. *Mentum* with 11–12 teeth either side of central one, basal teeth larger than apical. *Lateral hairs* on *thorax* and *abdomen* only moderately developed. *Comb* of about 20 teeth, each ending in a terminal spine and with a few strong lateral spines. *Siphon* very long, 6–7 times length of diameter at base; pale brown or yellow; surface covered with minute ridges, a narrow black ring at base. *Aous* present, but small. *Pecten* of 9–13 teeth, the three most distal widely spaced and without lateral denticles. *Hair-tuft* very small, usually lying about mid-way between most distal pecten-tooth and apex of tube. *Form of anal segment* as shown in fig 62, d.

HABITAT.—Open pools formed by rain or seepage, water-filled dykes, ditches, borrow-pits, etc.

DISTRIBUTION.—A fairly common species from the PUNJAB to BENGAL and through PENINSULAR INDIA to Ceylon. There are no records from Assam or Burma, or from outside the Indian region.

137. *Aedes* (*Aedimorphus*) *nigrostriatus* Barraud, 1927.

Ind Journ Med. Res xiv, p 549 (♂ & ♀). TYPE-LOC.: Golaghat, Sibsagar dist, Assam, xi. 1926 (*Barraud*). TYPE: ♂ & ♀ in Brit. Mus.

ADULT*.—Distinguished by black stripes on thorax and black knee-spots. Wing-length 3·5–4·5 mm.

♀—*Head*: covered with narrow bright yellow scales, numerous, rather narrow upright pale scales on vertex and nape, some broader yellow scales at sides. *Tori*, *clypeus*, *palpi*, and *proboscis* yellow or golden, *labella* and *flagellum* of *antenna* brown; *palpi* nearly $\frac{1}{2}$ length of *proboscis*. *Thorax*: mesonotal scales narrow and yellow, a pair of conspicuous submedian black stripes running from front margin back to lateral lobes of *scutellum*. A round black spot in

* Barraud 1928 a, p. 666

front of each wing-root, which, in some specimens, is continuous with a dark marking running downwards across the pleurae towards coxa of fore leg. Scutellar scales narrow and bright yellow. Postnotum pale yellowish or brownish, with a dark stripe on each side, corresponding with the dark stripes on mesonotum. Pleurae pale yellow or brownish, with a dark marking commencing at postspiracular area and continued downwards towards fore coxa. Some small lanceolate scales on sternopleura, mesepimeron, and postspiracular area, narrow yellow scales on *ppm*. *Wings* scales yellow and black the latter on basal $\frac{1}{2}$ of vein 1, on 2 and 4 from cross-veins to margin of wing, including branches, on 5 and 5.1. Membrane in region of cross-veins slightly darkened. *Legs* yellow or golden, with black knee-spots to femora. In some specimens anterior surface of mid-femur is brownish. *Abdomen* uniformly clothed with bright yellow scales.

♂—Antennal plumes yellow. Palpi mainly yellow, longer than proboscis by about length of apical segment, last two segments turned downwards and with tufts of yellow hairs, a dark ring on long segment and dark scales at base and apex of penultimate segment and along length of last segment. Other details as in ♀. *Hypopygium* style highly modified, as shown in fig. 59, 1.

LARVA.—Unknown.

DISTRIBUTION.—Known only from type-locality* and from Rangoon*, BURMA, 1926 (*Hamilton Miller*), and 1930 (*Feegrade*).

138. *Aedes* (*Aedimorphus*) *mediolineatus* (Theobald), 1901.

MC II, p. 113 (*Culex*) (♀) TYPE-LOC. *Thayetmyo*, Burma (*Watson*). TYPE: ♀ in Brit. Mus.

Culex trilineatus Theobald, 1901, MC II, p. 105 (♀) (name pre-occupied) TYPE-LOC. *Thayetmyo*, Burma (*Watson*). TYPE: ♀ in Brit. Mus.

ADULT †.—Distinguished from the last two, especially by abdominal markings. Wing about 4 mm.

♀—*Head*: very similar to that of *A. pallidostratus*, but clypeus and palpi darker. *Thorax*: mesonotal scales yellowish-brown, with a pair of submedian pale yellow stripes running back from front to scutellum. Scutellar scales narrow, yellow. *Wings*: dark scaled. *Legs*: brown or yellow, according to angle at which they are seen, no definite stripes or bands. *Abdomen*: a pale ochre-yellow stripe down middle of dorsum, bordered on either side with dark brown, pale yellow again at sides.

† Barraud 1928 a, p. 605; Borel 1928 b, p. 63.

♂ —Palpi yellowish, apex of long segment and last two segments brownish above, hair-tufts yellow, or brownish when viewed from side against a white background, penultimate segment slightly swollen, whole palp longer than proboscis by about length of last segment. Dorsum of abdomen with triangular dark area in middle of each tergite, with apex towards base of segment, yellow scales at sides. *Hypopygium*: style as in fig 59, b

LARVA * —Not isolated in India. From description and figures by Borel appears to be very similar to *A. pallidostriatus*, and from the details given by him it is not possible to say in what respects the two larvæ may differ. The siphon in both is very long, from 5–7 times length of diameter at base.

DISTRIBUTION —Known only from the type-locality as regards India. Recorded also from JAVA and COCHIN CHINA.

139 *Aedes (Aedimorphus) trimaculatus* (Theobald), 1905 †

Ann Mus Nat Hung. iii, p 86 (*Culex*) (♀). TYPE-LOC: Bonyay, 1902 (*Berd*) TYPE ♀ in Nat. Mus Hungary, Buda Pest

Aedes (Aedimorphus) littoralis Barraud, 1927, Ind Journ Med Res xiv, p 551 (♂ & ♀) TYPE-LOC: Trombay, Bombay Harbour, vii 1921 (*Barraud*) TYPE ♂ & ♀ in Brit Mus

ADULT ‡ —Readily distinguished by colouring of thorax. Wing 3.5–4 mm

♀ —*Head*: vertex and sides with broad pale scales, fairly numerous dark upright scales on vertex and nape, narrow golden scales along eye-margins. Palpi yellowish-brown, only about $\frac{1}{2}$ length of proboscis, latter largely yellow on basal $\frac{2}{3}$, especially beneath, dark brown apically. *Thorax*: integument of mesonotum pale at sides towards front, a median brown area, broad in front, gradually narrowing posteriorly, and continued to mid-lobe of scutellum, an oval dark area posteriorly on each side over wing-root, scales fairly dense but unusually small, golden and brown mixed. Scutellar scales narrow, golden on lateral lobes, dark brown at base of mid-lobe. *Wings*: dark scaled, except on subcosta, the scales on this vein being all yellowish. *Legs*: brown, yellowish when seen from behind, hind femur almost entirely yellow, except dorsally on anterior surface. *Abdomen*:

* Borel 1928 b, p. 64.

† [The types of *trimaculatus* and *littoralis* being from the same locality, and agreeing in the rather remarkable thoracic markings, there seems little doubt that they are conspecific. The type ♀ of *littoralis* differs somewhat from Theobald's description in regard to abdominal markings, but these may well be variable.]

‡ Barraud 1928 a, p. 666

almost entirely clothed dorsally with violet-black scales, sides of each tergite yellow from base to hind border, the yellowish area wider at base [According to Theobald abdomen dull brown, with basal dull yellow bands spreading out laterally]

♂—Antennal plumes yellowish. Palpi only slightly longer than proboscis, last two segments of about equal length, turned slightly downwards, and with tufts of brownish-yellow hairs, a tuft of hairs also at apex of long segment and a dark ring near middle, dark brown scales at apex and along the last two segments. *Abdomen* yellow scaled, with wide triangular violet-black apical bands on tergites II-IV, apex of triangle pointing towards base of each segment, narrower dark apical bands on I and V-VII. Sternites pale yellow, with faintly darker apical bands on terminal segments. *Hypopygium*. style (fig 59, a) of unusually simple form, with an appendage projecting laterally at about middle.

LARVA—Unknown

DISTRIBUTION—BOMBAY type-locality*, as given above, Deolali*, Nasik dist., x 1928 (Brit Mil Hosp) SOUTH INDIA Cannanore*, ix 1929 (*M O T Iyengar*)

Not known from elsewhere

140. *Aêdes* (*Aêdimorphus*) *nummatus* Edwards, 1923.

Bull Ent Res. xiv, p 4 (♂ & ♀) TYPE-LOC Meenglas, Jalpaiguri, N Bengal, vii 1922 (*M. O T Iyengar*) TYPE ♂ in Brit Mus

ADULT†—An aberrant species in markings and structure, differing from all others of subgenus in many respects. Wing about 3 mm

♀—*Head* entirely clad with broad flat scales, mainly dark brown, but lighter brown scales forming a median line and a border to eyes. Tori, clypeus, palpi, and proboscis dark brown, the last rather slender and about length of fore femur, palpi about $\frac{1}{2}$ length of proboscis. *Thorax*: integument of mesonotum dark brown; a large, nearly round, silvery-white patch of narrow scales anteriorly, not quite reaching front margin, scales over remainder dark brown, bristles black, numerous, and long on posterior $\frac{1}{2}$ of mesonotum. Scutellar scales dark brown, narrow on lateral lobes, broad on mid-lobe. Narrow dark brown scales on *apn* and *ppn*, some patches of flat pale scales on pleuræ. *Wings*: dark scaled. *Legs*: dark brown, undersides of femora and tibiae light, hind femur with basal $\frac{1}{2}$ almost entirely pale, also on anterior, or outer, side as far as knee, but no definite


† Barraud 1928 a, p. 667.

knee-spots *Abdomen*: dark brown dorsally, pale ventrally, not tapering towards extremity, segment VII being large, VIII not completely retractile, cerci comparatively short.

♂—Differs from ♀ as follows—Head-scales nearly all light brown. Palpi slender, about $\frac{1}{2}$ length of proboscis, tips not swollen and with few hairs. Scales on mid-lobe of scutellum light brown. *Hypopygium* style of very characteristic form (fig. 59, 1) being divided into two prongs, one of which has a moderately large terminal, pointed appendage Proctiger as in fig 59, h.

LARVA (fig 63).—Known only from one or two damaged skins, from which the following details can be made out—*Head* light brown, antennæ darker. Antennal shaft moderately long, curved, and fairly stout, thickest in middle and

Fig 63



Larval structures of *Aedes (Aedimorphus) nummatus* a, head, b, mentum, c, siphon, d, pecten-teeth, e, comb-teeth

tapering to tip, surface smooth, with very few minute spinules, hair-tuft missing, point of attachment at about middle of shaft on outer side. Pre-clypeal spines long and slender. Frontal hairs missing, except *d* on one side. *B*, *C*, and *d* placed rather far back, apparently at about middle of clypeus, point of origin of *C* slightly posterior and internal to *B*, *d* internal to, and in transverse line with, *B*, of moderate size, with about 12 fine branches. Mentum with 5-6 large teeth on either side towards base and 6-7 smaller teeth between these and a large central tooth. Mouth-brushes normal. Lateral hairs on *thorax* and *abdomen* moderately developed. Comb of about 24 fairly large pointed teeth. *Siphon* widest in middle and tapering slightly to base and apex, surface covered with minute ridges; length about 1 mm. and about 3 times length of diameter at base. Pecten extending along

basal $\frac{1}{2}$ of siphon, of about 12 very long slender teeth, fringed from near base to tip along one side, distal teeth only slightly more widely spaced than those nearer base. Hair-tuft at $\frac{2}{3}$ length of siphon from base, of 2-4 fairly long fine branches. Anal segment enclosed, except ventrally, within chitination, *etc* of 2 branches, one much longer than the other, $\frac{1}{2}$ of 3-4 moderately long fine branches. Fan-hairs fairly long (many missing), arising from fan-plate.

HABITAT.—Tree-holes and bamboos. Larvæ once found in water in broken bottle in jungle.

DISTRIBUTION.—N BENGAL. Meenglas, type-locality, as given above, Sukna*, Darjeeling dist, 500', ix 1922 (*Barraud*). ASSAM. Nongpoh*, Khasi Hills dist, vii 1922 (*Barraud*), Haflong*, Cachar Hills, viii 1922 (*Barraud*).

141. *Aedes* (?) *Aedimorphus*) *ostentatio* (Leicester), 1908†.

Cul. Malaya, p. 193 (*Aizetomyia*) (♀) TYPE-LOC. Kuala Lumpur, Malay Penin (*Leicester*) TYPE ♀ in Brit Mus.

Pseudohowardina chrysoscota Theobald, 1901, MC v, p. 228 (♀) TYPE-LOC. Peradeniya, Ceylon, iv 1907 (*Green*) TYPE ♀ in Brit Mus.

Danielsia pages Ludlow, 1911, Psyche, xviii, p. 128 (♀) TYPE-LOC. Fort Pikit, Mindanao, Philippine Is. TYPE ♀ in US Nat Mus.

ADULT †.—Rather small species, with dark legs and median golden line on thorax. Wing about 3 mm.

♀.—*Head*: a wide patch of narrow golden and pale upright scales on vertex and nape, flanked by large areas of flat black scales, alternating lateral patches of yellow and black scales. Tori pale yellow. Clypeus, palpi, and proboscis brownish-black, labella lighter; palpi rather thin and about $\frac{1}{2}$ length of proboscis. *Thorax*: mesonotum with a central stripe of narrow golden scales, wider in some specimens than in others, a broad border of similar scales round front margin, continued laterally and bending inwards in front of wing-roots; a small collection of similar scales over wing-root; remainder of mesonotum covered with dark brown narrow scales. Scutellar scales narrow, golden; *apn* rather heavily clothed with flat blackish-brown scales, *ppn* nearly bare, with dark scales on upper part. Two patches of broad white flat scales on sternopleura, and a small one on mesepimeron. *Wings*: heavily scaled with rather broad scales. *Legs*: brownish-black, femora paler beneath, and with very

† Until the ♂ has been discovered it is not certain whether this species is correctly placed in this subgenus.

‡ Edwards 1913 b, p. 228 (*Ochlerotatus*, syn.), 1922 d, p. 468 (syn.), Barraud 1928 b, p. 374.

small pale knee-spots *Abdomen* brownish-black, with large lateral pearly-white markings, continued over the dorsum as narrow basal bands on II-VII, VIII retracted within VII, cerci long and projecting from VII

♂ and LARVA.—Unknown

LARVAL HABITAT.—Pool in jungle (*Leicester*)

DISTRIBUTION.—MALABAR COAST *, x 1915 (*Khazan Chand*).

CEYLON: Peradeniya (type-locality of *P. chrysoscuta*, as given above), Bible, vii 1929 (*G. M. Henry*)

Also known from MALAYA, BORNEO, and PHILIPPINE IS

Subgenus *INDUSIUS*, subg n. †

142 *Aedes (Indusius) pulverulentus* Edwards, 1922

Ind Journ Med Res x, p 273 (♀). TYPE-LOC. Nowshera, Peshawar dist, NW India, vi 1911 (*Smith*) TYPE ♀ in Brit Mus

ADULT ‡ —Small, thick-set species with flat-scaled head, but without any distinctive ornamentation Wing 2.5–3 mm

♀ —*Head* entirely covered with broad flat scales, those on vertex grey or dull white, patches of brown ones at sides Palpi and proboscis brown, latter with sprinkling of pale scales in some specimens, palpi nearly $\frac{1}{2}$ length of proboscis, latter about length of the rather short fore femur *Thorax* mesonotum very densely covered with light yellow or fawn narrow scales, bristles only present on front margin, in front of scutellum, and over wing-roots Scutellum with flat scales on mid-lobe, dark at base, light at apex, pale narrower scales on lateral lobes Some dark brown narrow scales on upper part of *ppn*, paler ones below and on *apn* Sternopleura with patches of broad flat white scales, mesepimeron with one similar patch Bristles on *ppn* and upper part of mesepimeron fairly numerous (10–12), no lower mesepimeral *Wings*: costa dark, with sprinkling of pale scales, other veins for the most part covered with dull creamy scales *Legs*: rather short and thick femora and tibiae pale scaled, without definite markings, but with sprinkling of dark scales; tarsi darker, with, in some specimens, indefinite basal pale rings on first three segments, hind tibiae unusually bristly *Abdomen* dorsum dark greyish-brown or dark brown, with basal creamy bands, widening out at sides, on II-VII. In some specimens V-VII mainly pale scaled Segment VIII usually retracted within VII, cerci long and projecting from VII Sternites pale scaled

† [For definition of this new subgenus, *vide* Appendix.]

‡ Barraud 1928 b, p 374

♂ (?)—Two specimens from Dera Ismail Khan in the Central Research Institute collection, Kasauh, may possibly belong to this species. Although both are completely denuded of scales, they are briefly described here because of some remarkable peculiarities of structure which distinguish them from all other species of *Aedes*. Proboscis short and thick, somewhat swollen before the labella. Palpi equal in length to proboscis, slightly swollen apically. Front tarsi unusually short and thick, segment 2 scarcely more than twice as long as broad and shorter than 5, 3 and 4 each only slightly longer than broad, and together barely as long as 5, hairy beneath and with a pair of strong apical ventral spines, claws equal, simple, and as long as segment 5 or longer, empodium as long as claws and feathery. Hypopygium large and of unusual form, style broad and without appendage, a long, simple, curved process arising from protiger or base of coxite.

LARVA—Unknown

DISTRIBUTION—INDUS VALLEY only. Nowshera*, type-locality, as given above, Karachi*, viii 1926 (C M B records), Mari Indus*, vii 1927 (C M B records), Larkana*, Smd, x 1927 (*Abdul Majid*), Dera Ismail Khan* (♂ described above).

Subgenus **BANKSINELLA** Theobald, 1907.

M.C. iv, p 468. Genotype, *B. luteolateralis* Theo.

This subgenus is represented by only one species in India. It may be recognized by the characters given in the Key to Subgenera and in the following description. The females readily attack man, particularly at dusk. *A. lineatopennis* and six other species occur in Africa, and one other in Papua.

143 *Aedes* (*Banksinella*) *lineatopennis* (Ludlow), 1905

Can Ent xxxvii, p 133 (*Taniorhynchus*) (♀). TYPE-LOC Luzon, Philippine Is. TYPE ♀ in U S Nat Mus.

Culex luteolateralis Theobald, 1901, M.C. ii, p 71 (in part).


ADULT ♀†—General coloration dark brown. Sides of mesonotum conspicuously yellow scaled. Legs dark brown, unbanded. Wings with pale yellow scales on vein 1 and stem of vein 5. Abdomen with pale yellow basal bands. These characters are sufficient to distinguish this species from any other Indian mosquito; the only one with which it might be confused is *Ficalbia chamberlaini* (Ludl.), which has lateral yellow markings on the mesonotum, but has pale rings on hind tarsi.

† Leicester 1908, p 160 (*Culex luteolateralis*), Edwards 1915 a, p 274, 1924, p 371, Barraud 1928 a, p 668.

♂—Resembles ♀ in markings. Palpi apparently of only two segments, longer than proboscis by nearly length of 2nd segment, thus being upturned and hairy. Antennæ with dense plume-hairs projecting chiefly from dorsal and ventral sides of shaft. *Hypopygium* (fig 59, k) coxite fairly long, with numerous hairs on inner surface and with a projecting basal lobe carrying several strong spines. Style articulated below apex of coxite, short, widened in middle, and with a fairly long terminal appendage. Phallosome with toothed lateral plates, as in *Aedimorphus*. Paraprocts terminating in a blunt point.

LARVA * (fig 64)—*Head* moderately chitinated, antenna dark brown, with strong spicules along length of shaft, and

Fig 64



Larval structures of *Aedes (Banksinella) lineatopennis*: a, tail-end; b, head, c, mentum, d, pecten-teeth, e, comb-teeth

a hair with several fairly long branches at about middle. Frontal hairs *A*, *B*, and *C* about equally developed, each with a moderate number of fairly long branches, these three pairs forming a concave row a little posterior to level of bases of antennæ. Hair *d* very small, internal to other three pairs, and about level with *B*. Preclypeal spines slender, pale, tapering to a fine point. Hairs of mouth-brushes simple. Mentum triangular, with numerous small teeth. Lateral hairs of *thorax* and *abdomen* moderately well developed. Comb of 6-8 large sharply-pointed teeth, fringed with hairs at bases, irregularly arranged. *Siphon* about 1 mm. long. Pecten extending along rather more than basal $\frac{1}{2}$ of siphon, of 12-16 teeth, the two or three furthest from base more widely

* Ingram & de Meillon 1927, p. 61

spaced than those in main rank, each tooth with a long thin point, and one or two lateral denticles (most distal tooth apparently simple in some specimens) Hair-tuft small, of 5-8 short branches, attached slightly nearer apex of siphon than most distal pecten-tooth *Anal segment* enclosed, except ventrally, within chitination, *lh* single, fine, and fairly long, *isc* of about 8 fairly long subequal branches, *osc* single and long Fan of moderate size, 4-5 small branched hairs arising from membrane between fan-plate and base of segment Papillæ very long, narrow, and pointed

HABITAT —Open natural pools.

DISTRIBUTION —Widely spread from the PUNJAB to ASSAM and BURMA, and through PENINSULAR INDIA to CEYLON and ANDAMANS

Also known from AFRICA, MALAYA, PHILIPPINES, and AUSTRALIA

Subgenus *DICEROMYIA* Theobald, 1911.

4th Rept Welle Res Lab p 151 Genotype, *D. africana* Theo

Dendroekusea Edwards, 1929, Bull. Ent Res xx, p 341 Genotype, *Culex micropterus* Giles

ADULT —Rather small, dark mosquitoes, without conspicuous ornamentation on mesonotum, but with two white bands on head either side of middle line Vertex of head, *ppn*, and scutellum with broad flat scales Proboscis about length of fore femur Palpi of ♂ about length of proboscis, terminal segments short, somewhat thickened, hairy, and turned downwards Antennæ of ♂ as in *Aedimorphus* Mesonotal bristles very strong and numerous Usually one or more lower mesepimeral bristles Tarsal claws of ♀ all simple Segment VIII of ♀ abdomen not completely retractile, cerci short and fairly broad Coxite of ♂ hypopygium without definite lobes, style simple, with terminal, or sub-terminal, appendage, harpago absent; phallosome divided into lateral plates, with teeth, as in *Aedimorphus* The chief distinctions from the last-mentioned are to be found in the simple tarsal claws of ♀, presence of broad flat scales on *ppn* and of lower mesepimeral bristles (except in *periskeletus*), in the form of the terminal segments of abdomen of ♀, and of the style of ♂ hypopygium.

LARVA.—Resembles those of subgenus *Stegomyia*, as far as known, but comb-teeth are usually blunt-edged and fringed, and pecten-teeth have lateral denticles along one side for whole length.

DISTRIBUTION and BIONOMICS.—Five species are known from the Indian region, four being confined to that area,

the fifth having been recorded from Java also. Four other species are confined to Africa. So far as is known the early stages are passed in tree-holes and bamboo-stumps. Nothing appears to be known of the habits of the adults.

Key to Adults

- | | |
|---|---|
| 1 Femora with preapical pale rings | 2 |
| Femora without preapical pale rings | 4 |
| 2 Fore and mid-tibiae spotted in front (on anterior aspect), 1st hind tarsal segment with pale ring in middle | 3 |
| Fore and mid-tibiae not spotted in front, 1st hind tarsal segment without pale ring in middle | |
| 3 Abdomen with some round white admedian spots on dorsum | <i>perskeletus</i> , p. 272. |
| Abdomen without white round spots on dorsum | <i>nyengari</i> , p. 273. |
| 4 Pleurae dark | <i>punctipes</i> , p. 273 |
| Pleurae pale ochreous | <i>micropterus</i> , p. 275.
<i>reginae</i> , p. 277 |

144 *Aedes* (*Diceromyia*) *periskeletus* (Giles), 1902.

Handbook, 2nd ed p. 371 (*Stegomyia*) (♂) TYPE-LOC.. Shah-jahanpur, x (*Giles*) TYPE. lost

Ochlerotatus annulifemur Edwards, 1914, Bull Ent. Res v, p 77 (♂) TYPE-LOC. Jhansi, Central India, viii 1900 (*Giles*) TYPE ♂ in Brit Mus

ADULT ♂* —No specimens of this are available for re-description, none have apparently been obtained during the past thirty years, which seems remarkable, as a good deal of collecting has been done in the north of India.

The following details are taken from Edwards's description of *O. annulifemur* —*Head* clothed with broad flat scales, white at sides, black above, except for a white stripe either side of middle line. Proboscis dark brown, with narrow white ring in middle. Palpi longer than proboscis by length of last segment, dark brown, except last segment, which is pale. *Thorax*: mesonotal scales dark brown, except for some lighter scales on front margin and in front of wing-roots. Some flat white scales on scutellum (mostly denuded). *Abdomen*. black, with white basal lateral patches. *Legs*: brownish-black, preapical white rings on femora. All femora and tibiae narrowly white at tips, and all tarsal segments narrowly white at both ends. *Wings*: scales brown, a few white ones near base of subcosta.

♀ and LARVA —Unknown

DISTRIBUTION —Known only from type-localities given above.

* Barraud 1928 b, p. 360 (*A. (Skusea)*).

145. *Aedes* (Diceromyia) *punctipes* Edwards, 1921.

Bull Ent Res. xii, p. 77 (*Aedes* (Skusea)) (♀) TYPE-LOC. : Maymyo, Upper Burma, xii 1913 (Bennett) TYPE ♀ in Ind. Mus

ADULT ♀* —*Head* two patches, or bands, of pale scales on vertex, and another pale patch each side laterally. Palpi and proboscis dark brown. *Thorax*: mesonotal scales light bronzy-brown, some lighter ones on front margin. Scutellar scales broad, flat, brownish-black, some similar scales anterior to scutellum and over wing-roots. *Wings* densely clothed with dark brown scales, some pale scales at bases of costa and vein 5. *Legs* preapical pale rings and pale knee-spots to all femora, a small pale marking basal to ring on fore and mid-femora. Pale scaling over tibio-tarsal joints, fore tibia with three, mid- and hind tibiae with four, small pale spots, evenly spaced, not forming rings. Fore tarsi with pale spot at joint between segments 1 and 2, similar markings on mid- and hind legs and at following joint, segment 1 of mid-tarsi with pale medial spot, a pale ring on hind tarsi in same position. *Abdomen*: tergite I with large pale patch in middle and one at each side, II-V with pale lateral patches which are not quite basal, otherwise dorsum is dark brown.

♂ and LARVA —Unknown

DISTRIBUTION —Type-locality only. Type ♀ examined by author.

146. *Aedes* (Diceromyia) *iyengari* Edwards, 1923

Bull Ent Res. xiv, p. 4 (*Aedes* (Skusea)) (♀) TYPE-LOC. Meenglas, Jalpaiguri, vi & viii 1921 (M. O. T. Iyengar) TYPE one co-type ♀ in Brit Mus, one in Iyengar coll., Calcutta

Aedes (Skusea) *punctissimus* Barraud, 1928, Ind Journ Med Res. xvi, p. 360 (♂ & ♀) TYPE-LOC. Karwar, N Kanara, ix 1921 (Barraud) TYPE ♂ & ♀ in Brit Mus


ADULT ♀†. *Head* flat-scaled, a very few upright scales on nape, two patches, or bands, of pale scales on vertex, well separated by central dark area, another pale patch each side laterally. Palpi and proboscis dark brown. *Thorax*: mesonotal scales dark brown, with some paler ones intermixed, and a pale border to front margin. Scutellar scales as in *A. (D.) punctipes*. Broad pale scales on *apn*, and a broad line of flat white scales running from this across pleurae to upper part of mesepimeron; a small patch of similar scales on *ppn*, and one on lower part of sternopleura. Two lower mesepimeral bristles. *Wings* scales dark brown and dense, some pale scales at extreme base of costa. *Legs*: dark

* Barraud 1928 b, p. 359 (*A. (Skusea)*)

† Barraud 1928 b, p. 360 (*A. (Skusea)*).

brown, with creamy or white markings. Femora with preapical pale rings, a row of 3-4 spots basal to ring anteriorly, and small pale knee-spots. Tibiæ with about six pale markings, some not forming rings, and pale scaling over tibio-tarsal joints. Segment I of tarsi of all legs with two pale markings on middle $\frac{1}{3}$, not forming rings, and a few pale scales at tip, remaining segments dark brown. *Abdomen* dark brown, with pale markings. Tergite I with patch of pale scales on apical border in middle and large lateral silvery patches; II with lateral pale markings curving on to dorsum and meeting at base in middle line, also a pair of admedian round white spots near apical border, III-VII each with a pair of similar spots, and small lateral white markings, not visible dorsally.

Fig 65



♂ hypopygium of *Aedes*, subgenus *Diceromyia*. a, *iyengari* (with style detached); b, *micropterus*.

Sternites with basal pale bands. In some specimens, including the two co-type ♀♀ of *iyengari*, the round white spots are absent from tergites II-IV and VII. Various intermediate specimens have since been obtained.

♂—Differs from ♀ as follows:—Palpi about length of proboscis, tips slightly turned downwards and moderately hairy; a pale ring in middle of long segment, and pale markings at bases of last two segments. Proboscis dark brown, with narrow white ring at about middle. Antennal plumes directed chiefly dorsally and ventrally. Mid- and hind legs with a few pale scales at tips of tarsal segments 2 and 3. *Hypopygium* (fig. 65, a): coxite with dense patch of hooked hairs towards apex, internally. Style with large appendage, articulated some distance from apex.

LARVA †—[Not known in India, two skins from Java presented by Col Brug to the British Museum show the following characters.—*Antenna* rather shorter than in *micropterus*, scarcely tapering, about 6 times as long as broad. *Frontal hairs* as in *micropterus*, except that *B* is double. Dorsal hairs on *thorax* and *abdomen* small and inconspicuous. Mesopleural and metapleural tubercles alike, smaller than in *micropterus*, each with a very short spine. Comb much as in *micropterus*. *Siphon* longer than in *micropterus* (2–2½ times as long as breadth at base), pecten-teeth variable in number (8–14), fringed along one side and at tip, but apparently not so broad as in *micropterus*. *Siphonal hair* single, at middle of tube. *Anal segment* with moderately large saddle, with posterior edge smooth, *lh* single, at lower corner of saddle, *sc* 6-branched. *Papillæ* long and thick, about 3 times as long as segment, with rounded ends.]

HABITAT.—Bamboo-stumps

DISTRIBUTION.—Type-localities, as given above. N. BENGAL Sukna *, Darjeeling dist, 500', viii 1928 (*Puri*). S BENGAL. Calcutta *, Matlabruz, Garden Reach, x 1931 (*Senior-White*). BURMA: Rangoon *, 1930 (*Feegrade*).

Recorded also from JAVA (*Brug*)

147 *Aedes* (*Diceromyia*) *micropterus* (Giles), 1901

Journ. Bomb Nat Hist Soc xii, p. 609 (*Culex*) (♂ & ♀). TYPE-LOC Allahabad and Lucknow (*Giles*). TYPE ♂ & ♀ in Brit Mus

ADULT ♀ †.—*Head* marked with 4 pale stripes alternating with 3 dark spaces, scales mainly flat, but numerous upright scales on nape, some extending forwards on to vertex. *Palpi* and *proboscis* brownish-black. *Thorax* mesonotal scales deep brown, with lighter scales intermixed, not forming a definite ornamentation, bristles fairly numerous and strong. Scutellar scales broad, flat, deep brown or black. *Pleuræ* dark brown; 4–6 large *ppn* bristles; about 10 upper mesepimeral and 3 lower. *Wings*: dark scaled, short. *Legs*: dark brown, slightly paler on undersides and at tips of femora, also at tips of tibiae. *Abdomen*: brownish-black, pale hairs along apical margin of each tergite, and small lateral pale patches; sternites narrowly pale at base.

♂—Markings as in ♀. *Palpi* longer than *proboscis* by about length of short terminal segment, last two segment turned downwards, and with outstanding hairs. *Antennal plumes* brownish or yellowish. *Hypopygium* (fig. 65, b): coxite densely clothed with scales on outer side, numerous


† Brug 1932, p. 76.

‡ Barraud 1928 b, p. 358 (*A. (Skusea)*).

hairs along dorsal border; inner surface with small hairy lobe. Style rather long, slightly tapering, with terminal appendage. Phallosome divided into lateral plates, with teeth. Paraproct without teeth or small hairs at crown.

LARVA (fig 66).—Head and antenna moderately chitinated, latter long, somewhat tapering, with smooth shaft, and single hair on outer side, at about $\frac{2}{3}$ from base. Frontal hairs *A*, *B*, and *d* in transverse line at about level of bases of antennae: *A* with 6-8 branches, *B* single, *d* fairly large, with about 12 fine branches, *C* single, standing almost directly behind but at some distance from *B*. Preclypeal spines fairly long and slender. Mouth-brushes rather small,

Fig. 66



Larval structures of *Aedes (Diceromyia) micropterus*: a, tail-end, b, comb-teeth; c, head, d, mentum; e, pecten-teeth.

hairs of *thorax* and *abdomen* moderately developed. Some hairs on dorsum of stellate form, with fairly short branches, usually fine, but stronger in some specimens, number of branches varying from 4-8, even among specimens from the same batch. [Mesopleural tubercle rather large, with one long sharp spine, metapleural tubercle smaller, spine much shorter and quite inconspicuous]. Comb of 5-8 large teeth in a single row, each tooth ending in a blunt fringed tip. Siphon short and broad, usually almost black, 0.5-0.8 mm long. No acus. Pecten of 5-8 short strong teeth, with lateral denticles from base to tip along one side, in some specimens denticles can be seen along both sides of one or more teeth. Hair-tuft of usually only 2 branches, beyond middle point of tube from base. A fairly large chitinated

saddle covering larger part of *anal segment*, *lh* of 3 strong branches, *isc* of 3 branches. Fan of about 10 hairs arising from fan-plate, each hair split into 2-3 branches, longest of which are about length of *isc*. Ventral pair of papillæ about length of segment, dorsal pair about twice this length or more, both pairs with rounded ends.

HABITAT.—Tree-holes.

DISTRIBUTION.—Common in the PUNJAB and as far east as BIHAR (Pusa), and southwards to ORISSA and through WESTERN INDIA to MADRAS. So far as is known it is absent from Bengal, Assam, and Burma.

Not recorded from elsewhere.

148 *Ædes* (*Diceromyia*) *reginæ* Edwards, 1922

Ind Journ. Med Res v, p. 272 (*A. (Skusea)*) (♂ & ♀) TYPE-LOC.: Colombo, Ceylon, 1913 (*James*) TYPE ♂ & ♀ in Brit Mus.

ADULT †.—Differs from *A. (D.) micropterus* as follows.—Pleuræ and coxæ pale ochreous instead of dark brown, contrasting with darker mesonotum. Fewer white scales on pleuræ.

♂—*Hypopygium*: coxites with more numerous bristles on outer sides.

LARVA.—A few skins available, received many years ago from Col James, from Ceylon, closely resemble *A. (D.) micropterus*. The chief difference appears to be in the mouth-brushes, the hairs of which are stronger; the median series somewhat hooked and minutely serrated. [Chitinated plate of mesopleural tubercle larger, metapleural tubercle apparently without spine. Dorsal hairs of thorax and abdomen with only 3-4 fine branches, *isc* with 6 branches.]

HABITAT.—Tree-holes.

DISTRIBUTION.—Known only from type-locality, and from Kharghpur*, Bengal-Nagpur Rly, 26. vii. 32 (*Senior-White*).

Subgenus *ÆDES* Meigen, 1818 (s. str.).

Syst. Besch. i, p. 13. Genotype, *A. cinereus* Mg.

Verrallina Theobald, 1903, M.C. iii, p. 205. Genotype, *Ædes butleri* Theo.

Neomacleaya Theobald, 1907, M.C. iv, p. 238. Genotype, *N. indica* Theo.

Aioretomyia Leicester, 1908, Cul. Malaya, p. 185. Genotype, *A. varietas* Leic.

ADULT ‡.—Small or medium sized, dark brown or reddish-brown mosquitoes, with little or no ornamentation. The majority have lateral pale markings on the abdomen, and

† Barraud 1928 b, p. 359 (*A. (Skusea)*).

‡ Barraud 1928 b, p. 368; Edwards 1932, p. 174.

in a few these are produced on to the dorsum to form transverse bands on one or more segments. Head entirely or mainly flat scaled; palpi very short in both sexes. Antennae of ♂ with the hairs evenly spread round the segments. Proboscis about length of fore femur or slightly longer. Scutellar scales always narrow, those on *apn* also narrow, if any present. Tarsi always entirely dark. Claws of fore and mid-tarsi of ♀ toothed (in Indian species). Many species resemble one another so closely that identification can only be made with certainty by examination of the hypopygial structures. These are remarkably specialized in both sexes, and apparently quite different from those of any other genus or subgenus. In the ♂ the tip of the abdomen appears slightly swollen. The coxites are comparatively small and of remarkable form, as shown in the accompanying figures. Style articulated some distance below the apex of coxite, latter often with a process arising from inner side, or from dorsal or ventral borders, in some there are spines at ventral root. Chitinisations of proctiger (paraprocts) well developed and horn-like, sometimes forked. Phallosome usually with chitinisations, but without small teeth. In the ♀ segment VIII is partially retractile, and cerci of moderate length. The cowl, postgenital plate, spermathecal eminence, and atrial plates differ in form in various species, all these being usually heavily chitinised. Anterior lip of atrium usually membranous, insula not well developed.

LARVA.—Those of only two Indian species are at present known, they resemble those of the subgenus *Aednorphus*.

DISTRIBUTION and BIOLOGICALS.—The subgenus is chiefly Oriental, only a few species occur in the north temperate zone and in Australia, none known from Africa or Tropical America. About 40 species in all are known, 20 of these being found in the Indian region, the majority of which have been described in recent years. Little is known of the habits of the adults, but some of the commoner species attack man. Many appear to have a restricted distribution, and there is little doubt that further, as yet undescribed, species exist. So far as is known the larvæ live in open pools, water-filled ditches, or in mangrove swamps.

Key to Adults

♂♂.

- | | |
|---|----------------------------|
| 1 Tarsal claws of hind legs toothed; abdomen without any pale lateral patches on segments II-VII | <i>uniformis</i> , p. 281. |
| Tarsal claws of hind legs simple; abdomen with lateral pale patches or pale transverse bands, | 2. |

- 2 Paraprocts forked, or divided into two arms 3
 Paraprocts not forked nor divided 5.
- 3 Paraprocts divided into two pointed processes of different lengths 4
 Paraprocts divided into two arms, both truncated and slightly clubbed *clavatus*, p. 204
- 4 Style truncated at tip, with subterminal hair; shorter processes of paraproct internal to longer [p. 286.
pseudomediofasciatus,
 Style pointed, with a hair some distance from tip, shorter process of paraproct external to longer *cautus*, p. 288
- 5 Paraprocts long and pointed 6.
 Paraprocts short and rounded 10
- 6 A group of 4 or more spines at ventral root of coxite, tip of latter drawn out into several horn-like processes 7
 No spines at ventral root of coxite; tip of latter with not more than a single process 8
- 7 Style comparatively long and sinuous, with two small subterminal hairs *andamanensis*, p. 200.
 Style comparatively short and wide, with a long hair at about middle *vallestris*, p. 200
- 8 Coxite without processes on inner side, but with large tergal lobe *yerburyi*, p. 298
 Coxite with one or two processes on inner side, but no tergal lobe 9
- 9 A single elbowed process (apart from the style) arising from inner side of coxite; apex of coxite wide and truncate *nudicus*, p. 283.
 Two straight processes (apart from the style) arising from inner side of coxite; apex of coxite terminating in a single pointed process *rami*, p. 205.
- 10 Paraprocts very small, coxite terminating dorsally in two long arms, one bearing 4 spines *hirsutipleura*, p. 291.
 Paraprocts very small; coxite terminating in two curved points; sternite with a pair of terminal processes *ceylonicus*, p. 288.
 Paraprocts very small; coxite with one arm, style deeply trifid *butleri*, p. 206.
 Paraprocts comparatively very wide (other details unknown; hypopygium of type ♂ damaged) *pseudodihurnus*, p. 295.

♀♀.

1. Abdomen without any lateral pale marking on II-VII *uniformis*, p. 281.
 Abdomen with lateral pale markings or pale transverse bands 2
2. *apn* covered with silvery-white scales *yusafi*, p. 282.
apn not so covered 3.
3. Tarsal claws of hind legs toothed *abditus*, p. 282.
 Tarsal claws of hind legs simple 4.


- 4 Abdominal tergites marked with one or more complete pale bands 5
 Abdominal tergites marked with lateral pale patches not forming complete bands 6.
- 5 Tergites II-IV marked with complete pale bands; cowl markedly curved, with wide chitinated lip; brownish species of moderate size *indicus*, p. 283
 Tergite II only marked with a complete pale band, following segments with small lateral pale patches; cowl not markedly curved, membranous in middle, and without a wide chitinated lip, smaller blackish species *lugubris*, p. 294.
- 6 Numerous rather long hairs covering larger part of mesepimeron, especially lower $\frac{1}{2}$. Hairs on mesepimeron less numerous, none on lower $\frac{1}{2}$ of this sclerite 7.
 8
- 7 Mesonotal integument almost black, scales dark brown, cowl narrow in middle, widening out at sides *hirsutiplura*, p. 291.
 Similar, but cowl trilobed *comatus*, p. 292.
 Mesonotal integument and scales reddish-brown, cowl widest in middle, narrowing at sides *agrestis*, p. 293
- 8 Fairly numerous upright scales on head (nape), hypopygium comparatively very large *atrus*, p. 293.
 9
- 9 Few, if any, upright scales on head
 10 Mesonotal scales golden-brown rather than dark brown; a very large shield-like chitination on either side of spermathecal eminence *andamanensis*, p. 290
 Mesonotal scales dark brown rather than golden-brown, hypopygium of different form 10.
10. Small blackish species, lateral pale markings on abdomen hardly, if at all, visible dorsally 11.
 Dark brown species; lateral pale markings on abdomen visible from above and usually produced well on to dorsum 12
11. Postgenital plate a single large lobe, apex not usually emarginate, no definite hairy processes anterior to spermathecal eminence *butleri*, p. 296.
 Postgenital plate emarginate; a pair of hairy processes anterior to spermathecal eminence *sigmoides*, p. 295.
 12. Cowl markedly rounded 13
 Cowl much flatter and very wide *cavus*, p. 288.
- 13 A fairly large shield-like plate on either side of spermathecal eminence *vallistris*, p. 290.
 Atrial plate with characteristic pointed chitination, anterior to spermathecal eminence, nearly meeting that of other side in middle line [p. 286.
pseudomediofasciatus,

149. *Aedes* (*Aedes*) *uniformis* (Theobald), 1910*.

Rec. Ind. Mus. iv, p. 33 (*Skusea*) (♀) TYPE-LOC Pallode, 20 miles north-east of Trivandrum, Travancore, South India, xi. 1908 (*Annandale*). TYPE: ♀ in Ind. Mus.

ADULT ♀ †—A small dark brown species. Wing 2.5–3 mm. Abdomen without pale markings, except for small lateral patches of pale scales on tergite I. Head apparently without any upright scales on nape. Palpi about $\frac{1}{2}$ length of proboscis. Mesonotum deep brown, 4–5 *ppn* bristles, about 7 along posterior border of sternopleura, about 12 upper mesepimeral, but no lower. Both tarsal claws of all legs toothed. *Hypopygium* (fig. 70, a). postgenital plate slightly

Fig. 67.



Mesepimeron of three species of *Aedes*, subgenus *Aedes*, showing distribution of hairs and bristles: a, *uniformis*; b, *hirsutipleura*; c, *vallistria* (spiracle on right of each figure).

emarginate; cowl membranous, anterior lip of atrium with a pair of submedian lobes (indicated by dotted line in figure)

♂.—Resembles ♀, except for plumose antennæ. Hind tarsal claws toothed. *Hypopygium* (fig. 68, a): coxites small, especially ventrally, where they are joined by harpaginal

* [This is evidently very closely related to the Malayan *A. fragilis* Leno. and *A. indecorabilis* Leic., which have a very similar hypopygium (though different in details). Both these species (which are probably distinct from one another—not synonymous, as quoted by Edwards, 1932) have white markings on sides of abdominal tergites.]

† Theobald 1910 b, p. 491; Barraud 1928 b, p. 366.

fold, produced apically into a point. Style widened in middle and with terminal appendage Paraproct single, strongly curved, and pointed, 9t with a pair of submedian lobes and a finger-like projection on each side laterally. A pair of very elongate chitunisations apparently forming part of the phallosome.

LARVA.—Unknown.

HABITAT.—Pools in jungle

DISTRIBUTION.—BOMBAY DECCAN Tavargatti *, Belgaum dist., viii 1921 (*Barraud*), Kambarganvi *, Dharwar dist., 1929 (C M B. records). MALABAR COAST: Pudupadi *, x. 1915 (*Khazan Chand*). TRAVANCORE: type-locality *, as given above. NORTH BENGAL: Sukna *, Darjeeling dist., 500', viii 1928 (*Sobha Ram*)

Not known from elsewhere

150 *Aedes (Aedes) abditus* Barraud, 1931.

Ind Journ Med. Res. xix, p. 614 (♀) TYPE-LOC. Sukna, Darjeeling dist., 500', viii 1928 (*Sobha Ram*). TYPE: ♀ in Brit Mus

ADULT.—A small brownish-black mosquito resembling *A. (A.) uniformis* Wing 2·6 mm

♀—*Head* with a few upright scales on nape; palpi about $\frac{1}{2}$ length of proboscis. Mesonotum and scutellum brownish-black; 4 *ppn* bristles; no lower mesepimeral bristles. Tarsal claws of hind legs toothed. Abdominal tergites almost black, lighter on lateral borders, but with no distinct pale markings. *Hypopygium* (fig 70, b). postgenital plate a single lobe, basal parts resembling those of *uniformis*, but differing as shown in figures

♂ and LARVA.—Unknown.

DISTRIBUTION.—Known only from type-locality.

151. *Aedes (Aedes) yusafi* Barraud, 1931.

Ind Journ. Med Res. xix, p. 612 (♀). TYPE-LOC.: Rohanara Gardens, Delhi, iv 1914 (*Christophers*). TYPE: ♀ in Brit. Mus.

ADULT ♀—A medium-sized species (wing 2·7–3 mm.), resembling *A. (A.) indicus* in general appearance, but differing from that, and from all other known Indian species, in having *apn* completely covered with silvery-white scales. Head-scales mainly brownish-black, a patch of white scales in middle of vertex in front extending forwards between eyes, a patch of similar scales at each side of head; apparently no upright scales. Palpi about $\frac{1}{2}$ length of proboscis. Integument of mesonotum brownish-black, scales rather

lighter, many white ones around margin; 6 *ppn* bristles, 15 upper mesepimeral but no lower. Both tarsal claws on all legs toothed. Abdominal tergites almost black, with rather large basal lateral white patches extending on to dorsum but not forming complete bands, tergite I entirely dark, sternites with basal white bands and apical dark bands of about equal width. *Hypopygium* (fig 70, c) atrium and associated chitinisations small compared with size of cerci. Postgenital plate usually a single lobe, occasionally slightly emarginate at apex. Cowl lobed at each side, with marked median depression.

♂ and LARVA.—Unknown

DISTRIBUTION.—DHRI *, type-locality, as given above, type and 15 other ♀♀ PUNJAB. Amritsar *, viii 1910 (*Christophers*), Karnal *, viii. 1928 (*Barraud*)

Not known from elsewhere.

152. *Aedes* (*Aedes*) *indicus* (Theobald), 1907 †.

MC iv, p. 238 (*Neomacleaya*) (♀). TYPE-LOC : India (probably Lahore, Punjab) (*Christophers*) TYPE · ♀ in Brit. Mus

Skusea mediofasciata Theobald, 1907, MC iv, p. 544 (♂ & ♀). TYPE-LOC . India (probably Lahore, Punjab) (*Christophers*)

TYPE ♂ & ♀ in Brit Mus

ADULT ‡ —A moderate-sized dark brown species. Wing 3·3·5 mm

♀.—Abdomen brownish-black, marked with complete transverse pale bands on tergites II–IV, a pale band also on V in some specimens, in others this is divided into lateral patches as on VI and VII. The bands and patches are slightly removed from bases of segments, but that on II is often basal. Apparently no upright scales on head; 5–7 *ppn* bristles, 12–16 upper mesepimeral, but no lower. Tarsal claws of hind legs simple. *Hypopygium* (fig 70, d) · cowl strongly curved and with wide chitinised lip. Numerous long hairs, apparently arising from spermathecal eminence, obscuring the form of the atrial plates.


♂ —Differs from ♀ in having lateral pale spots on abdominal tergites instead of bands, plumose antennæ, and in its smaller size. *Hypopygium* (fig. 68, b) : coxite with broad, blunt projection at apex. Style slender, curved, with small pointed terminal appendage. A single slender, elbowed process arising from inner surface of coxite. Paraproct single, long, slightly curved, and pointed. No spines at ventral root

† [The Philippine *A. nigrotarsis* Lucil. is closely allied, differing in details of hypopygial structure.]

‡ Edwards 1913 b, p. 229 (syn.); Barraud 1928 b, p. 366.

LARVA (fig. 69).—*Head* brown, darker posteriorly. Antenna light brown, a few small dark spicules on shaft, hair-tufts usually of 3 stout frayed branches, attached rather nearer

Fig. 68.




♂ hypopygia of *Aedes*, subgenus *Aedes* a, *uniformis*; b, *indicus* (dorsal, ventral, and posterior views), c, *clavatus*. pc, prolongation or processes at end of coxite, ic, process on inner side of coxite. Other lettering as on p. 4.

base than apex; apical bristles all arising very near tip of shaft. Preclypeal spines pale, slender, fairly long, and tapering to a point. Median hairs of mouth-brushes without

obvious serrations or teeth. Frontal hairs placed as shown in figure. *A* with 7-13 branches, *B* and *C* usually of 3 branches, the middle one stout. In one specimen hair *B* is a single, very stout, frayed bristle on one side of head, hair *d* small, with 3-4 very fine branches, hair *e* fine and usually single, but may be split. Mentum with about 17 teeth on either side of central one, the teeth very regular in size. Larger lateral tufted hairs on *thorax* moderately developed, meso- and metapleural tubercles with small spines at bases. *Comb* of about 10 large teeth arranged more or less in a row, teeth delicately fringed on basal $\frac{1}{2}$. *Siphon* pale brown, surface covered with minute ridges, a very narrow dark ring at base, length along dorsal border in side view from 0.70

Fig. 69.

Larva of *Aedes* (*Aedes*) *indicus* Theo.

to 0.82 mm., and about $2\frac{1}{2}$ times length of diameter at base. Acus moderately developed. Pecten of 11-14 long teeth, all with lateral denticles, though those on distal teeth may be quite small. The more distal one or two teeth usually more widely spaced than those in main rank. Hair-tuft small, with 3-5 short, fine branches, attached between most distal pecten-tooth and apex of siphon. *Anal segment* enclosed, except ventrally, with chitination; *isc* of about 8 rather short branches all about same length; *lh* single. Fan-hairs of only moderate length, about 12 arising from fan-plate, each split into a number of branches; 2 or 3 branched hairs arising from membrane between fan-plate and base of segment. Papillæ apparently about length of longest fan-hairs.

HABITAT.—Open pools, rain-filled ditches, etc

DISTRIBUTION.—PUNJAB. Amritsar *, viii 1910, and iv 1911 (*Christophers*); Lahore * (*Christophers*), and viii 1923 (*Sinton*), Karnal *, vii. 1927, vii 1928, viii 1930, and x 1931 (*Barraud*) DELHI * iv 1914 (*Christophers*) SIND. Larkana *, x 1928 (*Baily*) UNITED PROVINCES. Saharanpur *, ix 1927 (*Sinton*) BIHAR. Pusa *, various dates, vi –x 1913–1929 (*Shaffi*) MADRAS. Madras Town *, iv 1911 (*Justice*)

Not known from beyond limits of Indian region

153 *Aedes (Aedes) pseudomediofasciatus* (Theobald), 1910.

MC v, p 489 (*Skusea*) (♂) TYPE-LOC. Paradeniya and Hakgala, Ceylon, iii & iv 1907 (*Green*) TYPE. 2 co-type ♂♂ in Brit. Mus

ADULT ♀†—Moderate-sized species (wing 3.5 mm.), thorax dark reddish-brown, abdomen darker. Lateral pale markings on tergites II–VII, usually produced well on to dorsum, and in some specimens forming almost complete bands on VI or VI–VII. Head with a few upright scales on nape and usually a few narrow pale scales in middle of vertex, 5–6 *ppn* bristles, 8 upper mesepimeral, and a few hairs in middle of the sclerite just below a patch of scales. *Hypopygium* (fig 70, e). cowl strongly rounded. Atrial plates with a characteristic pointed chitunisation on each side

♂.—Very similar to ♀ *Hypopygium* (fig 71, a) coxite with 4–5 blunt teeth at apex, all about same length. Style truncate at tip, with small subterminal hair. Numerous hairs and some strong bristles on inner surface of coxite. Paraproct forked, the shorter arm internal to longer


LARVA.—This has not been isolated in India, but has been figured by Senior-White (1927, pl xi) from Ceylon. It appears to be very similar to that of *A. (A.) indicus* described above, but the comb-teeth are shown as having lateral fringes to the tip, whereas in the larva referred to the fringe is confined to the basal $\frac{1}{4}$

HABITAT.—Hill-stream, in company with *Anopheles maculatus*, also in swamp

DISTRIBUTION.—BOMBAY DECCAN. Kambarganvi *, Dharwar dist., 1929 (C.M.B. records) SOUTH INDIA: Nilgiri Hills *, x 1915 (*Khazan Chand*), Kallar, xii. 1922 (*T. B. Fletcher*) MADRAS: Gundy * (Madras Town), viii. 28 (C.M.B. records) CEYLON: type-localities, as given above, and Colombo, 1913 (*James*), Marble, Kurunegalla dist. (*Senior-White*)

† Edwards 1917, p. 222, Barraud 1928 b, p. 367.

Fig 70



♀ hypopygia of *Acides*, subgenus *Acides*: a, *uniformis*; b, *abditus*; c, *grisea*; d, *indicus*; e, *pseudomediofasciatus*; f, *cautus*; g, *andamanensis*; h, *vallistris*. Floor of atrium not shown.

Lettering: AA, anterior lip of atrium; AP, shield-like plate on either side of spermathecal eminence; CL, cowli; CR, cercus; RGP, postgenital plate; SEM, spermathecal eminence.

154. *Aedes (Aedes) cantus* Barraud, 1928.

Ind Journ Med Res xvi, p 368 (♂ & ♀) TYPE-LOC Tavargatti, Bombay Deccan, Belgaum dist, viii 1921 (Barraud) TYPE. ♂ & ♀ in Brit Mus

ADULT—Resembles *A. (A.) pseudomediofasciatus* very closely in both sexes, but differs in structure of hypopygium.

♀—*Hypopygium* (fig 70, f): cowl wide and nearly flat in middle, outer angles forming nearly square shoulders Atrial plates with a characteristic oval chitination on each side.

♂—*Hypopygium* (fig 71, b). apex of ooxite produced into 5 strong tooth-like processes, the one at extreme tip longer than others Style curved and pointed, with a hair arising some distance from tip. Four strong spines arising from inner surface of coxite towards ventral root. Paraproct forked, shorter arm external to longer

LARVA—Unknown

HABITAT—Ground-pools in jungle

DISTRIBUTION—BOMBAY DECCAN: Tavargatti*, type-locality, as given above, Kambarganvi*, Dharwar dist., 1929 (CMB records) MALABAR COAST: Pudupadi*, x. 1915 (*Khazan Chand*)

155. *Aedes (Aedes) ceylonicus* Edwards, 1917.

Bull. Ent Res. vii, p 221 (♂) TYPE-LOC.: Colombo, Ceylon (*K. McGahey*). TYPE. ♂ in Brit. Mus.

[ADULT—Closely resembles *A. pseudomediofasciatus*, no external differences being apparent No scales on *apn*, a few small hairs in middle of mesepimeron


♂—*Hypopygium* (fig. 73, g): sternite rather large and swollen, with a pair of pointed bare projections on margin, coxite with tip slightly produced and divided into two sharp points, tergal point with a small tooth, sternal point with a small patch of hairs below it, but no spines, style minute, with one hair at middle and another at tip; no chitinous processes from anal segment, but some small internal processes at root of coxite, phallosome with a pair of long rods.

♀ (if correctly associated with ♂) —*Hypopygium*: somewhat intermediate between *A. pseudomediofasciatus* and *A. atrius*, with rather more resemblance to the latter

PUPA and LARVA.—Unknown.

DISTRIBUTION—Known only from Colombo, CEYLON (*K. McGahey*, 1914, *G. M. Henry*, 14–19. x. 1914); specimens captured indoors]

Fig 71.



♂ hypopygia of *Aedes*, subgenus *Aedes*: a, *pseudomediofasciatus*; b, *cautus*; c, rami; d, *vallisstris*; e, *hirsutipleura*; f, *pseudodurinus*; g, *andamanensis*. PC, prolongation or processes at end of coxite; VR, spines at ventral root of coxite. Other lettering as on p. 4.

156. *Aedes (Aedes) andamanensis* Edwards, 1922 †

Ind Journ Med. Res. x, p 272 (♂) TYPE-LOC Andaman Is.,
ix 1911 (Christophers) TYPE ♂ in Brit Mus

Neomacleaya indica var *simplex* Theobald, 1908, Rec Ind Mus ii,
p 291 (♀) TYPE-LOC Sylhet, Assam, vi 1905 (Hall) TYPE
♀ in Ind. Mus

ADULT ♀ † —Dark brown, of moderate size (wing about 3.5 mm) Lateral pale markings on abdominal tergites, very variable in size, but usually produced on to dorsum Mesonotal scales deep golden brown, 7-8 *ppn* bristles; about 15 bristles and some hairs along upper and posterior borders of sternopleura, about 15 upper mesepimeral bristles, and below these numerous hairs scattered over the larger part of the sclerite, except lower $\frac{1}{3}$ Head with narrow yellowish border to eyes, some pale scales in middle of vertex, and a few upright scales on nape. *Hypopygium* (fig 70, g). cowl flat, wide, and narrow A very large shield-like plate on either side of spermathecal eminence, apparently a development of the atrial plate.

♂ —Resembles ♀. Wing about 2.5 mm Antennae plumose *Hypopygium* (fig. 71, g): coxite very much drawn out at apex into a wide arm with two horns. Ventrally a three-pronged process and two other processes arise at, or near, apex of coxite Style slender, curved, and pointed, with 2 hairs arising some distance from apex Six strong spines at ventral root Paraproct single, long, and slender.

LARVA —Unknown.

DISTRIBUTION —ANDAMAN Is * and Sylhet *, ASSAM, type-localities, as given above, Golaghat *, Sibsagar dist., xii. 1924, i 1925 (Barraud) E BENGAL. Chittagong *, viii. 1922 (Barraud) MALABAR COAST Pudupachi *, x. 1915 (Khazan Chand)

Known also from COCHIN CHINA, SUMATRA, and PHILIPPINES *

157. *Aedes (Aedes) vallistris* Barraud, 1928.

Ind Journ Med Res. xvi, p 369 (♂ & ♀). TYPE-LOC.: Golaghat, Sibsagar dist., Assam, xi & xii 1924 (Barraud). TYPE: ♂ & ♀ in Brit Mus.

ADULT ♀.—A small species (wing about 3 mm.) with lateral pale markings on abdominal tergites, variable in size, but usually slightly produced on to dorsum, 5 *ppn* bristles;

† [A sketch sent by Dyar of the hypopygium of *A. margarsen*, described by him in 1926 from the Philippine Is., shows great similarity in structure to *A. andamanensis* As the present author notes having received *andamanensis* from the Philippines it is possible that *margarsen* may be a synonym of this species.]

‡ Barraud 1928 b, p. 368; Edwards 1928 a, p 273.

about 10 sternopleural along upper and posterior borders, 10-12 upper mesepimeral, and some small hairs at about middle of this sclerite (fig. 67, c). *Hypopygium* (fig. 70, h) atrial chitinisations somewhat resembling those of *A. (A.) andamanensis*, but shield-like plates smaller and cowl much more rounded.

♂.—Resembles ♀ but is rather smaller. Antennæ plumose. *Hypopygium* (fig. 71, d) : coxite with two terminal pointed processes, and a third thumb-like process at about middle of dorsal border. Four spines at ventral root. Style wide, curved, and pointed, with a long hair arising at about middle. Paraproct single, and only of moderate length.

LARVA.—Unknown.

DISTRIBUTION.—ASSAM. Golaghat*, type-locality, as given above. BURMA. Rangoon*, 1931 (*Feegrade*).

Not recorded from elsewhere.

158 *Aëdes (Aëdes) hirsutipleura* Barraud, 1928.

Ind Journ. Med Res. xvi, p. 370 (♂ & ♀). TYPE-LOC. Golaghat, Sibsagar dist., Assam, xii 1924 & i 1925 (*Barraud*). TYPE ♂ & ♀ in Brit Mus.

ADULT.—A medium-sized brownish-black species with small lateral pale markings on abdomen, not usually produced on to dorsum. Mesonotal scales dark brown, some lighter ones on front margin, and a roundish patch of light brown scales over each wing-root; 5-6 large *ppn* bristles and about 6-7 smaller. At a point on sternopleura opposite lower corner of mesepimeron there are numerous hairs and some scales. Larger part of mesepimeron covered with bristles and fine hairs, conspicuous even in pinned specimens (fig. 67, b) †.

♀.—*Hypopygium* (fig. 72, a) comparatively large. Post-genital plate with two pronounced lobes. Cowl wide and narrow, outer corners bluntly pointed.

♂.—*Hypopygium* (fig. 71, e) : apex of coxite produced into two arms on tergal side, one arm with two strong spines and two more slender, the other divided at the tip. On sternal side there are a few small spines at apex of coxite. Style curved, widened in middle, pointed, and without an obvious appendage, but with hairs along apical $\frac{1}{2}$. Paraproct unusually short.

LARVA.—Unknown.

DISTRIBUTION.—Known only from type-locality.


† [Other species of the subgenus showing numerous fine hairs on lower part of mesepimeron are *A. singularis* Leis., *A. virilis* Leis., *A. unicus* (Theo.) Edw., and *A. leicesteri* Edw., all found in Malaya. Of these, *A. unicus* most closely resembles *A. hirsutipleura* in hypopygial structure, the differences being quite small.]

159. *Aedes* (*Aedes*) *comatus* Barraud, 1931

Ind Journ. Med Res xix, p 615 (♀). TYPE-LOC. Sukna, N Bengal, Darjeeling dist, 500', viii 1928 (*Sobha Ram*) TYPE: ♀ in Brit Mus.

ADULT ♀.—Resembles *A. (A) hirsutipleura* closely in markings and in presence of fairly numerous bristles on lower part of mesepimeron, but with differences in structure of hypopygium (fig 72, b): postgenital plate with pronounced

Fig 72



♀ hypopygia of *Aedes*, subgenus *Aedes*: a, *hirsutipleura*; b, *comatus*; c, *agrestis*; d, *lugubris*; e, *atrius*; f, *sigmoides*; g, *bulleri*. APF, hairy processes anterior to spermathecal eminence. Other lettering as in fig. 70.

lobes. Cowl with wide median lobe and smaller lateral lobes (differing in this respect from all other Indian species). Atrial plates fairly large and rounded

♂ and LARVA.—Unknown

DISTRIBUTION.—Known only from type-locality.

160. Aedes (Aedes) agrestis Barraud, 1931.

Ind. Journ. Med. Res. xix, p. 613 (♀). TYPE-LOC. Nilgiri Hills, S India, ix. 1915 (*Khazan Chand*). TYPE. ♀ in Brit. Mus.

ADULT.—Resembles *A. (A.) hirsutipleura* both in markings and in presence of numerous bristles and hairs on mesepimeron, but with marked differences in hypopygial structures

♀.—*Hypopygium* (fig. 72, c): comparatively large. Cowl only slightly curved. Postgenital plate large and not markedly emarginate on apical border.

♂ and LARVA.—Unknown.

DISTRIBUTION.—NILGIRI HILLS*: type-locality, as given above. BOMBAY DECCAN: Nagargali*, viii. 1921 (*Barraud*)

Not known from elsewhere.

161. Aedes (Aedes) yerburyi Edwards, 1917.

Bull. Ent. Res. vi, p. 222 (♂). TYPE-LOC.: Kith Station, Ceylon, xi. 1891 (*Yerbury*). TYPE. ♂ in Brit. Mus.

[ADULT.—Closely resembles *A. hirsutipleura*, the mesepimeron having numerous hairs on the middle $\frac{1}{2}$ (the type is rather mouldy and denuded and the ornamentation not well shown).

♂.—*Hypopygium* (fig. 73, a): very large, and very different from all other species, sternite apparently with a pair of finger-like processes; coxites swollen, each with a large tergal flap extending the whole length and produced apically into a point, and on the sternal side at tip with a long hook-like process; style rather large, broad, and flattened; paraprocts represented by a pair of very long, stout, sinuous rods reaching beyond end of coxite.

♀ and LARVA.—Unknown.

DISTRIBUTION.—Known only from CEYLON, type-locality, as given above, and Trincomali, xi. 1906 (1 ♂ in British Museum, mounted by Theobald; collector unknown)].

162. Aedes (Aedes) atrius Barraud, 1928.

Ind. Journ. Med. Res. xvi, p. 371 (♀). TYPE-LOC.: Nongpoh, Khasi Hills dist., Assam, vii. 1922 (*Barraud*). TYPE: one co-type ♀ in Brit. Mus.; one in M.S.I. coll., Kaseuli.

ADULT ♀.—*Head* mainly covered with flat dark scales, some pale ones on vertex, along eye-margins, and low down

at each side, fairly numerous upright scales on nape. Mesonotal scales dark brown, with some lighter ones on front margin, 4-5 strong *ppn* bristles, about 18 upper mesepimeral, no lower. Abdomen almost black, with lateral pale markings, produced well on to dorsum on IV-VII, especially on V, those on II and III hardly visible in dorsal view. *Hypopygium* (fig. 72, e): comparatively very large. Cerci broad and rather short. Postgenital plate divided into a pair of large broad lobes. Numerous hairs arising from spermathecal eminence. Atrial plates of distinctive shape.

♂—Unknown, unless the species here described is the ♀ of *A. (A.) yerburyi*, known only from Ceylon.

LARVA.—Unknown.

DISTRIBUTION.—Known only from type-locality.

163 *Aedes (Aedes) lugubris* Barraud, 1928

Ind. Journ. Med. Res. xvi, p. 372 (♀). TYPE-LOC. Rangoon, Burma, v. 1926 (*Hamilton Miller*). TYPE one co-type ♀ in Brit. Mus., one in M.S.I. coll., Kasauli.

ADULT ♀.—A very dark species of moderate size. Head mainly covered with flat black scales, a few pale ones on vertex, a narrow pale patch low down at each side. Integument of mesonotum black, scales deep brown, 5-6 *ppn* bristles, about 12 upper mesepimeral and a few hairs below, scattered over the sclerite, chiefly towards posterior border. A complete, or almost complete, narrow band of white scales on tergite II, curving on to dorsum at sides, and reaching base of tergite in middle line, small lateral pale markings on III-VII. *Hypopygium* (fig. 72, d) postgenital plate only slightly emarginate at apex. Cowl membranous. A pair of hairy processes, apparently connected with the spermathecal eminence, directed anteriorly.

♂ and LARVA.—Unknown.

DISTRIBUTION.—BURMA Rangoon*, type-locality, as given above, also 1930 (*Feegrade*). ANDAMAN Is.: Port Blair*, iii. 1925 (*Shaffi*), and vii. 1926 (*Sobha Ram*).

Not recorded from elsewhere.

164 *Aedes (Aedes) clavatus* Barraud, 1931

Ind. Journ. Med. Res. xix, p. 614 (♂). TYPE-LOC. Sukna, Darjeeling dist., 500', viii. 1928 (*Sobha Ram*). TYPE: ♂ in Brit. Mus.

ADULT ♂.—Medium-sized, wing about 2.8 mm. Head mainly covered with dark brown flat scales, those along eye-margins and at sides light brown. Fairly numerous upright scales on nape. Palpi about $\frac{1}{2}$ length of proboscis. Mesonotum chestnut-brown, with slight reddish tinge, 6 *ppn*

bristles, 10 strong upper mesepimeral, and 6 smaller, arranged in a row extending downwards, and within the patch of scales. Dorsum of abdomen brownish-black, small basal lateral pale patches, not produced on to dorsum. *Hypopygium* (fig. 69, c) a short finger-like process at apex of coxite dorsally, and a pointed process below apex ventrally. Style articulated some distance below apex of coxite on inner side. Paraproct divided into two arms, one longer than the other, both slightly clubbed and truncated.

♀ and LARVA —Unknown.

DISTRIBUTION —Known only from type-locality.

165 *Aëdes* (*Aëdes*) *pseudodiurnus* (Theobald), 1910

Rec. Ind. Mus. iv, p. 32 (*Skusea*) (♂). TYPE-LOC. Sukma, Darjeeling dist., 500', vii. 1908 (*Annandale*). TYPE ♂ in Ind. Mus.

ADULT ♂* —A moderate-sized dark brown species with lateral pale markings on abdomen. Head-scales brownish-black, with some light ones in middle of vertex and at sides. Mesonotal scales dark rich brown. *Hypopygium* (fig. 71, f, drawn from type ♂, which is damaged, coxites being broken); paraproct apparently unusually broad and short.

♀ and LARVA —Unknown.

DISTRIBUTION —Known only from type-locality.

166. *Aëdes* (*Aëdes*) *rami* Barraud, 1928

Ind. Journ. Med. Res. xvi, p. 370 (♂). TYPE-LOC.: Golaghat, Sibsagar dist., Assam, xi. 1924 (*Barraud*). TYPE ♂ in Brit. Mus.

ADULT ♂.—A small dark brown species with small lateral pale markings on abdomen. *Hypopygium* (fig. 71, e): a rather long pointed process, with an accessory point near its tip, projecting from apex of coxite. Two very strong spines, or straight processes, on inner side of coxite towards base. Style enlarged towards tip, a number of hairs on terminal $\frac{1}{2}$ and a pointed terminal appendage. Paraproct single and very long, longer than coxite.

♀ and LARVA —Unknown.

DISTRIBUTION.—Known only from type-locality.

167. *Aëdes* (*Aëdes*) *sigmoides* Barraud, 1928.

Ind. Journ. Med. Res. xvi, p. 373 (♀). TYPE-LOC.: Port Blair, Andaman Is., ix. 1911, caught in crab-hole (*Christophers*). TYPE ♀ in Brit. Mus.

ADULT ♀.—A small blackish species with very small lateral pale markings on abdomen, not produced on to dorsum.

* Barraud 1928, p. 371.

Hypopygium (fig. 72, f) postgenital plate emarginate at apex. Cowl membranous. Two hairy processes, apparently connected with the spermathecal eminence, directed outwardly and somewhat posteriorly.

♂ and LARVA.—Unknown

DISTRIBUTION.—Known only from type-locality.

168 *Aedes* (*Aedes*) *butleri* Theobald, 1901

M.C. II, p. 230 (♀) TYPE-LOC Selangor, Malay Penin., x. 1899
(Butler) TYPE. ♀ in Brit. Mus.

Skusea diurna Theobald, 1903, Entom. xxxvi, p. 259 (♀) TYPE-LOC. Jugra, near Kuala Lumpur, Malay Penin. (Durham)
TYPE ♀ in Brit. Mus.

Stegomyia haitiensis Carter, 1910, Entom. xlin, p. 275 (type ♀ only)
TYPE-LOC: Saigon (or Ha-tien?), Cochinchina (Bouquet)
TYPE: ♀ in Brit. Mus.

ADULT ♀†—A small blackish species with lateral pale markings on the abdomen, not produced on to the dorsum. Thoracic integument black. Mesonotal scales very dark brown. About 12 upper mesepimeral bristles, and below these a number of hairs scattered over the middle part of the sclerite towards posterior margin. *Hypopygium* (fig. 72, g) postgenital lobe comparatively large and not usually emarginate on apical border. Cowl wide and nearly flat, the outer corners rounded. No definite hairy processes connected with spermathecal eminence.

[♂.—*Hypopygium* (fig. 73, e, f). size small, coxites short and rather broad, rounded apically, with an arm arising beyond middle of internal border on sternal side, this arm ending in two small teeth, and with another tooth above its base, style comparatively large, divided almost to base into three arms, one slender and sickle-shaped, one round-tipped, the third stouter, pointing in a different direction, and ending in 2 sharp teeth, paraprocts very small.]

LARVA.—Unknown.

HABITAT.—Brackish pools in mangrove swamps (Leicester).

DISTRIBUTION.—ANDAMAN IS., xi. 1911, caught in forest where biting (Christophers). MALABAR COAST*, x. 1915 (Khazan Chand).

Known also from COCHIN CHINA and MALAY PENINSULA.


† Theobald 1903 a, p. 295 (*Verrallina*), 1907, p. 547 (*Skusea diurna*); Leicester 1908, p. 196 (*Verrallina*). Edwards 1913 b, p. 229 (syn.); 1917, p. 220, Barraud 1928 b, p. 373.

Subgenus *CANCRAÊDES* Edwards, 1929.

Bull. Ent Res. xx, p. 342. Genotype, *Aedes* (*Skusea*) *cancricomes* Edwards.

ADULT—Small dark brown mosquitoes, without special ornamentation. Scales on dorsum of head and on scutellum broad and flat. Proboscis about length of fore femur. Palpi very short in both sexes, about $\frac{1}{4}$ length of proboscis. Plumes

Fig 73



♂ hypopygia of *Aedes*, subgenera *Aedes* and *Cancraêdes*: a, *yerburyi*; b, *cancricomes*; c, d, *simplex*; e, f, *bulleri*; g, *ceylonicus*. Lettering as on p. 4.

of antennæ of ♂ evenly spread round shaft (hairs few and short in *cancricomes*). One or two lower mesepimeral bristles present. All tarsal claws simple in both sexes. Segment VIII of abdomen of ♀ retractile, cerci fairly long, but usually almost hidden within ring of VII. Coxite of ♂ hypopygium short, with group of leaf-like processes and strong spines near apex. Style short and thick, without appendage. Harpago absent. Phallosome divided into lateral plates, with one or two teeth or spines.

LARVA.—As in subgenus *Aedes*.

DISTRIBUTION and BIONOMICS—The subgenus is entirely Oriental. Four species are known, of which two have been found in the Andaman Islands, one of these also occurs in Ceylon and the Nicobar Islands. The larvæ live in crab-holes, or in holes in mangrove swamps. The habits of the adults are not known.

169. *Aedes (Cancraëdes) canericomes* Edwards, 1922.

Ind Journ. Med Res x, p 272 (*Aedes (Skusea)*) (♂ & ♀). TYPE-LOC. Andaman Is, vii 1911, breeding in crab-holes (*Christophers*). TYPE: ♂ & ♀ in Brit Mus.

ADULT †—Resembles *A. (C.) simplex* closely, but differs as follows:—Abdomen with lateral basal creamy-white triangular patches on II–VII, *apn* and *ppm* largely dark, like mesonotum. Antennæ of ♂ longer and much less hairy, not appearing plumose, the hairs not borne on raised ridges; last two segments together about same length as segments 2–12 together.

♂—*Hypopygium* (fig 73, b) differs from *A. (C.) simplex* in form of processes near apex of coxite.

LARVA.—Unknown.

DISTRIBUTION—Known only from type-locality.

170. *Aedes (Cancraëdes) simplex* (Theobald), 1903.

MC m, p 297 (*Ficalbia*) (♂). TYPE-LOC. Kurunogalla, Ceylon (Green). TYPE ♂ in Brit Mus.

ADULT †—A small brownish-black mosquito, distinguished from *A. (C.) canericomes* by absence of pale lateral markings on abdomen, and by the pale pleuræ, including *apn* and *ppm*, contrasting with the dark colour of mesonotum. Distinguished from *A. (R.) longirostris* by shorter proboscis, and by segment 1 of hind tarsi being shorter than tibia.

♂—*Hypopygium* (fig. 73, c, d): in addition to the small thick style, there are several leaf-like and spine-like processes at apex of coxite, differing in form from similar processes in *A. (C.) canericomes*.

LARVA—Unknown.

HABITAT—Crab-holes.

DISTRIBUTION.—CEYLON. type-locality, as given above. ANDAMAN IS *, viii. 1911, breeding in crab-holes (*Christophers*). NICOBAR IS., * iii. 1925 (*Isaac*).

It has been recorded from DUTCH EAST INDIES (*Brug*), but this record probably refers to the closely-allied *A. (C.) curtipes* Edw.


† Barraud 1928 b, p. 362 (*A. (Skusea)*).

† Edwards 1917, p 224 (*Skusea*), 1928, p. 274 (*A. (Skusea)*); Barraud 1928 b, p 362 (*A. (Skusea)*).

Genus **HEIZMANNIA** Ludlow, 1905Can. Ent. xxxvii, p. 130 Genotype, *H. scintillans* Ludl*Bolbodomyia* Theobald, 1910, Rec. Ind. Mus. iv, p. 31. Genotype, *B. complex* Theob.

ADULT *—Small, dark-coloured mosquitoes, distinguished by the presence of a collection of small hairs on the postnotum.

Fig. 74.



♂ hypopygial structures of *Heizmannia*: a-d, style and process on inner surface of coxite; e, coxite from inner side, showing hair-tufts and appendages, also process on inner surface (shown separately). a, *humalayensis*; b, *indica*; c, *chandi*; d, *metalhoa*; e, *covelli*.

* Barraud 1929 b, p. 260.

(The only other Indian mosquitoes having this character are *Tripteroides edwardsi* and *Armigeres (Leucesteria) flavus*, both these species have other well-marked characters distinguishing them from *Heizmannia*) Palpi short in both sexes, antenna in ♂ only slightly plumose, but last two segments usually very long, other flagellar segments short, *apn* lobes large and, in some species, practically touching behind the head* Mesonotum covered with rather broad recumbent scales, usually with metallic lustre, no bristles on dorsum, even immediately in front of scutellum. Usually 3 or 4 *ppn* bristles, no spiracular, postspiracular or upper sternopleural, 1 or 2 lower and several upper mesepimeral.

♂—*Hypopygium*: coxite comparatively short, wide, and rounded, with subapical lobe carrying 1 or 2 stout spines. Usually several hair-tufts on inner side of coxite towards base, which, in some species, are remarkably long and twisted or matted. Arising from inner side of coxite, towards sternal side, there is also a peculiar process, apparently representing the "harpago" or "claspette", the apical part, or blade, of this structure is very transparent, and sometimes divided into leaflets, the form of which cannot usually be clearly seen except in heavily stained preparations. Style short, widened apically, appendage stout and pointed. Phallosome divided into small lateral plates with teeth. Paraproct ending in single blunt tooth. In two species the 9th sternite is very large.

LARVA.—Very similar to *Aedes*. The larvae of four species are known, and descriptions of three of these are given below.

DISTRIBUTION and BIONOMICS.—About twelve species are at present known, these are all Oriental except for one which occurs on the island of Halmahera. The larvae live in tree-holes and bamboo-stumps. Nothing is known regarding the habits of the adults.

Key to Adults.

♀♀.

- | | |
|---|-------------------------------|
| 1 Proboscis more than $1\frac{1}{2}$ times length of fore femur | <i>indica</i> , p. 303. |
| Proboscis only slightly, if at all, longer than fore femur | 2 |
| 2 A large patch of silvery scales on mid-lobe of scutellum, extending anteriorly on to mesonotum, hind femur entirely pale on basal $\frac{1}{2}$ | <i>himalayensis</i> , p. 306. |
| Only a small patch of silvery scales on mid-lobe of scutellum, or none, hind femur dark along dorsal edge, from base to knee-joint | 3 |

* [The lobes themselves are not much larger than usual, but appear so owing to their very dense covering of large flat scales.]

- 3 *apn* entirely dark scaled *funcrea*, p. 302.
apn with some white or silvery-white scales 4
- 4 Mesonotal scales dull greyish-brown, with little or no metallic lustre, *apn* large and touching behind head, with white scales anteriorly *complex*, p. 302.
 Mesonotal scales with strong blue, green, or bronzy metallic lustre, *apn* more or less separated 5.
- 5 *apn* covered with silvery-white scales, mesonotal scales with bright green or bluish-green lustre 6
apn with white scales in front only, mesonotal scales with dark blue or bronzy lustre. 7
- 6 Outstanding plume-scales on veins 2.1 and 3 2 linear *chandi*, p. 308.
 These scales rather narrow, but ligulate in shape *viridis*, p. 306
- 7 Proboscis shorter than fore femur, mesonotal scales blackish, with bronzy lustre. Proboscis as long as fore femur or slightly longer; mesonotal scales with deep blue lustre 8.
 8. Outstanding plume-scales on veins 2.1 and 2 2 linear *giceni*, p. 308.
 These scales ligulate in shape *metallica*, p. 307.

♂♂.

- 1 *apn* lobes entirely dark scaled *funcrea*, p. 302
apn with some white scales 2.
- 2 Ninth sternite very large, much more than $\frac{1}{2}$ length of coxite 3.
 Ninth sternite not more than $\frac{1}{2}$ length of coxite 4
- 3 Proboscis longer than fore femur *indica*, p. 304.
 Proboscis shorter than fore femur *covelli*, p. 305.
- 4 Style not markedly expanded apically, and divided nearly to the base into two arms. *complex*, p. 302
 Style expanded apically, and not divided into two arms 5.
- 5 Subapical lobe of coxite with 2 strong spines. Subapical lobe of coxite with only 1 strong spine *giceni*, p. 308.
 6.
- 6 Style greatly expanded apically; coxite with tufts of very long hairs from base; subapical lobe produced into an arm directed basally, with a single strong spine at extremity *himalayensis*, p. 307.
 Style less expanded; coxite with tufts of rather short hairs all about same length; subapical lobe not produced into a long arm 7.
- 7 Coxite with a large dense tuft of rather short twisted hairs near base; style with an elbow-like projection near base *chandi*, p. 308.
 Coxite without any large dense tufts of hairs; style without an elbow-like projection near its base *metallica*, p. 308.

[Key to known Larvæ

- | | |
|---|-----------------------------|
| 1 Antenna entirely pale, shaft smooth, terminal papilla short | <i>himalayensis</i> , p 307 |
| Antenna longer, darkened on distal $\frac{1}{2}$, shaft with some fine spicules, terminal papilla long | 2 |
| 2 Pecten-teeth with long basal denticles | <i>covelli</i> (?), p. 305 |
| Pecten-teeth all simple | 3 |
| 3. Siphon under 3×1 | <i>funerea</i> (see below). |
| Siphon about 4×1 (Sumatran species) | <i>pilosa</i>] |

171. *Heizmannia complex* (Theobald), 1910.

Rec. Ind Mus iv, p 31 (*Bolbodeomyia*) (♂ & ♀) TYPE-LOC -
Dawna Hills, Lower Burma, iii 1908 (*Anmandale*) TYPE
♂ & ♀ in Ind Mus

ADULT ♀†.—General coloration dark brown. Proboscis about length of fore femur Mesonotal and scutellar scales dull greyish-brown, with little or no metallic lustre, *apn* large and practically touching behind head, with white scales anteriorly Plume-scales on veins 2 1 and 2 2 narrow and linear. Abdomen deep brown, with small lateral pale markings, rather square in shape on segments 2 and 3, small and triangular on remaining segments, and not forming bands over dorsum

♂—Very similar to ♀ Antennæ of usual form in this genus *Hypopygium*. style of characteristic form, being divided nearly to base into two separate arms Coxite rather square-ended, with a few flattened bristles at tip, no hair-tufts, but a short subapical arm bearing one long spine, *lp* more truncate apically than in the other species.

LARVA —Unknown

DISTRIBUTION —BURMA type-locality*, as given above
ASSAM Nongpoh*, Khasi Hills dist, vii 1922 and iii 1927 (*Barraud*)

Recorded from COCHIN CHINA (*Borel*)

172. *Heizmannia funerea* (Leicester), 1908.

Oul Malaya, p 252 (*Wyeomyia*) (♀). TYPE-LOC : Kuala Lumpur,
Malay Pennin (*Leicester*) TYPE ♀ in Brit. Mus

ADULT ‡.—Distinguished from all other known Indian species by characters given in keys. ♂ hypopygium very similar to that of *H indica*, but with slight differences in form of style.

LARVA § —This has been described by Edwards and Given as follows:—"Head rounded, scarcely as broad as long,

† Edwards 1922 d, p 447; Barraud 1929 b, p. 263

‡ Edwards 1922 d, p. 448; Barraud 1929 b, p. 263

§ Edwards & Given 1928, p 340.

pale in colour, except round occipital margin and mouth. Clypeal spines long and slender, not set on distinct tubercles: a small accessory hair external to each. Dorsal hairs placed well back from the front. *A* with about 15 plumose branches; *B* with 2-3 branches, placed in front of but a little external to *C*, which has about 10 branches, *d* smaller, but still rather large, with about 20 plumose branches arranged in a fan-shaped manner round a pear-shaped base, *e* small, 2-3-branched. Antennæ two-thirds as long as the head, somewhat curved and tapering, darkened on the outer half, shaft with a few very fine spicules, tuft of 4 branches placed about the middle, preapical bristles close to tip, one much longer than the other. Mouth-bristles normal, the constituent hairs simple. Mentum triangular, a rather large central tooth, on each side of which are first five closely set teeth, and then five more, shorter and more widely separated. *Thorax* with 3-4 almost simple hairs in the lateral tufts, no associated spines, and no definite plates. *Abdomen* with the lateral hairs mostly double, dorsal hairs inconspicuous. Comb a patch of about 15 scales, most of which are finely and evenly fringed all round, though a few have a long terminal point. *Anal segment* with a moderately large saddle, the posterior margin of which is fringed with little clumps of very fine setæ. Outer dorsal hair long and simple, inner dorsal hair double. Lateral hair rather long and simple. A small barred area present, with six long, double, non-plumose hairs, two small single hairs before the barred area. Siphon very dark, broadest a little before the middle, then rather strongly tapering; index about 3. No acus. Pecten composed of 5-9 simple teeth, reaching hardly more than one-third the length of the siphon, tuft represented by a single long hair placed a short distance beyond the last pecten-tooth. Valves small, of normal structure."

HABITAT.—Tree-holes.

DISTRIBUTION.—As regards India, this species is known from one ♀ only, from Meenglas, Jalpaiguri, N BENGAL (*M O T. Iyengar*).

Known from MALAY PENINSULA (Kuala Lumpur and Singapore).

173. *Heizmannia indica* (Theobald), 1905.

Ann. Mus. Nat. Hung. lii, p. 115 (*Phonomysia*) (♂). TYPE-LOC.: Singapore, 1902 (*Biró*). TYPE: ♂ in Nat. Mus. Hung., Budapest.

ADULT ♀*.—*Head*: mainly covered with purplish-black

* Theobald 1907, p. 601; Leicester 1908, p. 253; Edwards 1922 *d*, p. 447; Barraud 1929 *b*, p. 265.

flat scales, a small silvery area on vertex, continued downwards between eyes. Antennæ, clypeus, palpi, and proboscis black. Proboscis unusually long, from $1\frac{1}{4}$ – $1\frac{1}{2}$ times length of fore femur. *Thorax*: mesonotum with scutellum covered with greenish-blue broad scales with metallic lustre, *apn* more or less separated, with white scales on upper and anterior parts, dark scales below. Patches of silvery-white scales on pleuræ and coxæ. *Wings* dark scaled, outstanding plume-scales on veins 2.1 and 2.2 rather narrow, but ligulate in shape. *Legs*: dark brown, mid-femur paler beneath, hind femur white along both surfaces from base nearly to knee, except along dorsal edge, which is dark for the whole length. *Abdomen* bluish-black, with lateral triangular basal white markings not usually produced on to dorsum to form basal bands, as is the case in some other species.

♂.—Proboscis unusually long and palpi quite short, as in ♀, last two flagellar segments of antennæ very long, as usual, rest short (about 2×1)*. *Hypopygium*: 9th sternite very large and hood-like, as long as coxite, with a few scales and short hairs on its distal portion. Form of style and "harpage" shown in fig. 74, b.

LARVA.—Undescribed

HABITAT.—Tree-holes and bamboo-stumps

DISTRIBUTION.—Fairly common in EASTERN HIMALAYAS from 500' (Sukna) up to 6–7,000' (Kurseong, Darjeeling, etc.). Not known at present from other parts of Indian area, but is found in MALAY PENINSULA

174 *Heizmannia covelli* Barraud, 1929

Ind. Journ. Med. Res xvii, p 265 (♂ & ♀) TYPE-LOC Andaman Is, vii 1926 (♂) (*Covell*), Sukna, N Bengal, x 1922 (allotype ♀) (*Barraud*) TYPE co-type ♂ and allotype ♀ in Brit Mus

ADULT.—Differs from *H. chandi*, *greeni*, and *metallica*, to which it appears to be related, in structure of ♂ hypopygium, and in following characters present in both sexes:—Proboscis distinctly shorter than fore femur, mesonotal scales blackish, with a bronzy metallic sheen, scutellar scales

* [In the key to the species of *Heizmannia* published by Edwards (1922 d) the antennæ of *H. indica* were stated to be alike in the two sexes, i. e., with all the flagellar segments in ♂ subequal in length, the last two not elongate. This is true of the Malayan ♂ in the British Museum, assumed to be conspecific with the type from Singapore, of which the hypopygium was figured by Edwards. This Malayan ♂ may be abnormal as regards its antennæ, as it is identical in hypopygial structure with one in the British Museum from Kurseong (*M. O T. Iyengar*), in which the antennæ are as described by Barraud.]

greenish-black, with a few silvery scales at apex of mid-lobe, *apn* approximated, but not touching behind head, with silvery-white scales anteriorly, outstanding plumescales on veins 21 and 22 narrow but hgulate in shape. Abdomen marked with the usual lateral triangular basal silvery patches, with narrow white basal bands over dorsum on VI and VII in ♀. 3-4 *ppn* bristles, 1-2 lower mesepimeral. Last two flagellar segments of antenna of ♂ very long, as usual, terminal segment about twice length of penultimate, last two segments together a little more than twice length of first 11. segments 2-11 each only about twice as long as broad. Tarsal claws of fore and mid-legs of ♂ with a slender sharp tooth arising from base.

♂—*Hypopygium* (fig. 74, e) coxite short and rounded, with two large tufts of hairs arising from inner side towards base, subapical lobe with a single stout pointed process. Style less expanded apically than in several other species. Harpago of characteristic shape.

LARVA.—Not known with certainty. Some specimens sent from Rangoon, thought to be this species, show the following characters:—Length about 5 mm. *Antennae* comparatively long, darkened on apical $\frac{1}{2}$, shaft rather more than $\frac{1}{2}$ length of head, with small, thin spines scattered along its length, hair-tuft of about 4 branches at middle of shaft, 2 subapical bristles a little below tip, terminal papilla very long. *Frontal hairs*: *A* missing, *B* with few fairly long branches, *C* and *d* with fairly numerous fine branches. Hair *C* has some distance behind *B*, *d* internal to, and almost level with, *B*. Preclypeal spines fine and long. Lateral hairs of *thorax* moderately well developed, meso- and metapleural tubercles with rather small, but distinct, thorn-like spines. Lateral hairs of *abdomen* with 2-3 long branches. Comb of about 12 fringed teeth, each ending in a number of fine points, teeth apparently arranged more or less in two rows. *Siphon* not quite 1 mm. long, tapering from middle to apex. Pecten of 5-8 moderate-sized pointed teeth, each with several long, but fine, denticles, which are divided from main tooth nearly to base. Hair-tuft of 3-5 fine branches, base of tuft before middle of tube. *Anal segment* with fairly large chitinous saddle, long, single, lateral hair and small fan-plate, *asc* of 3-4 long branches; *osc* single and long, as usual. Fan of few hairs.

HABITAT.—Tree-holes and bamboo-stumps.

DISTRIBUTION.—ANDAMAN Is. *, and N. BENGAL, Sukna *, type-localities, as given above. BURMA: Rangoon *, 1930 (*Feegrade*).

Not known from elsewhere.

175 *Helzmannia viridis* Barraud, 1929.

Ind Journ Med Res xvii, p 266 (♀) TYPE-LOC N Kanara,
Yellapur, x. 1921 (*Barraud*) TYPES co-type ♀ in Brit Mus.,
two co-types in M.S.I. coll., Kasauli

ADULT.—Most nearly resembles *H. covellii* and *H. metallica*, but differs as indicated in key Wing about 3 mm

♀—*Head* a median patch of flat silvery scales on vertex in front, continued downwards between eyes, remainder of dorsal surface covered with flat blue-black scales, a patch of silvery scales low down at each side. Antennae, olypens, palpi, and proboscis black, the last about equal in length to fore femur. *Thorax*. mesonotal scales with bright green metallic lustre, contrasting with blue-black scales on head and abdomen. Scutellar scales flat, with green or greenish-blue metallic lustre, a few white scales in middle of centro lobe, *apn* large and fairly closely approximated, covered with silvery-white scales, except on lower part, where scales are black. A well-marked tuft of hairs on postnotum. Integument of pleurae black, with patches of silvery-white scales; 3-4 *ppn* bristles and 1 lower mesepimeral. *Wings* dark scaled, outstanding plume-scales on veins 2.1 and 2.2 rather narrow, but ligulate in shape. *Legs* deep bronzy-brown, fore and mid-femora paler posteriorly, hind femora more extensively pale on both surfaces, dark along dorsal edge and around knee-joint. *Abdomen* blue-black, with the usual basal triangular white markings laterally, and more or less complete narrow basal bands over dorsum on terminal segments

♂ and LARVA—Unknown

DISTRIBUTION.—NORTH KANARA Yellapur*, typo-locality, as given above. BOMBAY DECCAN Nagargah*, viii 1921 (*Barraud*) MALABAR COAST*, x 1915 (*Khazon Chand*).

Not known from elsewhere

176. *Helzmannia himalayensis* Edwards, 1922.

Ind Journ Med Res x, p 290 (♂) TYPE-LOC Darjeeling,
vii 1918 (*Kemp*) TYPE: ♂ in Ind. Mus

ADULT †.—Distinguished from other species on following characters:—Hind femur entirely pale on basal $\frac{1}{2}$, even along dorsal edge. Mesonotal scales black, with bluish-black metallic lustre. A large silvery patch covering mid-lobe of scutellum and extending on to mesonotum in middle. Proboscis about length of fore femur, *apn* slightly separated, with silvery-white scales covering the larger part. Outstanding plume-scales on veins 2.1 and 2.2 rather narrow but ligulate. Abdomen with basal triangular silvery patches

† Barraud 1929 b, p. 267

laterally, and narrow basal white bands over dorsum on segments IV-V in ♀. Larger claw of fore tarsi of ♂ with a long slender tooth.

♂—*Hypopygium* (fig 74, a) style characteristic, with very large expansion on apical $\frac{1}{2}$. Harpago with hairy stem, and processes resembling petals of a flower.

LARVA.—The following details are described from one incomplete larval skin from Koti, near Kalka, from which the resulting ♀ has been preserved.—*Antenna* pale except at extreme base, of moderate length, considerably shorter than in *covelli* and *funerea*, tip about level with ends of longer mouth-brush hairs in the mounted skin. shaft quite smooth, hair-tuft of 2 branches, nearer apex than base, terminal papilla quite small. *Frontal hair* *A* of about 10 branches, *B* and *C* each of 3 fairly long branches, *C* standing directly behind, but some distance from, *B*. Mouth-brushes of moderate size, hairs simple. Preoclypeal spines pale in colour, very long and tapering. Mentum triangular, with 11–12 teeth either side of central one. Hairs on *thorax* and *abdomen* moderately developed. Comb of a few large sharp teeth, each tooth with lateral fine hairs towards base. *Siphon* about 0.8 mm long. a dark ring at base, no acus. Pecten of only 4–5 fairly large sharp teeth, each with 2–3 large basal lateral denticles. Hair-tuft of 3 branches, slightly nearer base than apex. *Anal segment* almost enclosed in chitinous ring, *lh* of 2 rather long branches, *isc* of 2 branches.

HABITAT.—Tree-holes, and bamboo.

DISTRIBUTION.—EASTERN HIMALAYAS. Darjeeling*, type-locality, as given above, Kurseong*, 6,000' Darjeeling dist., viii 1928 (*Sobha Ram*). NORTH BENGAL. Sukna*, Darjeeling dist., 500', ix. 1922 (*Barraud*), and viii 1928 (*Sobha Ram*). WESTERN HIMALAYAS. Koti*, near Kalka (Kalka-Simla road), c 4,000', vii. 1923 (*Barraud*).

Not known from elsewhere.

177. *Heizmannia metallica* (Leicester), 1908.

Cul Malaya, p. 251 (*Wyeomyia*) (♀). TYPE-LOC Bukit Kutu, Malay Penin. (Leicester). TYPE. ♀ in Brit. Mus.

ADULT†.—Very similar to *H. covelli* and *viridis*, but the mesothoracic scales have a deep bluish-metallic lustre (not green or bronzy). Proboscis slightly longer than fore femur; *apn* large and nearly touching behind head, with silvery-white scales on outer anterior parts. Outstanding plume-scales on veins 2.1 and 2.2 ligulate. Abdomen with the usual large basal triangular silvery patches. Antenna of ♂ with first 11 flagellar segments very short, scarcely longer than

† Edwards 1922 d, p. 448; Barraud 1929 b, p. 267

broad, each with few long hairs, last two segments with long and dense pubescence and very long basal hair-whorls

♂—*Hypopygium* (fig 74, d) coxite without any large dense tufts of hairs, subapical lobe with a single strong spine-like process. Harpago represented by two processes, one of which is very transparent, with broad stem and leaf-like blade

LARVA—Unknown

HABITAT—Tree-holes

DISTRIBUTION—MALABAR COAST Pudupadi*, x. 1915 (*Khazan Chand*) EASTERN HIMALAYAS Kurseong, Darjeeling dist., 5,000', ix 1909 (*Annandale*)

Known also from MALAY PENINSULA

178 *Heizmannia chandi* Edwards, 1922

Ind Journ Med Res x, p 291 TYPE-LOC Pudupadi, Malabar Coast, x 1915 (*Khazan Chand*) TYPE ♂ and allotype ♀ in Brit Mus

ADULT †—Resembles *H. greeni* and *complexa* in having scales on veins 2.1 and 2.2 very narrow (linear), but differs in other respects as indicated in key. Also closely resembles *H. viridis*, differing in the narrower wing-scales.

♂—*Hypopygium* (fig 74, e) coxite with a collection of rather short hairs, all about same length, on inner side, harpago with several leaf-like expansions representing the blade, style with an elbow-like projection near base on outer side

LARVA—Unknown

DISTRIBUTION—Known only from type-locality

179 *Heizmannia greeni* (Theobald), 1905.

Journ Bomb Nat Hist Soc xvi, p. 247 (*Hypomyia*) (♂ & ♀) TYPE-LOC. Peradeniya, Ceylon, i & ii. 1902 (*Green*). TYPE ♂ & ♀ in Brit Mus

ADULT †—Proboscis about length of fore femur, mesonotal scales with dark blue-metallic lustre, *apn* with white scales only along anterior edge and laterally: outstanding plume-scales on veins 2.1 and 2.2 very narrow (linear); basal white, triangular, lateral patches on abdomen, not forming bands over dorsum

♂—*Hypopygium* subapical lobe of coxite with two strong spine-like processes, long twisted hair-tufts on inner side of coxite

LARVA.—Unknown.

DISTRIBUTION—MALABAR COAST*, x. 1915 (*Khazan Chand*). CEYLON type-locality, as given above.

† Barraud 1929 b, p 268.

† Theobald 1907, p 597, Edwards 1922 d, p 449, Barraud 1929 b, p 268.

Genus *HÆMAGOGUS* Williston, 1896

Trans. Ent. Soc. Lond. 1896, p. 271. (Genotype, *H. splendens* Will.)

Two Indian mosquitoes have been provisionally referred to this tropical American genus, but, as no males of the Indian species have yet been obtained, their true generic position is not definitely settled*. In general appearance they resemble the species of *Heizmannia*, but there are no hairs on the postnotum.

ADULT ♀.—Densely scaly mosquitoes, general coloration black, with brilliant silvery markings, more definitely metallic than in *Heizmannia*. *apn* large, collar-like, closely approximated behind head. Proboscis slender, distinctly longer than fore femur, palpi not more than $\frac{1}{3}$ length of proboscis. Larger part of dorsum of thorax without bristles, none present even immediately in front of scutellum.

♂ and LARVA.—Unknown.

DISTRIBUTION and BIONOMICS.—Both Indian species appear to be rare and confined to heavily forested localities. The ♀♀ are day-flying, and most probably suck human blood.

180 *Hæmagogus discrepans* Edwards, 1922

Ind. Journ. Med. Res. x, p. 201 (♀). TYPE-LOC. Pudukudi, Malabar Coast, N. 1915 (*Khazan Chand*). TYPE ♀ in Brit. Mus.

ADULT †.—A medium-sized dark mosquito with silvery markings. Wing 3.2–3.5 mm.

♀.—*Head*. a large patch of silvery flat scales, with bluish tint, on vertex, a silvery patch at each side, remainder of head covered with flat black scales, except for very few upright scales far back on nape. Antennæ, clypeus, palpi, and proboscis black. Palpi about $\frac{1}{3}$ length of proboscis. *Thorax*: integument of mesonotum black, covered with deep bronzy scales, for the most part fairly narrow, but broader ones posteriorly, a few bristles on anterior margin and over wing-roots. Mid-lobe of scutellum densely clothed with flat silvery scales, some black scales apically; lateral lobes with dark scales; *apn* large, closely approximated behind head, covered outwardly with flat silvery scales; *ppn* bare. A patch of flat silvery scales below margin of mesonotum, in front of wing-root, another patch covering larger part of sternopleura and upper part of mesepimeron; other patches on propleura and coxæ; 3 or 4 *ppn* bristles or

* [A third Oriental species (*achetæ* Leir.) is also referred here, and is likewise known only in the female.]

† Barraud 1929 b, p. 269.

2 postspiracular, 1 or 2 lower mesepimeral and 3 or 4 upper mesepimeral. Integument of postnotum and pleura almost black. *Wings* dark scaled, outstanding plume-scales rather narrow, membrane with bronzy-green or blue reflections. *Legs* brownish-black, mid-femur pale posteriorly on about basal $\frac{1}{2}$, hind femur pale on both sides, except dorsally, on about basal $\frac{1}{2}$. Tarsal claws of fore and mid-legs toothed. *Abdomen* greenish-black, with large basal lateral silvery patches, usually produced on to dorsum of segments V-VII, and forming narrow basal bands. Segment VIII and cerci small, tip of abdomen square-ended in side view.

♂ and LARVA — Unknown

HABITAT — Bamboos (*Barraud*)

DISTRIBUTION — MALABAR COAST. type-locality*, as given above BOMBAY DECCAN. Nagargali*, viii 1921 (*Barraud*). N KANARA Yellapur*, x 1921 (*Barraud*)

Not known from elsewhere

181 *Hæmagogus tripunctatus* (Theobald), 1908.

Rec Ind Mus ii, p 288 (*Stegomyia*) (♀). TYPE-LOC. Assam, Lushai Hills, vi 1904 (*Macleod*) TYPE. ♀ in Ind. Mus

ADULT ♀ † — Very similar to *H. discrepans*, described above, but differing as follows — Hind femur pale on anterior, or outer, side nearly to the knee, tarsal claws simple; outstanding wing-scales rather broad

♂ and LARVA — Unknown

DISTRIBUTION — Known only from the type-locality.

Genus *ARMIGERES* Theobald, 1901.

MC i, p 322 Genotype, *A. obturbans* Walk

(For synonymy see under subgenera)

ADULT ‡. — Medium-sized to rather large mosquitoes, not usually highly ornamented, but otherwise resembling *Aedes* (subgenus *Stegomyia*) in general appearance and structure. Scales of head broad and flat, a few upright scales on nape, at least in some species, scutellar scales all broad and flat. Proboscis rather stout, slightly curved downwards towards tip, the curved part somewhat laterally flattened. Palpi of ♂ as long as, or longer than, proboscis, of 3 segments, the last two of which are long, slender, turned upwards, and without hair-tufts. Palpi of ♀ of variable length, from about $\frac{1}{2}$ to $\frac{3}{4}$ length of proboscis. Antennæ of ♂ plumose, the plume-hairs projecting fairly evenly round shaft. Abdomen of ♀ somewhat tapering, segment VIII partially retractile, cerci short.

† Theobald 1910 b, p. 182; Barraud 1929 b, p. 269.


‡ Barraud 1927 b, p. 534; Edwards 1932, p. 182.

Coxite of ♂ hypopygium with lobe on inner surface bearing spines. Style of variable form, but usually with spines along apical part or at tip. Phallosome divided into small lateral plates with teeth. Paraprocts without teeth or hairs at crown. The only character distinguishing the genus, as a whole, from *Aedes* appears to be the form of the proboscis.

PUPA —As in *Aedes*

LARVA —Resembles *Aedes* except in absence of a pecten on siphon. *A. flavus* is distinct from all other known Culicine

Fig 75




Adult characters of *Armigeres*: a, *omisus*, ♀, side view (legs and wings omitted), with enlargement of postspiracular area, showing scales, b, *obturbans*, ♀, as in a; c, *obturbans*, ♂, head

larvæ in having a conspicuous ventral chitinated plate on the anal segment separate from the dorsal saddle. Descriptions of this, and of the five other known larvæ, are given in the following pages. Of these five the author has seen only a few specimens of three, and a reliable key cannot be given at present. [In all known larvæ of the genus except the aberrant *A. treubi* the antennæ are short, with smooth shaft, frontal hairs rather small and placed far forwards; mandibles large, with strong black teeth, comb-scales in a patch or very

irregular row, siphon very short and with trachea with siphonal tube very wide, anal fan poorly developed, no ventral tufts on anal segment before the fan, papillae very large and sausage-shaped.]

DISTRIBUTION and BIONOMICS.—The genus is mainly Oriental, a few species extend into the Australian region, and one as far as Japan. Of about 30 known species, 13 have,

Fig. 76.



Larval structures of *Armigeres* a-d, *obturbans* (head, tail-end, mentum, comb-scales), e-g, *flavus* (anal segment, comb-scales, mentum).

so far, been found in India. These occur chiefly in forested localities where there is a heavy rainfall, and the larva live chiefly in bamboos, coconut-shells, etc. Those of *A. obturbans* sometimes occur in tree-holes, cement sinks in connection with rain-water channels, and in very foul domestic collections of water. The eggs are usually laid singly, but *A. flavus* has been observed on more than one occasion with a mass of eggs attached to its hind leg, with the object, it is thought,

of introducing those to water in bamboo-stems, through a small hole. The ♀♀ are vicious blood-suckers, and attack man both in the day and at dusk.

Subgenus ARMIGERES Theobald, 1901.

M C i, p. 322. Genotype, *A. obturbans* Walk.

Desvoidya Blanchard, 1901, C R Soc Biol lin, p. 1046, nom. nov. for *Armigeres*.

Blanchardiomyia Brunetti, 1912, Rec Ind Mus iv, p. 440, nom. nov. for *Desvoidya*.

Desvoidia Theobald, 1903, M C iii, p. 134, lapsus for *Desvoidya*.

The following characters distinguish this subgenus from *Lecosternia*—Postspiracular area with several bristles, and some white scales on this part posteriorly, but no black scales, mesonotum not distinctly produced over the head, palpi of ♀ less than $\frac{1}{2}$ length of proboscis. One lower mesepimeral bristle usually present (absent in *Lecosternia* except in *flavus*). Style of ♂ hypopygium with close-set spines along about $\frac{1}{2}$ its length.

Key to Adults *

1. Abdominal tergites with median apical yellow markings *theobaldi*, p. 319.
- Abdominal tergites without such markings 2
2. Mesonotum with a pair of well-defined submedian golden lines, and a curved line of similar scales over each wing-root, lateral white markings on abdominal tergites produced on to dorsum *auricolmatus*, p. 319.
- Mesonotum without a pair of submedian golden lines (though there may be a single median yellowish line), lateral white markings on abdominal tergites not produced on to dorsum 3
3. Abdominal sternites III-VI with wide apical black bands, pale margin of mesonotum usually yellowish; palpi of ♀ about $\frac{1}{2}$ length of proboscis; basal lobes of coxite of ♂ usually with 2 straight spines which are round and not flattened, tip of style not reaching to basal lobe. *obturbans*, p. 314.
- Abdominal sternites III-VI variously marked (see description of varieties below), scales on margin of mesonotum usually white (or else no distinct pale border present); palpi of ♀ about $\frac{1}{2}$ length of proboscis; basal lobe of coxite of ♂ with 3 curved flattened spines; tip of style reaching to basal lobe *kuchingensis*, p. 317.

* [In addition to the four species already recorded as occurring in India, it is probable that some others occur in Burma, as about ten have been found in the Malay Peninsula and Archipelago. Of these *A. malay* is one of the commonest and most widely distributed; it is distinguished from *obturbans* by hypopygial structure and by presence of white scales on the clypeus.]

182 *Armigeres (Armigeres) obturbans* (Walk.), 1860*

Proc Linn Soc Lond iv, p 91 (*Culex*) (♀) TYPE-LOC

Amboyna (Moluccas) (Wallace) TYPE ♀ in Brit Mus

Culex ventralis Walker, 1861, Proc Linn Soc Lond v, p 144

(♀) TYPE-LOC Celebes (Wallace) TYPE ♀ in Brit Mus

Culex subalbatus Coquillett, 1898, Proc U S Nat Mus xxi, p 302

(*Culex*) (♀) TYPE-LOC Japan TYPE U.S. Nat. Mus

Armigeres panalectoros Giles, 1902, Handbook, 2nd ed p 386

(♂ & ♀) TYPE-LOC Calcutta (Giles) TYPE ♂ & ♀ in Ind Mus

ADULT † (fig. 75) — Rather large species Wing 4–5 mm


♀ — *Head* a border of flat creamy or white scales to eyes, narrow in middle, widening out at sides; a patch of broad white scales in middle of vertex, remainder covered with bluish-black flat scales, some upright scales on nape. Tori dark brown or black, covered with small broad pale scales, flagellum and hairs of antenna brown. Clypeus black, with greyish sheen. Palpi and proboscis brownish-black, palpi about $\frac{1}{3}$ length of proboscis. *Thorax*: a border of creamy or yellow scales to mesonotum from wing-roots round front margin, dorsum covered with dark brown narrow scales, with, in many specimens, lighter scales intermixed, sometimes forming a fairly well-defined median line from front margin to scutellum, and curved lines laterally. Scutellar scales broad and flat, varying in colour from black to pale brown. Postnotum dark brown, *apn* with flat white scales

* I have assumed that the species which is generally common in India is the same as the one described by Walker as *C. obturbans* and *C. ventralis*, from Amboyna and Celebes, but there would seem to be some doubt on the point. There is at least one other species (*A. kuchingensis*) which is variable, and in some forms resembles *obturbans* closely. The latter is far less common in Upper Assam. Specimens sent to me as *A. obturbans*, from the Philippine Islands, proved to be *A. kuchingensis*, and I am in doubt as to whether *Desvoidia fusca* var. *joloensis*, described by Ludlow from those islands (1904 a, p 236), should be given as a synonym of *A. obturbans* or not. [If it should be shown that the species here called *obturbans* does not occur in Amboyna or Celebes, it should, perhaps, take the name *subalbatus* Coq., the application of which is not in doubt, as all specimens from China and Japan appear to be *A. obturbans* as here described, having the same hypopygial structure and unequal middle claws of ♂. On the other hand, it is possible that the name *joloensis* should replace *kuchingensis*, as this appears to be a common Philippine form. The types of *joloensis* were from the island of Jolo, south of Mindanao, P I., and are in the U.S. Nat. Mus. Specimens in the British Museum received from Miss Ludlow as *joloensis*, and agreeing with the descriptions, are labelled as from Ilo-Ilo, Panay, P I., these have the ♂ hypopygium as in *kuchingensis*, and colouring almost as in var. *durhami*, sternites II and VI being almost all white, III–V with broad black apical bands.]

† Leicester 1908, p 75, Edwards 1917, p. 205; 1921 c, p. 320; 1924, p 368, Brug 1924 b, p 46. Barraud 1927 b, p 537, Martini 1930, p 257

anteriorly, dark scales posteriorly, *ppn* with narrow scales on upper part, usually white or yellowish, but in some specimens dark brown, broader white scales on lower part, and patches of similar scales on pleuræ and coxæ. A few bristles and a small patch of white scales a short distance behind

Fig 77



♂ hypopygial structures of *Armigeres*, subgenus *Armigeres*. a, *obtusatus* (with details), b, *kuchingensis* (with style in different positions), c, *aureolineatus* (style and basal lobe). Lettering as on p. 4

anterior spiracle. *Wings*: dark scaled. *Legs*: brownish-black, hind femur white on outer side from base to knee-joint, usually a small pale knee-spot dorsally on mid-femur; all legs pale when seen from behind; tibiae and tarsi not marked with pale rings. *Abdomen*: brownish-black dorsally, lateral white markings on tergites, hardly, if at all, visible dorsally. Sternite II white scaled, III-VI white at base, with black

apical bands, usually broad and all about same width VII dark at base, with narrow subapical white band, tergite VIII usually pale

♂.—Palpi longer than proboscis by about $\frac{1}{2}$ length of apical segment. Antennæ plumose. Markings as in ♀. Claws of middle leg rather markedly unequal (though larger claw is much smaller than those of front leg) *Hypopygium* (fig 77, a) *bl* with 2 (occasionally 3) fairly strong, straight spines, pointing inwards towards opposite coxite. Style slightly curved, with about 18 short teeth along rather more than apical $\frac{1}{2}$, tip does not reach basal lobe when style is depressed against coxite

LARVA * (fig 76, a-d) —Described from several skins definitely identified as this species by examination of hypopygium of resulting ♂♂. *Antenna* short and thick, about 6-7 times length of greatest width, shaft bare except for one very small hair at about middle, sometimes a little nearer base than apex. *Frontal hair* *A* usually of 3 fairly long branches, *B*, *C*, and *d* placed far forward, *d* being most anterior and slightly external and dorsal to preclypeal spines, *B* external and slightly posterior to *d*, *C* slightly internal and posterior to *B*, *d* fairly large and usually with 6 fine branches. *B* single and long, fairly stout on basal part, fine apically, *C* usually of 3 fairly long branches, *e* fine and single or bifid. Preclypeal spines arising from tubercles, pale, very long, apical part fine and hair-like. Median hairs of mouth-brushes slightly curved, minutely serrated along one side. Mentum triangular, 6-8 teeth either side of central one. Lateral hairs of body fairly well developed. Those on *thorax* many-branched, some arising from large chitinated tubercles, without spine-like processes, except a small one on larger meta-thoracic (metapleural) tubercle. Number of branches in dorso-lateral tufted hairs on first five abdominal segments are respectively 7-10, 5-8, 2-3, 2, 2 (in *A. kuchingensis* the corresponding numbers are 4-6, 3-4, 2-3, 2, 2). Slightly ventral to the tufted hairs referred to there is another smaller hair, which on segments I and II is usually fine and divided into several branches in *A. obturbans*, whereas in *A. kuchingensis* it is larger, and either single or 2-branched. Comb of 6-11 strong teeth, with large basal and smaller apical parts, the latter bluntly pointed and fringed with hairs on both sides to apex. One stout subplumose hair slightly ventral to comb. Subsiphonal tuft large, with about 8-10 long subplumose branches. *Siphon* short and wide, about $1\frac{1}{4}$ to twice length of width at base (when pressed flat, width across base is about equal to length). No pecten. Hair-tuft

* Brug 1924 a, p 439; Senior-White 1927, p 74; Edwards 1927, p 119; Martini 1930, p 258

of 2 fine inconspicuous branches, at about $\frac{1}{3}$ of length from base. Tracheæ within siphon much swollen, nearly filling tube on basal $\frac{1}{3}$, some perforations in tracheæ near base of tube. Anal segment with fairly large chitinated saddle, $1\frac{1}{2}$ inconspicuous, lying slightly below posterior corner of saddle, with 4-5 fine branches, osc of 4-5 long barbed branches all about same length, asc of 3 similar branches. Both pairs of papillæ very long, sausage-shaped, with rounded ends. Fan not large, of about 10 hairs, each divided into several branches, longest hairs about length of subdorsal pairs. no definite fan-plate.

HABITAT.—Tree-holes, bamboos, domestic collections of water, often very foul.

DISTRIBUTION.—The commonest species of the genus. Occurs from the PUNJAB to ASSAM and BURMA, and throughout PENINSULAR INDIA to CEYLON.

Known from MALAY PENINSULA (♂♂ examined by Author), CHINA, JAPAN, SUMATRA (♂♂ examined by Editor). Further distribution doubtful (see footnote, p. 314).

183 *Armigeres (Armigeres) kuchingensis* Edwards, 1915

Bull. Ent. Res. v, p. 283 (♂ & ♀). TYPE-LOC. Kuching, Sarawak, vii 1914 (Moulton). TYPE ♂ & ♀ in Brit. Mus.

var. *durhami* Edwards, 1917 (*Armigeres durhami*), Bull. Ent. Res. vii, p. 206 (♂ & ♀). TYPE-LOC. Bukit Kutu, Malay Penin. 3,300', v 1903 (Leicester). TYPE ♂ & ♀ in Brit. Mus.

var. *nongpohensis* Barraud, 1927, Ind. Journ. Med. Res. xiv, p. 539.

var. *shillongensis* id., ib.

var. *dibrugarhensis* id., ib.

ADULT *.—Variable in markings of mesonotum and in extent of white scaling on abdominal sternites, in some forms resembles *A. obturbans* so closely that it is often difficult to separate ♀♀ with certainty. Scaling of head and appendages, markings of pleuræ, and coloration of legs as in *A. obturbans*, but palpi of ♀ about $\frac{1}{2}$ length of proboscis instead of about $\frac{1}{3}$. Abdominal sternite VII, in all forms, black scaled, with narrow subapical white band, as in *A. obturbans*. Claws of middle legs of ♂ equal in size, both small, as in most other species of the genus.

Type-form.—Mesonotum covered with yellowish-brown scales; no distinct pale margin. Abdominal sternites II-VI white scaled, without apical black bands. In the ♂ mesonotal scaling is often darker, with indications of pale margin, thus being intermediate between this form and var. *nongpohensis*, described below.

* Barraud 1927 b, p. 538. The type-form and the four forms described as varieties represent types of individual variation only, intermediate forms frequently occur.

var *nongpohensis*.—Resembles *A. obturbans* in scaling of mesonotum, but pale border is white, not creamy or yellow. Sternites marked as in type-form

var. *shillongensis* —Mesonotum marked as in *nongpohensis*. Abdominal sternites V and VI with narrow apical black bands, III and IV without complete apical black bands, being almost entirely white.

var *durhami* —Mesonotum marked as in var *nongpohensis*. Abdominal sternites III–VI with black apical bands, those on III and IV much wider than those on V and VI

var *dibrugarhensis* —Closely resembles *A. obturbans*, but pale margin of mesonotum is white, and black bands on venter usually narrow

♂ —*Hypopygium* (fig 77, b) very similar to that of *A. obturbans*, but differing as follows:—*bl* with 3 (occasionally 4) flattened curved spines which bend towards anal segment, in dissected and flattened preparations these spines curve inwards on to the coxite, and not outwards as in *A. obturbans*. Coxite shorter than in *obturbans*, style very similar, but tip reaches to basal lobe when style is depressed against coxite. In some specimens there is a swelling on the inner side of style (seen occasionally also in *obturbans*)

LARVA (described from skins definitely identified as this species by examination of hypopygia of resulting ♂♂) —Very similar to *A. obturbans*, but the following small differences have been found in the material available:—Number of branches in dorso-lateral tufted hairs on first five abdominal segments are, respectively, 4–6, 3–4, 2–3, 2, 2 (the corresponding numbers in *A. obturbans* are 7–10, 5–8, 2–3, 2, 2). Slightly ventral to the tufted hairs referred to there is another, rather smaller, hair, which, on segments I and II in *A. kuchingensis*, is long, fairly stout, and either single or 2-branched. In *A. obturbans* this hair is usually smaller, finer, and divided into several branches. Distance between frontal hairs *B* and *C* more than twice that between *B* and *d* (in *obturbans* the three hairs are about equidistant). Comb-teeth, when seen flat, with wide, frayed, apical margin, not bluntly pointed, as in *obturbans*. (This last character may not be entirely reliable, as the teeth vary in shape, and may have a very different appearance when not mounted flat. It appears, however, that in *kuchingensis*, at least, some of the teeth have a wide, frayed, rounded tip, whereas this appearance is seldom seen in *obturbans*.)

HABITAT —Bamboos.

DISTRIBUTION —Common in ASSAM, where the type-form, the four varieties referred to above, and intermediate forms all occur. The type-form is found chiefly during the cold weather or more or less dry season, the other varieties being found during the rains. Of the four varieties, all except

var *dibrugarhensis*, occur chiefly on the hills var *nongpohensis* has also been taken in the EASTERN HIMALAYAS, Kurseong, ix. 1922 (*Barraud*)

Specimens from Nagargah, BOMBAY DECCAN, viii 1921 (*Barraud*), and from the ANDAMANS, ix 1911 (*Christophers*), resemble var. *durhami*

184 *Armigeres* (*Armigeres*) *theobaldi*, nom. nov

Stegomyia crassipes Theobald (nec v. d. Wulp), M. C. I., p. 320, 1901.

Desvoidia apicalis Theobald, 1910 (nec Theobald, 1908), Roc. Ind.

Mus. iv., p. 5 (♀) TYPE-LOC. Balighat, near Puri, Orissa,
x 1908 (*Annandale*) TYPE ♀ in Ind. Mus.

ADULT ♀.—A large yellowish species with median apical yellow markings on abdominal tergites, and lateral white patches, not visible dorsally, venter white scaled. Distinguished from *A. (L.) flavus*, which it resembles in general appearance, by the shorter palpi and unbanded tarsi, by absence of black scales behind anterior spiracle, and by presence of white scales and bristles in this position.

Palpi about $\frac{1}{2}$ length of proboscis. Clypeus dark brown, with a number of white scales on each side. Mesonotum covered with dark brown narrow scales, and with a border of yellowish scales all round margin. Legs very similar to those of *A. obturbans*, but tibiae and tarsi yellowish when viewed from certain angles.

♂ and LARVA.—Unknown.

DISTRIBUTION.—ORISSA type-locality*, as given above, Singhbhum*, 1923 (*Wats*). N. KANARA. Karwar, 1902 (*Cogill*). BURMA. Thayetmyo, viii. 1893 (*Watson*) † ASSAM. Cherrapunji, 18 x. 1920 (*Senior-White*).

Not known from elsewhere.

185. *Armigeres* (*Armigeres*) *aureolineatus* (*Leicester*), 1908.

Cul. Malaya, p. 70 (*Desvoidia*) (♂ & ♀). TYPE-LOC. Ampang, Malay Penin. (*Leicester*) TYPE ♂ & ♀ in Brit. Mus.

ADULT ♂.—Distinguished by golden lines on thorax, as noted in key. Abdominal markings resemble those of some species of subgenus *Leicesteria*. Sternites III–VI with narrow black apical bands. Tibiae and tarsi dark brown, without pale bands.

♂.—*Hypopygium* (fig. 77, c) *bl* with 3 strong spines. Style rather wide, with comb of 15 or 16 stout teeth along a little more than apical $\frac{1}{2}$. A conspicuous tuft of hairs at apex of coxite near base of style.

† Barraud 1927 b, p. 588.

‡ Described by Theobald (1901) in error as *crassipes* v. d. W.

§ Edwards 1917, p. 206; Barraud 1927 b, p. 541.

LARVA † —Apparently very similar to that of *A. obturbans* but according to Senior-White's figure there are fewer teeth in the comb (5 or 6 are shown), and the siphon-tube is a little longer. In his figures of these larvæ the subposterior siphonal hair and antennal hair are not shown, and one pair of fronta hairs (*C*) has been omitted from the figure of *A. obturbans* larva, so that a comparison on these points is not possible.

HABITAT —Coconut-shells, etc

DISTRIBUTION —N. KANARA Karwar *, ix 1921 (*Barraud*)
MALABAR COAST *, x 1915 (*Khazan Chand*). Ceylon
Colombo, 1913 (*James*), Suduganga, xii. 1919 (*Senior-White*).

Known also from MALAY PENINSULA, COCHIN CHINA and BORNEO

Subgenus **LEICESTERIA** Theobald, 1904.

Entom xxxvii, p 211. Genotype, *L longipalpis* Loew.

Chaetomyia Leicester, 1908, Cul Malaya, p 100. Genotype
L flava Loew

Brevirhynchus Theobald, 1908, Rec Ind Mus ii, p 293. Genotype, *B magnus* Theo

Leicesteromyia Brunetti, 1912, Rec Ind Mus iv, p 152, nom nov for *Chaetomyia*

Distinguished from subgenus *Armigeres* as follows:—No postspiracular bristles, but this area with flat black scale anteriorly and white scales posteriorly. Palpi of ♀ from about $\frac{1}{2}$ to about $\frac{3}{4}$ length of proboscis. Mesonotum more or less produced over the head and in some species narrow and laterally compressed. Style of ♂ with a group, or transverse row, of spines at the tip, and a small basal lobe on inner side of coxite bearing spines or processes.

A flavus is peculiar in having a few scales and minute hairs on postnotum, one or two lower mesepimeral bristles and unusually short hind tibiae.

Key to Adults †.

- | | |
|---|---|
| 1 Hind tarsi with narrow pale rings (sometimes faint) | 2 |
| Hind tarsi entirely dark | 0 |

† Senior-White 1927, pl vii

‡ [Four Malayan species have not yet been found in the Indian region, but may occur. These are:—(1) *dolichocephalus* Loew, resembling *armulicarsus*, but without scales on olypous and tarsal rings faint or absent, ♂ hypopygium with 7 spines on style, 4 on *bl*; (2) *cingulatus* Loew, resembling *longipalpis*, but ♂ style without membranous expansion and no scale-patch on coxite, style short, with 4 spines longitudinally placed and some scales, (3) *pectinatus* Edw., resembles *digitatus* in colouring, but ♂ style more expanded at tip, with 9 spines, *bl* with 2+ spines, coxite more hairy, (4) *pendulus* Edw., with abdomen unbanded and without yellow markings, hind tarsi very faintly ringed, hypopygium small and pendulous, with only 3-4 spines on style and dense scale patch on coxite, 9th sternite unusually large.]

- 2 Abdominal tergites with median apical yellow markings, postnotum with a few scales and some minute hairs, one or two lower mesepimeral bristles, tibia of hind leg distinctly shorter than that of fore leg *flavus*, p. 321
Abdominal tergites marked otherwise, postnotum bare, no lower mesepimeral bristles, tibia of hind leg as long as, or longer than, that of fore leg . . . 3
- 3 Tergites with median basal yellow markings *maignus*, p. 324
Tergites otherwise marked . . . 1
- 4 Tergite II only with median basal white patch, ♀ with simple claws and distinct white ring on palpi just before middle *annulipalpis*, p. 327
No median basal pale mark on tergite II, ♀ with toothed claws (as usual) and without a distinct white ring on palpi . . . 5
- 5 Tergites IV-VII or V-VII with lateral yellow spots, as well as oblique white lateral marks *annulatus*, p. 325
Tergites without lateral yellow markings *nuchatus*, p. 328.
- 6 Tergites II-VII with lateral white markings, curving on to dorsum, where they become yellowish and form complete apical bands. *longipalpis*, p. 320.
Tergites with lateral white markings, not produced on to dorsum to form bands . . . 7
- 7 Tergites with some lateral yellow or creamy markings, style of ♂ with 9-14 spines at apex . . . 8
Tergites without lateral yellow or creamy markings (though with the usual white lateral spots), style of ♂ with only 5 spines at apex . . . *digitatus*, p. 330
- 8 Tergites VI and VII only with distinct lateral yellow patches, style of ♂ with about 15 stout tooth-like spines at apex. . *omissus*, p. 330
Tergites IV-VI with indistinct lateral creamy or yellowish markings; style of ♂ with 9 finger-like spines at apex . . . *dentatus*, p. 331

186 *Armigeres (Leicesteria) flavus* (Leicester), 1908.

Cul. Malaya, p. 101 (*Chatomys*) (♂ & ♀) TYPE-LOC. : Malay Penin. (*Leicester*). TYPE: 3 ♀♀ co-types in Brit. Mus.

Leicesteria apicalis Theobald, 1908, Rec. Ind. Mus. ii, p. 291 (♂ & ♀)

TYPE-LOC. : Lushai Hills, Assam (*B. L. Macleod*). TYPE: ♂ & ♀ in Ind. Mus.

Brevirhynchus apicalis Theobald, 1910, Rec. Ind. Mus. iv, p. 7 (♀) TYPE-LOC. : Sylhet, Assam (*Major Hall*). TYPE. ♂ in Ind. Mus.


ADULT *.—Medium-sized to large species (wing 3-5 mm), distinguished from all others of the genus occurring in India by the short hind tibia.

♀—*Head*. with scales on vertex mostly creamy-yellow, with three patches of black scales in front near eye-margins,

* Edwards 1914, p. 258 (syn.). Strickland 1917; Barraud 1927, p. 541

these patches sometimes joined. Palpi and proboscis all black scaled, palpi $\frac{2}{3}$ to nearly $\frac{1}{2}$ as long as proboscis. Clypeus without scales, dark brown, contrasting with orange-yellow tori. *Thorax* not compressed and not distinctly produced over head, mesonotal scales mixed dark brown and golden-brown, with a rather broad border of creamy-yellow round

Fig 78



♂ hypopygial structures (style and basal lobe) of *Leicosternus*, subgenus *Leicosternus*: a, *magnus*, b, *annularis*, c, *omissus*; d, *dentatus*, e, *longipalpis*, f, *flavus*, g, *inchoatus*, h, *deutatus*.

front, *apn* and *ppn* with blackish flat scales above, creamy-yellow scales below. A few short and inconspicuous pale hairs at tip of postnotum, one strong lower mesepimeral bristle present. *Wings* dark scaled. *Legs* dark brown, pale area on outer side of hind femur not quite reaching tip,

front and middle tarsi with faint pale rings, hind tarsi with more distinct, though narrow, pale rings over joints 1-2 and 2-3 and sometimes at base of 4 and 5. *Abdomen* dark brown above, tergites II-VI with narrow yellow apical bands, II-VII with large creamy lateral basal patches which tend to curve upwards beyond middle of each tergite but are not connected with the yellow bands and are scarcely visible dorsally.

♂—Palpi with narrow yellowish rings at middle of long segment and at bases of last two segments, longer than proboscis by about length of last two segments. Ornamentation as in ♀. *Hypopygium* (fig 78, f). *bl* with 3 strong spines placed close together and some long hairs. Style with 5 or 6 blunt spines at apex. *Lobes* of 9th more densely hairy than in other species. No definite scale-patch on coxite.

LARVA (fig 76, e-g)—This has been described by Edwards and Given (1928, p 342) as quoted below. Two skins from isolated larvæ obtained from bamboos at Nongpoh, Assam, vii 1922 (*Barraud*), agree with this description. The larva, as well as the adult, possesses some unique characters. "Head much as in the other species of the genus, roundish in shape and pale in colour. Clypeal spines hair-like but set on large tubercles. Dorsal hairs placed far forward. *d* placed nearest the front, dendriform, with about 8 branches, *B* simple or 2-branched, *C* placed obliquely behind *B*, 3-4-branched, both rather small. Antennæ nearly cylindrical, smooth, with a fine hair at one-third of their length. Mentum rather broad, with about five teeth on each side of the triple median tooth. *Thorax* with not more than two simple hairs in any of the lateral tufts, which are set in distinct chitinous plates that have no trace of a projecting spine. *Abdomen* with all the hairs of segments 1-7 simple except the two lateral hairs of the first segment and the dorsal hair of the seventh, which are double, the dorsal hairs are almost as long and thick as the lateral hairs. Eighth segment with one strong lateral tuft composed of 5-8 stout simple branches and set in a small plate, other four hairs of eighth segment small, each with several branches. *Comb* of about 10 small scales in a patch, each scale has a pear-shaped base, the free portion being divided to the base into 3-5 portions, each of which is irregularly frayed towards the tip. *Anal segment* with two chitinous plates, one dorsal and the other ventral, the dorsal plate larger than the ventral and extending below the middle, not quite meeting the ventral plate. Outer dorsal hairs are 2- or 3-branched, inner 3-5-branched, both slightly plumose. Brush composed of five pairs of double (rarely triple), slightly plumose hairs, the usual chitinous transverse

bars hardly indicated. A very small branched lateral hair set between the plates. Gills very stout, round-ended, over twice as long as the saddle. *Siphon* moderately dark, tapering, index about 1.0, without pecten, but with a small two-haired tuft at four-fifths of its length. No aous. Tracheæ very large, with numerous perforations near base of siphon.

"A unique feature of this larva, found in no other mosquito larva hitherto known, is the presence of a ventral chitinous plate on the anal segment. The comb-teeth are of unusual form, but the remaining characters are similar to those of other *Armigeres* larvæ."

HABITAT—Bamboos and coconut-shells, pitcher-plants (Singapore).

DISTRIBUTION—Fairly common in forested areas of ASSAM, BURMA, BENGAL, and EASTERN HIMALAYAS up to 5,000' or more, in the WESTERN GHATS from the Konkan to Malabar and MADRAS.

Known also from MALAY PENINSULA, SIAM, COCHIN CHINA, SUMATRA, JAVA, and BORNEO.

187. *Armigeres (Leicesteria) magnus* (Theobald), 1908

Rec Ind Mus u, p 293 (*Brevirhynchus*) (♂ & ♀) TYPE-LOC Sylhet, Assam, v 1905 (Hall). TYPE ♂ & ♀ in Ind Mus

Leicesteria annulata (♂ only) Leicester, 1908, Cul Malaya, p 99 TYPE-LOC Kuala Lumpur, Malay Penin (*Leicester*) TYPE ♂ in Brit Mus

Toworhynchites rectirostris (Giles MSS) Theobald, 1910, MC v, p 214 TYPE-LOC Philippine Is TYPE in Brit Mus

ADULT *.—Distinguished by basally-banded abdominal tergites. Rather large species. Wing 4.5–5 mm.

♀—**Head**: mainly covered with broad, flat, yellowish and dark brown scales, the latter forming a pair of sublateral and a pair of lateral patches. Tori with pale scales on inner sides. Palpi from $\frac{5}{8}$ – $\frac{3}{4}$ length of proboscis, scaling of palpi light brown, except towards base; proboscis dark brown. **Thorax**: mesonotum covered with light brown narrow scales, a pale border all round margin. Scutellar scales flat and dark brown. Flat white or creamy scales covering *apm*, lower part of *ppm*, and larger part of pleuræ, a small patch of black scales behind anterior spiracle; pale brownish flat scales on upper anterior part of *ppm*. **Wings**: scales brown, those towards base of costa and near base of vein 1 light when viewed from certain angles. **Legs**: mainly dark brown when seen from the front, almost entirely pale when

* Theobald 1910 b, p 145, Edwards 1914 a, p. 259 (syn.); Dyar & Shannon 1925, p 74, Barraud 1927 b, p 542

seen from behind, hind femur creamy or white except along dorsal edge pale scaling at tips of femora and tibiae and at bases of tarsal segments, more distinct in some specimens than in others. *Abdomen* tergites dark brown, with median basal yellow markings on II-V, wider dull yellow basal markings on VI and VII, wide lateral white markings from base of each tergite nearly to apex, the tips of these patches usually visible dorsally, lateral basal yellow markings on III-VII or IV-VII, those on VI and VII usually joined to the dorsal yellow markings, forming complete bands. Sternites white, with narrow apical black bands on III-VI.

♂ —Palpi longer than proboscis by length of apical segment, dark brown, with fairly distinct yellowish rings at bases of last two segments, and a wider pale area nearer base. Markings very similar to those of ♀. *Hypopygium* (fig 78, a). *bl* with 4 strong spines all about same length. Style expanded at tip, with a row of 12 finger-like spines and a few scales at about middle.

[LARVA —Four isolated skins sent by Dr. R. B. Jackson in 1933 from Hong Kong rather closely resemble *A. flavus*, having the same peculiar forms of comb-teeth, but differing in the following points —Mentum more triangular, one of the metathoracic tufts usually triple, as is the dorsal tuft on VII, mid-lateral tuft on VIII with only 2, or at most 3, branches, comb-scales fewer, about 6 in number, ventral plate of VIII very small and inconspicuous, perhaps sometimes absent.

HABITAT (of Hong Kong specimens) —Pitcher-plants.]

DISTRIBUTION —Similar to that given for *A. flavus*, except that it does not appear to have been recorded from Siam.

188. *Armigeres (Leicesteria) annulitarsis* Leicester, 1908

Cul. Malaya, p. 99 (♂ only). TYPE-LOC : Kuala Lumpur, Malay Penn (Leicester). TYPE : ♀ in Brit Mus.

ADULT* —Distinguished by compressed thorax, yellow lateral markings on abdomen, and ringed tarsi. Wing about 4 mm.

♀.—*Head* somewhat elongate, mainly covered with flat black scales, a line or stripe of white scales from nape to vertex in middle, a narrow pale border to eyes, and a large patch of white scales low down at each side. Tori brown, with pale scales on inner sides. Clypeus very dark, with some pale scales each side. Palpi and proboscis brownish-black, the former white-tipped and about $\frac{1}{2}$ length of proboscis. *Thorax*.

* Edwards 1914 a, 260, 1923, p. 3, Barraud 1927 b, p. 543

mesonotum distinctly produced over head, narrow and laterally compressed, dorsum covered with brown scales; a fairly wide pale border all round margin. Scutellar scales flat, dark and light brown mixed. Broad white scales covering larger part of *apn* and *pleuræ*, those on *ppn* dark brown, a patch of very dark scales behind anterior spiracle. *Wings* dark scaled. *Legs* brownish-black, femora extensively pale beneath, a few pale scales at tips of femora and tibiae, and narrow pale rings at bases of first two, three, or four tarsal segments. *Abdomen* tergites black, with lateral oblique white markings, tips of which are just visible dorsally, lateral basal yellow patches also on tergites IV-VII or V-VII.

♂.—Palpi a little longer than proboscis and marked with 3 pale rings. Other markings very similar to those of ♀ *Hypopygium*. *bl* with about 7-10 long, slightly clubbed processes, a few long hairs, and a dense patch of shorter hairs. Style with 5 spines at tip, the outer ones longer than inner. No scale-patch on coxite.

LARVA.—Very similar to those of other species of the genus. *Antenna* about 0.26-0.27 mm long and about 6 times length of greatest width. Hair-tuft represented by a single rather long fine hair at about $\frac{1}{2}$ length of shaft from base. Procoxal spines long, with hair-like terminal part as usual. *Frontal hair* *A* 3-4-branched, *B*, *C*, and *d* standing far forward, *d* slightly internal and anterior to *B*, fairly large, with a number of fine branches originating some little distance from a stout base, *B* single, stout, and long, *C* standing almost directly behind *B*, with about 6 fine branches. *Lateral thoracic hairs* 1-3-branched (one specimen with several hairs 4-5-branched), arising from small tubercles without spines. Number of branches in dorso-lateral hairs of first five *abdominal* segments are, respectively, 3, 3, 1, 1, 1, the single hairs sometimes divided into two. A second fairly large hair, slightly ventral to dorso-lateral series, on I and II, either single or 2-branched. A very long hair with 2 stout branches (in one specimen 3-branched) on VIII dorsal to comb, subsiphonal hair rather large and 4-5-branched, but much shorter than dorsal hair. Comb of about 20-25 teeth, very similar in form to those of *A. flavus*, but variable in shape; base large, usually longer than the terminal fringe. Siphon short and wide, about $1\frac{1}{2}$ - $1\frac{3}{4}$ length of diameter at base, diameter at tip less than $\frac{1}{2}$ that at base. Siphonal hair at about $\frac{3}{4}$ from base, either single or split distally into two. *Anal segment* with small chitinous saddle, its posterior margin smooth. Subdorsal hairs set in a single small dark plate; *osc* with 2-3, *isc* with 6-8, strong subplumose branches, subequal in length, *lh* at about middle of segment, towards

posterior margin, some distance below border of saddle, very small, as usual, with several fine branches. About 6 pairs of fan-hairs, each with 5-6 branches. No fan-plate, and no trace of ventral plate adjoining fan. Papillæ very long.

HABITAT —Bamboo-stumps (*Barraud*, Bombay Deccan), living bamboos bored by a beetle (*Hacker*, Malaya).

DISTRIBUTION —ASSAM. Shillong*, Nongpoh*, Khasi Hills dist. and Haflong*, Cachar Hills, vi-viii 1922 (*Barraud*). BOMBAY DECCAN. Nagargali*, Belgaum dist., viii 1921 (*Barraud*). NORTH KANARA. Karwar* and Yellapur*, ix & x 1921 (*Barraud*). NILGIRI HILLS*, x 1915 (*Khasan Chand*).

Recorded also from MALAY PENINSULA, SIAM, COCHIN CHINA, SUMATRA, and FORMOSA.

189 *Armigeres (Leicesteria) annulipalpis* (Theobald), 1910

Rec. Ind. Mus. iv, p. 6 (*Brevirhynchus*) (♀). TYPE-LOC. Madathoray, S.W. base of Western Ghats, Travancore, ix 1908 (*Ammandale*). TYPE ♀ in Ind. Mus.

ADULT † —Resembles *A. (L.) annulitarsis*, but easily distinguished by the ringed palpi and simple claws of ♀.

♀.—Palpi slightly more than $\frac{1}{2}$ length of proboscis, with a conspicuous white ring at about middle, and some pale scales on upper surface near base. Clypeus dark brown, with some pale scales. Hind femur with a rather wide subapical dark ring all round leg, tibiae with nearly complete basal pale rings, basal white rings on first two segments of fore and mid-tarsi, similar, but more pronounced, whiterings on all hind tarsal segments. In addition to markings on abdomen noted in key, there are on tergites III-VI white scales forming a short curved line towards venter, so that, on these segments, the straight white lateral markings divide near hind margin, one part curving on to dorsum and the other towards the venter. Usually lateral yellowish patches on tergites III-VII or IV-VII, and scattered yellow scaling on dorsum of last few segments. A median white line and subapical white bands on sternites.

♂ and LARVA —Unknown.

DISTRIBUTION —SOUTH INDIA. Travancore*, type-locality, as given above, Malabar Coast, Tamarchery*, x 1915 (*Khasan Chand*). BURMA. Papun*, near Moulmein, x 1926 (3 ♀♀ in M.S.I. coll.).

Recorded also from SUMATRA.

† Edwards 1921 b, p. 74, Barraud 1927 b, p. 544.

190 *Armigeres (Leicesteria) incheatus* Barraud, 1927.

Ind. Journ. Med. Res. xiv, p. 544 (♂ & ♀) TYPE-LOC Tindhara, Darjeeling dist., x 1922 (*Barraud*) (♂), Kurseong, Darjeeling dist., ix 1922 (*Barraud*) (♀) TYPE ♂ and allotype ♀ in Brit Mus

ADULT.—Resembles *A. (L.) annulatus*, but without yellow marks on abdomen. Wing 4-4.5 mm.

♀.—*Head*: scaling similar to *A. annulatus*. Clypeus dark brown, without scales. Palpi a little more than $\frac{1}{2}$ length of proboscis, brownish-black, with a narrow pale ring much nearer base than apex. Proboscis dark brown. *Thorax*: mesonotum moderately produced over the head. A narrow border of pale scales, which are broad from wing-root to *ppn*, narrow round front margin, dorsum covered with dark brown narrow scales, some lighter ones on either side of anto-scutellar space in some specimens. Scutellar scales dark and pale brown mixed. Postnotum dark brown, *apn*, pleura, and coxæ mainly covered with broad white flat scales, darker scales in middle of *ppn* and behind anterior spiracle, as usual. *Wings*: dark scaled. *Legs*: dark brown, with pale markings. Fore and mid-femora and tibiae dark brown anteriorly, almost entirely pale posteriorly, hind femur more extensively pale, but dark along dorsal edge and on outer side at knee. Tibia with a few pale scales at tips. Narrow basal pale markings on first two or three segments of fore and mid-tarsi, and on first four segments of hind tarsi. *Abdomen*: tergites II-VII with oblique white lateral markings, from base to about middle of each segment, produced on to dorsum, but not forming transverse bands. No lateral yellow patches. Venter white at base, following sternites with white patches and apical black bands.

♂.—Palpi a little longer than proboscis, dark brown, with three narrow white rings. Scutellar scales rather lighter than in ♀. *Hypopygium* (fig. 78, g). *bl* with 5 strong, pointed spines. Style expanded apically, with a row of 10 finger-like, rather blunt spines, the outermost longer than others. No scale-patch on coxite.

LARVA.—Unknown.

HABITAT.—Bamboos.

DISTRIBUTION.—Known only from type-localities.

perhaps, also of *angulatus*) are variable, at least in the ♂ sex. In the type ♀ of *longipalpis* the abdomen is unbanded, but there is no certainty that this is correctly associated with the type ♂, it may belong to one of the allied species which Leicester had undoubtedly included under *longipalpis*. There appear to be no adequate grounds for upsetting Edwards's restriction of *longipalpis*. This name is, therefore, used for the species called *angulatus* by Barraud, the true *angulatus* (as figured by Edwards) has not yet been found in the Indian region.]

191 *Armigeres (Leicesteria) longipalpis* (Leicester), 1904*†

In Theobald, Entom xxvi, p 211 (♂ & ♀) TYPE-LOC Kuala Lumpur, Malay Penin (*Leicester*) TYPE ♂ & ♀ in Brit Mus

ADULT—Usually distinguished by complete apical bands on abdominal tergites and dark tarsi Wing about 4 mm

♀.—*Head* scales mainly black, but a narrow pale border to eyes, and a white patch low down at each side Tori and first flagellar segment of antennæ with some pale scales. Clypeus, palpi, and proboscis brownish-black, the first with some narrow pale scales each side directed outwardly Proboscis with a more or less evident pale line beneath, as in several other species Palpi entirely dark, about $\frac{1}{2}$ length of proboscis *Thorax* rather narrow, and produced over the head Scales dark brown, with a narrow pale margin all round. *Wings* dark scaled *Legs* hind femur white except dorsally, tibiae and tarsi entirely dark, but paler when seen from behind, as in several other species *Abdomen*. tergites brownish-black, II–VII with oblique white patches, running from base to near hind margin of each segment, continued on to the dorsum, where they become yellowish and form complete bands on II–V these bands are usually very slightly subapical, on VI and VII they are usually apical. [In some specimens the bands may be incomplete dorsally on VI and VII or even on all segments] No lateral yellow patches

♂—Palpi a little longer than proboscis, entirely dark Markings as in ♀ *Hypopygium* (fig 78, e) *bl* with 3 very short stout spines Dorsal border of coxite with a thick patch of scales along basal $\frac{1}{2}$ Style characteristic, comparatively short and wide, divided apically into two parts, one of which carries 3 fairly short spines

LARVA—Unknown

HABITAT—Bamboos

* Edwards 1914 a, p 260, Barraud 1927 b, p 545 (*cingulatus*).

† [It was suggested by Barraud (1927 b, p 546) that Edwards (1914 a, pp 260 and 261) made a mistake in figuring the hypopygium of *A. (L.) longipalpis* and *cingulatus*, transposing the two figures The reason for this suggestion was that Indian specimens with the hypopygium as figured by Edwards for *longipalpis* agreed with Leicester's description of *cingulatus*, and differed from that of *longipalpis* in the markings of the abdomen But against this it must be stated that (1) the hypopygia of the two types were carefully labelled at the time of figuring, and it is unlikely that they were mixed in mounting; (2) the two ♂ types as now standing in the British Museum collection show no appreciable difference in ornamentation, both having complete apical bands on tergites II–V, incomplete bands on VI and VII, and (3) that there is a ♂ in the British Museum from Sarawak in which the hypopygium has the structure figured by Edwards for *longipalpis*, but the white abdominal markings do not form complete bands dorsally on any segment It seems clear, therefore, that the abdominal markings of *longipalpis* (and,

[Footnote continued on previous page.

DISTRIBUTION —ASSAM Nongpoh*, Khasi Hills dist., and Haflong*, Cachar Hills, vii & viii 1922 (*Barraud*). EASTERN HIMALAYAS Tindharia* and Kurseong*, both Darjeeling dist., ix & x 1922 (*Barraud*)

Known from MALAY PENINSULA, and recorded from SUMATRA and BORNEO

192 *Armigeres (Leicesteria) digitatus* Edwards, 1914

Bull. Ent. Res. iv, p. 262 (♂ & ♀) TYPE-LOC. Ulu Gombak, F. M. S. viii & ix 1903 (*Daniels*) TYPE ♂ & ♀ in Brit. Mus.

ADULT ♀ †—A comparatively small species resembling *A. inchoatus* in general appearance, but distinguished by absence of pale rings on tarsi, the leg-markings being similar to those of *A. cingulatus*. Palpi and proboscis entirely dark brown, former nearly $\frac{3}{4}$ length of proboscis. Mesonotum distinctly produced over the head. Abdomen without any lateral yellow patches, but with oblique white lateral markings not produced on to upper side, the tips being only just visible dorsally in some specimens.

♂.—Markings as in ♀. Palpi a little longer than proboscis, entirely dark brown. *Hypopygium* (fig. 78, h). *bl* with about 7 sharp spines, arranged more or less in two rows. Style rather short, not widened at tip, with 5 blunt spines at tip, somewhat resembling the spread-out fingers of a hand. No scale-patch on coxite.

LARVA.—Unknown

HABITAT —Bamboos

DISTRIBUTION —NORTH KANARA Yellapur*, x 1921 (*Barraud*). NORTH BENGAL (Darjeeling dist.) Sukna* and Tindharia*, ix & x 1922 (*Barraud*); Pashok, 3,500' (*Graveley*).

Recorded also from PHILIPPINE Is. and SUMATRA

193. *Armigeres (Leicesteria) omissus* Edwards, 1914.

Bull. Ent. Res. v, p. 76 (♂ & ♀) TYPE-LOC. Ceylon, Colombo (*James*). TYPE ♂ & ♀ in Brit. Mus.

ADULT †—Resembles *A. digitatus* in having entirely dark tarsi, but differs from that species in having lateral yellow spots on last two abdominal tergites, and from *A. cingulatus* in the absence of pale bands on dorsum of abdomen.

Palpi and proboscis entirely dark in both sexes, palpi of ♀ nearly $\frac{3}{4}$ length of proboscis, those of ♂ longer than proboscis by $\frac{1}{2}$ length of apical segment. Clypeus bare.

† Barraud 1927 b, p. 546

† Barraud 1927 b, p. 547.

Legs black scaled, undersides of fore and mid-tibiae white, hind femora white, except dorsally and near tip. Fore tibiae shorter than mid- or hind pair. Segment I of hind tarsi about same length as tibia. *Abdomen* black scaled dorsally, segment I with a broad white patch on each side, II-VII with large lateral oblique white patches, upper margin of each concave, extending nearly to hind margin of each tergite, and not produced on to dorsum, tips only visible from above in some specimens. Small basal lateral yellow patches on VI and VII only.

♂—*Hypopygium* (fig 78, c). *bl* with 3 strong sharp spines. Style much expanded apically, with a row of strong spines, one fairly long and about 14 short and stout. Dorsal border of coxite with a patch of scales near base.

LARVA —[Very similar to *annulitarsis*, differing chiefly as follows —Dorso-lateral hair on VIII shorter and much less conspicuous. Comb-teeth sometimes fewer (12-25), and rather different in average form, with smaller base and longer, more spiny fringe. Siphon shorter, diameter at tip $\frac{2}{3}$ that at base, tuft only slightly beyond middle of tube, *isc* with fewer branches (4-5)]

HABITAT —Bamboo-stumps

DISTRIBUTION —CEYLON. Colombo, type-locality, as given above, Peradeniya (*Green*). BOMBAY DECCAN Nagargali*, Belgaum dist, viii 1921 (*Barraud*)

Not known from elsewhere.

194 *Armigeres (Leicesteria) dentatus* Barraud, 1927.

Ind Journ Med Res xiv, p 547 (♂). TYPE-LOC. Nongpoh, Khasi Hills dist, Assam, xi 1921 (*Christophers*). TYPE ♂ in Brit Mus

ADULT —Very similar to *A. (L.) mchoatus*, but with dark tarsi. Wing (♂) about 3.5 mm

♂—*Head* mainly black scaled, a pale patch in middle of vertex and narrow pale border to eyes; a band of pale scales low down each side; a small collection of pale upright scales on nape. Tori brown, with small broad pale scales on inner sides and on first flagellar segment of antenna. Clypeus black, bare. Palpi and proboscis brownish-black, former a little longer than latter, proboscis with a narrow pale streak on underside for nearly whole length. *Thorax* mesonotum only slightly produced over the head, covered with narrow dark brown scales, a narrow pale border round front margin from wing-roots. A mixture of flat dark brown and paler scales on scutellum. Scaling of pleurae as in *A. mchoatus*. *Wings*: dark scaled. *Legs*: dark brown, paler posteriorly, hind femur pale ventrally on outer side

to knee no pale knee-spots dorsally on femora, tibiae and tarsi entirely dark *Abdomen* black dorsally, lateral oblique white markings from base of each tergite, not produced on to dorsum, and not reaching hind margins of segments, the tips just visible dorsally, IV-VI with indefinite and indistinct basal lateral yellowish patches *Sternites* with basal pale bands and apical dark bands *Hypopygium* (fig 78, d) . very similar to that of *A. inchoatus*

LARVA and ♀ —Unknown

DISTRIBUTION —Known only from type-locality

Genus **CULEX** Linnæus, 1758.

Syst Nat ed 10, p 602

(For synonymy see under subgenera)

ADULT —Distinguished from all other genera by the presence of well-developed pulvilli (these are best seen under a binocular

Fig 79




Upper figure side view of *Culex (Lutzia) fuscans*, showing typical form of palpi in genus *Culex*, also lower mesepimeral hairs of subgenus *Lutzia* (hairs on mesonotum, ppn, and sternopleura omitted). Lower figure wing of *Culex mimeticus*, showing markings, venation and scaling typical of *Culex*

microscope against a dark background, when they appear pale). Bucco-pharyngeal armature present in ♀♀ (absent in all other genera) No spiracular or postspiracular bristles. Mesonotal bristles well developed. Mesonotal and scutellar scales usually all narrow Other characters given under

subgenera. No very well-marked subgeneric characters have been discovered by which the ♀♀ may be separated, but ♂♂ can be distinguished on characters of the palpi, antennæ, or hypopygium

PUPA—Respiratory trumpet usually of moderate length and with rather small opening. Well-developed dendritic tufts on abdominal segment I. Paddles usually wide and

Fig 80



a-g, pharyngeal teeth of different species of *Culex*: a, *C. (Lutzia) vorax*; b, *C. (Mochthogenes) malaya*; c, *C. (Neoculex) tenuipalpis*; d, *C. (Lophoceratomyia) uniformis*; e, *C. (Oulicomyia) viridiventer*; f, *C. (Culex) bitaeniorhynchus*; g, *C. (Culex) fuscocephalus*. h-k, pharyngeal bar of other genera, showing absence of teeth. h, *Megarhinus*; i, *Uranotaenia*; j, *Armigeres*; k, *Aedomyia*. (Posterior hard palate not shown in any case).

rounded, with 2 short hairs at or near termination of midrib, posterior margin without fringe or serrations

LARVA—In the subgenus *Lutzia* the mouth-parts are modified for predacity, the mouth-brushes being composed of strong curved and fringed lamellæ, frontal hairs usually single, antennal tuft represented by a single short hair;

siphon very short, with pecten extending for nearly the whole length. In the other subgenera the mouth-parts are normal, one or more pairs of frontal hairs with several branches, antennal tuft usually well developed and beyond middle of shaft, siphon of very variable length, with pecten extending along at most basal $\frac{1}{3}$ of tube. The larvæ differ from those of most other genera in having several pairs, or a posterior row, of branched hairs on the siphon. Abdomen without chitinous plates, comb of segment VIII variable in form, but usually composed of a number of scale-like teeth in a triangular patch. Except in the case of *Lutzia* there are no very well-marked subgeneric characteristics.

Egg—So far as is known the eggs are laid in rafts, a number of eggs being cemented together in a boat-shaped mass, which floats on water. The individual eggs are narrowed at one end, and the larva emerges by bursting off a cap at the lower (larger) end, which is in contact with the water-surface.

Notes on bionomics are given under subgenera.

Key to Subgenera

♀♀

- | | |
|--|---|
| 1. Four or more lower mesepimeral bristles
One or two lower mesepimeral bristles, or none | LUTZIA, p. 338
2 |
| 2. Proboscis with a pale ring in the middle
Proboscis without a pale ring | CULEX (part), p. 387.
3 |
| 3. First hind tarsal segment distinctly shorter than tibia
First hind tarsal segment about as long as tibia or longer | BARRAUDIUS, p. 345
4 |
| 4. Dorsal surface of head mainly covered with flat scales
Dorsal surface of head mainly covered with narrow scales, flat scales, if present, confined to sides or eye-margins | [malaya, p. 358.
C. (<i>Mochthogenes</i>)
5 |
| 5. Rather large to medium-sized species, without any broad scales adjacent to eyes
Smaller, usually dark-coloured species, or else with small broad scales on vertex around eye-margins | CULEX (part), p. 387.
6 [p. 351] |
| 6. Wing-scales scanty, especially on vein 6
Wing-scales denser | LOPHOCERATOMYIA,
7 |
| 7. Vertex with small broad scales forming a border to eyes
Vertex without any broad scales along eye-margins | 8.
9. |
| 8. Moderate-sized species
Very small species | CULICOMYIA, p. 376.
C (<i>Mochthogenes</i>)
[pluvialis, p. 356. |
| 9. No lower mesepimeral bristles
One or two lower mesepimeral bristles | C (<i>Neoculex</i>) <i>brevi-</i>
10 [palpis, p. 348. |
| 10. Dorsum of abdomen with very narrow basal pale bands
Dorsum of abdomen unbanded | C (<i>Neoculex</i>) <i>tenui-</i>
[palpis, p. 351.
MOCHTHOGENES (part),
[p. 352. |

♂♂

- | | |
|---|---|
| 1. Four or more lower mesepimeral bristles
One or two lower mesepimeral bristles, or none | LUTZIA, p. 338
2 |
| 2. Antenna with matted tuft, or tufts, of hairs, and sometimes scales, on one or more flagellar segments | [p 359
LOPHOCERATOMYIA,
3 |
| 3. Palpi longer than proboscis
Palpi shorter than proboscis | 4
6 |
| 4. Palpi with a row of long scales projecting obliquely inwards and downwards from apical part of long segment
Palpi without such scales | CULICIOMYIA, p. 376
4 |
| 5. First hind tarsal segment shorter than tibia
First hind tarsal segment as long as, or longer than, tibia | BARRAUDIUS, p. 345
CULEX, p. 387 |
| 6. Palpi from $\frac{1}{2}$ – $\frac{2}{3}$ length of proboscis
Palpi not more than about $\frac{1}{2}$ length of proboscis | NEOCULEX, p. 347
[p 352
MOCHTHOGENES, |

*Key to known Larvæ (4th stage) **

- | | |
|---|---|
| 1. Mouth-parts adapted for predacity, pecten extending along nearly whole length of siphon, frontal hairs A, B, C usually single, but may be split (fig 82) | [pp 342-344
C (Lutzia), all species, |
| Mouth-parts normal, pecten confined to about basal $\frac{1}{2}$ of siphon or less, one or more pairs of frontal hairs with several branches | 2 |
| 2. Hair-tufts on siphon arranged in a close-set zigzag row along mid-posterior line, several hairs being between the pectens, no lateral hair-tufts | C (B) modestus, p. 347 |
| Hair-tufts on siphon usually arranged in pairs on either side of mid-posterior line, but, if some are more or less in a zig-zag row, none are between the pectens, or there are lateral tufts in addition | 3 |
| 3. Comb of 4-8 large sharp teeth
Comb of more numerous smaller scale-like teeth arranged in a triangular patch | 4
7 |
| 4. Mentum with very numerous minute teeth (fig 91)
Mentum with fewer teeth, some or all of which are of normal size | [p 393
C (C) bitensorhynchus,
5 |
| 5. Pecten of 2-4 small teeth close together at base of siphon, mentum with about 20 teeth either side of central one, the 12 teeth on either side nearest base very small, subapical bristles on antenna about mid-way between tuft and tip of shaft (fig 91) | [sinensis, p. 395.
C (C) cornutus, p. 396, |

* As the larvæ do not exhibit clearly-defined subgeneric characters, one key is given for all the known Indian larvæ of this genus. [At the suggestion of the author this key has been revised and largely re-written]

- Pecten of 8-14 teeth extending some distance from base of siphon, mentum with about 6 teeth on either side of central one, all about same size, sub-apical bristles on antenna much nearer to tip of shaft than to tuft
- 6 Anal papillæ as long as siphon, latter with 5-6 pairs of long 2-branched hairs, each about $\frac{1}{2}$ length of siphon, pecten-teeth smaller than comb-teeth (fig 96) [p 407
C (C) whitmorei,
 Anal papillæ about $\frac{1}{2}$ length of siphon or less, latter with 6-7 pairs of rather short hairs with 3-5 branches, of which longest are about $\frac{1}{2}$ length of siphon, larger pecten-teeth as large as comb-teeth (fig 93) C (C) vishnui, p 402
- 7 Body-integument covered with a dense pile of small dark hairs, chitinous ring of anal segment with numerous stiff bristly spines on and near posterior margins. [p 371
C (L) unifornis,
 Body without such pile, chitinous ring of anal segment smooth, or with at most a few short spines 8
- 8 Preclypeal spines remarkably thick, black and rather blunt-tipped, anal papillæ very short and rounded, hardly longer than wide, siphonal index 2-2.5 C (C) sitrens, p 399
- Preclypeal spines not so thick, sharply pointed, anal papillæ at least 3 times as long as wide, usually longer than anal segment 9
- 9 Preclypeal spines rather short, stout, black, and usually straight 10
- Preclypeal spines long, slender, pale, and curved 26
- 10 Frontal hairs *B* and *C* not reaching front of head, with only 1-2 almost simple branches, *C* quite small and much shorter than *B*, pecten-teeth with small denticles extending the whole length of one side, those on outer $\frac{1}{2}$ very fine, preapical bristles practically at tip of antenna 11
- Frontal hairs *B* and *C* both reaching front of head or beyond, with distinctly plumose branches 13.
11. Siphon fairly broad at base, very narrow at tip (hardly more than $\frac{1}{2}$ of basal diameter), siphonal tufts very long, longest about 4 times length of diameter of tube at point of attachment [p. 352.
C (N.) tenuisulcata,
 Siphon about $\frac{1}{2}$ as broad at tip as at base, and more evenly tapering, longest tuft hardly more than twice length of diameter of tube 12 [p 356.
C. (M.) castrensis,
 Comb-teeth fringed to tip C (M) malaya, p. 359*.
 Comb-teeth fringed at base only, ending in a long point

* Position in key uncertain, as frontal hairs have not been described

- 13 Siphon short (index 3-4), slightly swollen near middle *C (C) gelidus*, p 408
 Siphon longer (index at least 4.5, usually much more), not at all swollen near middle, but widest at base 14
- 14 Siphonal index 4.5-5, distal pecten-teeth strong and usually curved, *etc* with 4 fairly long branches *C (C) theileri*, p. 415
 Siphonal index at least 6, other characters not as above 15
- 15 Siphon long, narrow, slightly tapering on basal $\frac{1}{2}$, parallel-sided on apical $\frac{1}{2}$, index 8 or more, frontal hairs *B* and *C* 2-branched 16.
 Siphon shorter and wider, somewhat tapering from base to apex, index usually about 6-7, rarely more, frontal hair *C* 3-5-branched 22
16. Siphon with minute hair-tufts, index 11-14, *etc* single [p 350
 Siphon with moderate hair-tufts (about as long as diameter of tube), index 8-10.5, *etc* with 3-5 branches *C (N) brevipalpis*, 17
17. Siphon with a dark ring in middle 18
 Siphon without a dark ring in middle 19 [p 364.
- 18 Siphon with a dark ring also at base *C (L) minutissimus*,
 Siphon without a dark ring at base *C (L) fraudatrix*, [p 369.
19. Lateral hairs of abdominal segments I and II unusually stout *C (M) khazani*, 20 [p 357.
 Lateral hairs of abdomen normal *C (L) fraudatrix*, 21 [p 369.
20. Pecten of 8-12 teeth
 Pecten of about 16 teeth
21. Anal papillae more than twice length of segment; siphonal index 9-10 *C (L) minor*, p 371
 Anal papillae not much longer than segment, siphonal index about 10.5 *C (L.) mammillifer*, [p 375.
- 22 Siphonal tufts fully twice as long as diameter of tube, 8-10 in number (or 4-5 pairs), in addition to 2 pairs of short hairs near apex of tube, subapical bristles on antenna some distance from tip of shaft [p 411.
 Siphonal tuft scarcely longer than diameter of tube, subapical bristles on antenna near tip of shaft *C (C) mammetrus*, 23.
- 23 Siphon with 6 or 7 pairs of hair-tufts 24.
 Siphon with 4 or 5 pairs of hair-tufts 25
24. *th* of 4 very short branches, anal papillae about length of segment, ends bluntly pointed, pecten-teeth with 4-5 denticles, teeth of mentum regular [? *whites*, p. 402)
 [p 405 (and
C (C) tritaniorthynchus,
th of 3 short branches, anal papillae longer than segment, ends sharply pointed, pecten-teeth usually with 6 denticles *C (C) mimulus*, p 412.
- 25 Siphon with one pair of tufts, lateral in position, teeth of mentum irregular in size, antennal tuft at $\frac{2}{3}$; *etc* usually 3-branched *C. (C) barraudi*, p. 403.

- Siphon with 2 pairs of tufts, lateral in position, antennal tuft at $\frac{2}{3}$; *isc* 2-branched [p 420
C. (C.) univittatus, 27
 26 *isc* single 31
isc of 2-3 fairly long branches
 27 Posterior margin of chitinous ring of anal segment with short spines, siphon normal, tapering 28
 Posterior margin of chitinous ring of anal segment smooth 30
 28 Antennae moderate, tuft beyond middle, frontal hairs *B* and *C* 5-6-branched [p 386.
C. (L.) cinetellus?, 29
 Antennae long, tuft at or before middle, frontal hairs *B* and *C* 2-3-branched [p 379
 29 Siphon about 2.3 mm long *C. (Culex) viridiventer*,
 Siphon about 1.4 mm long *C. (Culex) shillibeae*,
 30. Siphon long and tapering, with a false joint beyond middle, due to lack of chitination in an irregular ring. [p 380
C. (Culex) tatus, p 384
 Siphon somewhat widened before middle, without false joint *C. (Culex) nigropunctatus*, p 381.
C. (Culex) pallidus.
 31. Siphonal tufts of 4-6 branches, index about 4, *lh* single *C. (C.) fatigans*, p 422
 Siphonal tufts of 2-5 branches, index 5-6 5; *lh* 2-branched 32.
 32. Mentum with about 11 teeth on either side of central one, *isc* 2-branched *C. (C.) vagans*, p 418
 Mentum with about 7 teeth on either side of central one; *isc* 3-branched [p 426.
C. (C.) fuscocephalus,

Subgenus LUTZIA Theobald, 1903.

M.C. III, p 155 Genotype, *L. bigoti* Theo

Jamesia Christophers, 1906, Sci Mem Med Ind. p 12. Genotype, *C. fuscans* Wied (as *C. concolor*)


ADULT—Large mosquitoes, distinguished from those in other subgenera of *Culex* by the presence of 4 or more lower mesepimeral bristles, and from all other genera by the presence of well-developed pulvilli. Palpi of ♂ longer than proboscis, last two segments upturned and hairy, of ♀ not more than $\frac{1}{2}$ length of proboscis. Wings with c-v's 2-3, 3-4, and 4-5 approximated and sometimes in straight line. Fore and mid-femora and tibiae mottled with spots of pale scales. Coxite of ♂ hypopygium without scales, but with subapical lobe bearing 3-5 stout spines. Style rather short, simple, curved, with small terminal appendage. Phallosome divided into lateral plates, with long pointed apex (ventral cornu) and some close-set teeth below this. Paraprocts with numerous spines at crown, a few minute hairs just below, and a projecting ventro-lateral arm.

The bucco-pharyngeal armature of ♀ resembles that of some species of subgenus *Culex*, the teeth being comparatively

short and blunt, and in some preparations the tips of the teeth appear to be thin and spoon-shaped (fig 80, a)

LARVA—Mouth-parts modified for predacity, mouth-brushes of a moderate number of strong curved rods, with minute hairs along one side on apical $\frac{1}{2}$. Frontal hairs single. Siphon short, with a postero-ventral row of hair-tufts and with pecten, both extending along whole length of tube. Anal segment large, longer dorsally than ventrally.

Fig 81




♂ hypopygial structures of *Culex*, subgenus *Lutzia*: a (seven upper figures), *fuscus*; b, *raptor*; c, *vorax*; d, *halifazr*. *ecb*, *db*, and *rb*, external, dorsal, and internal border of coxite, *bp*, *mp*, and *vp*, basal process, median processes, and ventral cornu of lateral plate of phalosome, *vlp*, *dp*, ventro-lateral and dorsal plates of proctiger. Other lettering as on p. 4.

DISTRIBUTION and BIONOMICS —All the four known Oriental species occur in India, three being fairly common, and one of these is found up to about 6,000' in the hills. The larvæ live in open pools of various kinds, in shallow wells, and in domestic collections of water, where they are entirely predaceous and devour the larvæ of other mosquitoes, chiefly

Culex fatigans The adult ♀♀ seldom attack man to any extent

One other species occurs in Africa, and four others in Central and South America

Fig 82



Larva of *Culex (Lutzia) halifax* a, siphon, b, anal segment; c, pecten-teeth, showing variation, d, head, e, mentum, f comb-teeth

Key to Adults*.

- | | | |
|---|---|---------------------------|
| 1 | Anterior surface of hind femur pale scaled on basal $\frac{1}{2}$ except dorsally, pale scaling on apical $\frac{1}{2}$ forming a line to the tip | 2 |
| | Anterior surface of hind femur with numerous dark scales on basal $\frac{1}{2}$, apical $\frac{1}{2}$ without a distinct line of pale scaling to the tip | 3 |
| 2 | Abdominal tergites all with whitish or ochreous apical bands, those on last few broader than those on anterior segments | <i>raptor</i> , p. 348. |
| | Abdominal tergites V-VIII entirely yellow scaled, or with broad apical bands, II-IV entirely dark, or with very narrow apical bands | <i>fuscatus</i> , p. 341. |
| 3 | Abdominal tergites all with pale apical bands of about the same width | <i>vorax</i> , p. 344. |
| | Abdominal tergites entirely dark, or with pale lateral markings only on the last few segments, or with very narrow pale apical bands | <i>halifax</i> , p. 344. |

* All four Indian species are very similar in markings, and some specimens are difficult to identify with certainty. The larvae appear to be indistinguishable. For these reasons it has been thought that they represent different forms of one species, but there are some differences in the form of the lateral plate of the phallosome of the ♂♂ which seem to be constant.

195. *Culex (Lutizia) fuscianus* Wiedemann, 1820*.

Dipt. Exot 1, p 9 (♀) [?]. TYPE-LOC 'Ind. or' TYPE non-existent (formerly in Vienna Mus)

Culex concolor Theobald, 1901, M C 11, p 107 (? *C concolor* Robineau-Desvoidy, 1825, Mem Soc. Hist. Nat. Paris, iv, p 405 TYPE-LOC. unknown TYPE: non-existent).

Culex schulosus Doleschall, 1857, Nat. Tijd Ned.-Ind. xiv, p 384 (♀). TYPE-LOC Java TYPE: non-existent (formerly in Vienna Mus)

? *Culex luridus* id., ib p 384 (♀) TYPE-LOC Java TYPE: ? non-existent.

ADULT ♂ —Wing about 5 mm

♀ —*Head* covered with narrow pale yellow scales and numerous upright scales, which appear dark from most angles, some broad white scales low down at each side. Antennae brown, tori and first flagellar segment with a few small pale scales. Clypeus and palpi dark brown, latter speckled with numerous pale yellow scales. Proboscis with wide pale yellow area in middle on both upper and undersides, variable in extent, otherwise dark brown. Palpi about $\frac{1}{2}$ length of proboscis. *Thorax* mesonotum covered with dark brown narrow scales, with lighter scales forming indefinite pattern of faint lines and spots, usually a pale spot each side at level of anterior spiracle, another pair of spots on dorsum in front of level of wing-roots, and a pair of faint lines from these to lateral lobes of scutellum, a faintly indicated median line anteriorly. Narrow, pale yellow, rather sparse scales on scutellum, *apm*, and posterior part of *ppn*; some darker scales on anterior part of *ppn*. Pleurae with few scales, some broad pale scales on upper part of sternopleura and mesepimeron, another small collection on lower part of sternopleura opposite lower corner of mesepimeron. *Wings*: dark scaled, c-vs 2-3, 3-4, and 4-5 sometimes one below the other in straight line, but position of 4-5 variable, either slightly nearer base of wing or slightly nearer apex than other two. *Legs* fore and mid-femora, when seen from the front, dark brown, plentifully sprinkled with pale scales, when seen from behind almost entirely pale yellow, hind femur on outer side pale on basal $\frac{1}{2}$ except dorsally, with pale scaling forming a more or less definite line to tip, on inner side almost entirely pale. Tibiae extensively speckled with pale scaling, tarsi brown or pale yellow, according to angle of view. No definite pale rings. *Abdomen*. markings variable, tergites II-IV either entirely

* The first part of Wiedemann's work was published in 1820, as recently shown by England (Ann. Mag. Nat. Hist. (10) viii, 1931, p. 613).

† Theobald 1901 b, p. 107 (*concolor*), Leicester 1908, p. 154 (*concolor*); Edwards 1922 c, p. 275 (syn.), Barraud 1924 c, p. 973, Brug 1924 b, p. 25.

dark brown, or with narrow yellow apical bands, and with or without a pair of submedian yellow spots. V with broad yellow apical band, VI-VIII usually entirely yellow scaled.

♂.—Antennæ densely plumose. Palpi longer than proboscis by rather more than length of apical segment; apical $\frac{1}{2}$, or more, of palp with long hairs projecting laterally from each side, hairs dark, except at tip of penultimate segment and along apical segment, where they are yellowish, scaling brown except on apical segment, where it appears mainly yellow, white scales on underside at joints between segments. Proboscis with pseudo-joint rather beyond middle, where there is a fairly narrow ill-defined pale ring. Otherwise scaling is dark brown. Other markings as in ♀ *Hypopygium* (fig. 81, a). subapical lobe of coxite with usually a fourth spine, more or less developed, and separate from the other three. *lp* with about 4 rather long teeth on the side of the ventral cornu, differing from the toothed median process present in the other three species.

LARVA*—*Head* moderately large. Antenna short, shaft smooth, with a single short hair at about $\frac{1}{3}$ from base; other hairs and spines all near tip of shaft. Mouth-brushes of a moderate number of strong curved rods, each with minute hairs along one side on apical $\frac{1}{2}$. *Frontal hairs* A, B, and C long and usually single, but may be split into two; A some little distance posterior to base of antenna; B and C placed far back, C being posterior to e, B about level with e, d anterior and internal to B, small and with several fine branches, e fairly long and usually single. Each individual pair of hairs widely spaced. Pre-clypeal spines inconspicuous, but fairly long, widely separated, and arising from small tubercles, external to each is a small "anterior frontal hair," as usual. Larger hairs on dorsum of *thorax* arising from small chitinated plates, lateral hairs arising from large tubercles, some of which have small thorn-like spines. *Abdominal segments* I-VI with projecting lateral tubercles carrying branched hairs. Comb of 35-45 fringed teeth on segment VIII. *Siphon* short and broad, with large acus at base and large valves at apex, from 1.2-2.5 times length of diameter at base. Pecten extending along length of siphon, with 7-11 teeth, the majority of which have a single lateral denticle, but the two or three most distal teeth simple. Posterior surface of siphon with a zigzag row of long hairs from near base to apex, one pair of smaller hairs anterior to pecten towards apex of siphon. *Anal segment* large, triangular in side view,

* Christophers 1906, p. 10 (*concolor*). Barraud 1924 d, p. 977. Senior-White 1927, pl. vi.

and usually longer than siphon Both *osc* and *isc* single and long, *lh* single, fairly long Fan of about 14 branched hairs arising from fan-plate Both pairs of papillae small

HABITAT.—Natural pools, shallow wells, domestic collections of water, etc

DISTRIBUTION.—Common over larger part of Indian region from the PUNJAB to ASSAM, BURMA, CEYLON, and ANDAMANS It has been found up to about 5,000' in the Assam hills, but is commonest in the plains

Known from MALAY PENINSULA, SIAM, COCHIN CHINA, DUTCH EAST INDIES, PHILIPPINES, and SOUTH CHINA

196 *Culex (Lutzia) raptor* Edwards, 1922.

Ind Journ Med Res x, p 275 (♂ & ♀) TYPE-LOC Ainitisar, Punjab, v. 1921 (*Barraud*) TYPE ♂ & ♀ in Brit Mus

ADULT ♀.—Very similar to *C (L) fuscans* but differing as indicated in key, e-v 4-5 usually much nearer base of wing than 3-4, but position variable Mesonotal scaling also subject to variation, in some specimens the scales are uniformly light brown without any indication of a pattern of pale scales, in others the scaling is mostly dark brown, with a similar pattern to that of *fuscans*, but more distinct. Some specimens, in which the abdominal banding is yellower than usual, are difficult to separate from certain specimens of *fuscans*

♂—Apical $\frac{1}{3}$ of long segment of palpi and base of penultimate segment usually with paler scaling than in *fuscans*, but this is not always very marked *Hypopygium* (fig 81, b): subapical lobe of coxite with usually 3 strong spines, but a fourth sometimes present, *lp* with a pronounced median toothed process Ventral cornu, in side view, less tapering than in the other species, lateral process rounded

LARVA †.—Very similar to that of *fuscans* In the material available it has not been possible to find any constant points of difference

HABITAT.—As given for *fuscans*

DISTRIBUTION—Commonest on western side of India from the PUNJAB, through BOMBAY and MADRAS PRESIDENCIES to CEYLON, but occurs also in CENTRAL PROVINCES, ORISSA, BENGAL, and BURMA No specimens from Assam have been seen. It is often found in the same localities as *fuscans*

Recorded from COCHIN CHINA (*Borel*)

* This and *C. fuscans* sometimes approach one another closely in abdominal markings and venation, but the difference in structure of the ♂ phallosome appears to be constant: *vide* Barraud 1924 c, p. 974.

† Barraud 1924 d, p. 977.

197 *Culex (Lutzia) vorax* Edwards, 1921

Bull. Ent. Res. xi, p. 327 (♂ & ♀) TYPE-LOC. Tokyo, Japan (Yamada). TYPE: ♂ & ♀ in Brit. Mus.

ADULT †—Fairly distinct in abdominal markings from the other three species, and differs from *C. (L.) fuscans* and *raptor* also in markings of hind femur, as indicated in key, c-v 4-5 generally nearer apex of wing than 3-4, but position variable, as in other species. Scaling of mesonotum variable and very similar to that of *raptor*. There is also some variation in width of pale apical bands on abdomen, and some specimens approach *halifaxi* in appearance. Palpi of ♂ with white scaling at bases of last two segments on upper side, otherwise scaling and hairs are mainly dark brown, but appearance varies according to angle of view, the apical segment sometimes appearing almost entirely light, but not definitely yellow as in *fuscans*.

♂—*hypopygium* (fig. 81, e) subapical lobe of coxite usually with only 3 strong bristles. Ventral cornu of *lp* tapering and pointed, and with a pronounced toothed median process, lateral process pointed.

LARVA †—Very similar to those of *fuscans* and *raptor*. In material available it has not been possible to discover any constant differences.

HABITAT—As given for *fuscans*.

DISTRIBUTION—Occurs almost exclusively in mountainous localities in India. WESTERN HIMALAYAS: Kasauli hills*, common during rains (Christophers & Barraud), Bhowali*, Naini Tal dist. (Rao), Mussooree, Dehra Dun dist., x. 1906 (Lefroy) all 4-7,000'. EASTERN HIMALAYAS: Kurseong* and Mungpoo*, Darjeeling dist., 5-6,000', ix. & x. 1922 (Barraud). ASSAM HILLS: Shillong* (Jukes, Shortt, Fletcher, Barraud), Cherrapunji* (Fletcher), Haflong*, Cachar Hills, viii. 1922 (Barraud), Mikir Hills*, Sibsagar dist., i. 1911 (Christophers) all 4-5,000'. BURMA: Kalaw*, Southern Shan States (Christophers). SOUTH INDIA: Coimbatore*, Nilgiri Hills, viii. 1928 (Shaffi).

Known also from HONG KONG and JAPAN.

198 *Culex (Lutzia) halifaxi* Theobald, 1903

M.C. iii, p. 231 (♀) TYPE-LOC. Dindings, Straits Settlements. TYPE: ♀ in Brit. Mus.

Culex multumaculosus Leicester, 1908, Cul. Malaya, p. 155 (♂ & ♀). TYPE-LOC. Malay Penin. (Leicester). TYPE: ♂ & ♀ in Brit. Mus.

Culex aureopunctus Ludlow, 1910, Can. Ent. xlii, p. 195 (♀). TYPE-LOC. Mindanao, Philippine Is. (Eastman). TYPE: ♂ in U.S. Nat. Mus.

† Barraud 1924 c, p. 975, Martini 1931, p. 348.

† Barraud 1924 d, p. 977.

ADULT †—Very similar to *C. (L.) vorax*, but differs in abdominal markings as indicated in key, and the pale scales on outer side of hind femur are usually less numerous. It is a darker species than the other three. In some specimens the mesonotum is covered with brownish-black scales, with paler scales forming a very distinct pattern, as in some specimens of *raptor*. The dorsum of the abdomen may be entirely brownish-black, or there may be narrow pale ochreous or white apical bands, much as in some forms of *vorax*. Palpi of ♂ very similar to those of last-named species.

♂—*Hypopygium* (fig. 81, d) subapical lobe of coxite with 3–5 strong spines. *lp* with very pronounced median toothed process and a rounded lateral process.

LARVA (fig. 82) †—Very similar to those of other species, no definite differences can be given.

HABITAT—Rice-fields (Assam, *Barraud*), street gully-traps, disused ships' tanks, garden water-barrels, and other water-holding rubbish about habitations, and rarely in fresh tea-tree swamp, nearly always with *Culex fatigans* (Brisbane, Australia, *Cooling*), pool of dirty water in ravine (Suduganga, Ceylon, *Senor-White*), small ponds, road-side ditches, and jungle-pool (Malay Peninsula, *Leicester*).

DISTRIBUTION—ASSAM, Nongpoh*, Khasi Hills dist., vii 1922 (*Barraud*). CEYLON, Suduganga, Matale dist., iii vi, x.–xii 1919, and i 1920 (*Senor-White*).

Known from MALAY PENINSULA, eastwards to AUSTRALIA, PHILIPPINES, and SOUTH CHINA.

Subgenus **BARRAUDIUS** Edwards, 1921.

Bull. Ent. Res. xii, p. 322. Genotype, *C. pusillus* Macq.

ADULT.—Small mosquitoes without conspicuous ornamentation. Segment I of hind tarsi distinctly shorter than tibia. Pulvilli well developed. Scales on vertex of head, *apn*, and scutellum all narrow. Palpi of ♂ rather longer than proboscis, slender, and without projecting hairs. Coxite of ♂ hypopygium with scales on outer side, lobe on coxite some distance from apex, with two projections, one carrying a single strong spine, the other several similar spines, but no leaflet. Style simple, with small terminal appendage. Phallosome divided into lateral plates, each with simple cornu, without teeth or processes. Paraproct with hairs or spines at crown, no lateral arm.

† Edwards 1913 b, p. 234, 1922 c, p. 275, 1922 d, p. 470 (syn.), 1924, p. 391, *Barraud* 1924 c, p. 975, *Brug* 1924 b, p. 26, *Pame* 1929, p. 309.

† *Barraud* 1924 d, p. 977, *Cooling* 1924, p. 14, *Hill* 1925, pl. viii [figured here from specimens in British Museum from Solomon Is.]

LARVA.—Mouth-parts normal, siphon short, with a single zigzag row of hairs on posterior border, much as in *Lutzia*.

DISTRIBUTION and BIONOMICS.—Only two species of this subgenus are known, both being found in the Mediterranean region, one extends eastwards to Kashmir, where it is fairly common. Little appears to be known of the habits of the adults, but *O. (B.) modestus* has been caught when biting man. The larvæ live in pools and swamps.

199 *Culex (Barraudius) modestus* Ficalbi, 1890.


Bull. Soc. Ital. xxi, p. 93 (♀). TYPE-LOC. Ravenna, Italy.
TYPE. Pisa (?).

Culex eadithæ Barraud, 1924, Ind. Journ. Med. Res. vi, p. 1004
(♂). TYPE-LOC. Sopor Village, Jhelum River, Kashmir,
2,200', x 1923 (Senton). TYPE. ♂ in Brit. Mus.

ADULT *.—Rather small species, wing (♂) 3-3.5 mm.,
(♀) 4-4.5 mm.

♀.—Head covered with brown or yellowish narrow scales, some broader pale ones low down on each side, and fairly numerous upright scales on nape. Tori yellowish-brown;

Fig. 83



♂ hypopygial structures of *Culex (Barraudius) modestus* Fic.
(end of coxite, with style; proctiger; phallosome)

flagellum of antenna, clypeus, palpi, and proboscis dark brown, the last pale beneath and usually at sides for whole length, palpi about $\frac{1}{2}$ length of proboscis. *Thorax*: integument of mesonotum brown, uniformly covered with chestnut-brown scales; scutellar scales rather lighter. Pleuræ pale brown, with a few small flat pale scales on sternopleura and upper part of mesepimeron. One lower mesepimeral bristle. *Wings* dark scaled. *Legs* dark brown; fore and mid-

* Ficalbi 1899, p. 211, Edwards 1921 c, p. 332; 1926 b, p. 624;
Martini 1931, p. 357

femora lighter posteriorly, hind femur pale, except dorsally, on both sides, segment 1 of hind tarsi shorter than tibia. *Abdomen*. Dorsum dark brown, lateral rectangular pale ochreous patches on tergites, forming a continuous pale border on either side of abdomen. Sternites uniformly pale ochreous.

♂—Markings very similar to those of ♀. Palpi slender, longer than proboscis by a little less than length of apical segment, dorsally dark brown, underside pale, except for the dark apical segment, underside of proboscis also pale, no hair-tufts. *Hypopygium*: *vide* remarks under subgenus, and fig 83.

LARVA *—Has not been isolated in India. The following details are taken from figures and brief description by Gil Collado. The larva appears to be very similar to that of *C. (B.) pusillus*, described and figured by Edwards (1921, p. 333). *Antenna* with spicules on shaft, and hair-tuft at nearly $\frac{3}{4}$ from base, part of shaft beyond tuft darkened, 3 long spines at tip. Mouth-brushes normal. *Frontal hair A* of about 8 branches, *B* and *C* at about middle of clypeus, *C* being slightly internal and posterior to *B*, both 3-branched. Numerous comb-teeth on abdominal segment VIII in a patch, individual teeth with rounded fringed apex. *Siphon* with a slightly zigzag row of branched hairs along posterior border from near base to apex. Pecten of about 12 teeth, the most distal one with about 5 lateral denticles along basal $\frac{1}{2}$. *Anal segment* figured (by Gil Collado) as having a small chitinous saddle†. Both pairs of papillae short, fan not large.

HABITAT—Ground-pools.

DISTRIBUTION.—Fairly common in the Jhelum Valley, KASHMIR (*Simton, Fletcher, Barraud*). Not known from other parts of Indian region, but occurs in ASIA MINOR and MEDITERRANEAN REGION as far west as SPAIN.

Subgenus NEOCULEX Dyar, 1905.

Proc. Ent. Soc. Wash. vii, p. 45. Genotype, *C. apicalis* Adams (as *C. territans* Walk.).

Protomelanoconion Theobald, 1910, M.C. v. p. 462. Genotype, *P. fuscum* Theo.

ADULT (Indian species) ‡—Small mosquitoes with dark tarsi, and without marked ornamentation. Mesonotal bristles strong and numerous. Scales on scutellum and on vertex

* Gil Collado 1930, p. 342, Martini 1931, p. 358.

† [The figure may represent a 3rd-stage larva; in the related *C. pusillus*, as well as in most other species of *Oulex*, the anal segment in the 4th-stage larva has a complete chitinous ring.]

‡ [Most of the non-Indian species differ from the two recorded here in having abdominal tergites apically banded and ♂ palpi as long as proboscis or longer. The Indian species belong to the *Protomelanoconion* group.]

of head all narrow. Palpi of ♂ from $\frac{1}{2}$ to $\frac{5}{8}$ length of proboscis, those of ♀ quite short. Coxite of ♂ hypopygium without scales, but with subapical lobe bearing spine-like processes. Style simple, with small appendage. Phallosome divided into two lobes, upon which are numerous small spines or teeth.

Bucco-pharyngeal armature of ♀ with a row or palisade of very numerous pointed teeth (fig. 80, c).

LARVA.—Very similar to those of several other subgenera. Siphon long and narrow, with about 5 pairs of hair-tufts. Pecten-teeth with lateral denticles along one side from base to apex.

DISTRIBUTION and BIONOMICS.—Twenty-three species have been classed in this subgenus by Edwards, only two of these are at present known in India, the others being scattered over Europe, Africa, Asia, and Australia, one occurring in America. *C. (N.) brevipalpis* is fairly common in parts of India where there is heavy rainfall, but *C. (N.) tenuipalpis* is known only from one locality. The larvae of the latter were found in road-side pools, those of the former live in tree-holes, bamboos, etc. Very little is known of the habits of the adults of these two species, but Giles stated that the ♀ of *brevipalpis* bites during the day. Adults of this species have been caught in houses. [Several species of this subgenus have been found to suck the blood of frogs, and it has been suggested that this may be a common habit of species of this group of *Culex*.]

200 *Culex (Neoculex) brevipalpis* (Giles), 1902.

Handbook, 2nd ed. p. 384 (*Stegomyia*) (♂ & ♀). TYPE-LOC.: Shahjahanpur, U.P. (Giles). TYPE: ♂ & ♀ in Brit. Mus.

Culex longipes Theobald, 1901, M.C. II, p. 68 (♀) (name preocc.). TYPE-LOC.: Singapore, vii. 1899 (*Hantsch*). TYPE: ♀ in Brit. Mus.

Culex macropus Blanchard, 1905, Les Moustiques, p. 327 (nom. nov. for *C. longipes*).

Melanocomia uniformis Leicester, 1908, Cul. Malaya, p. 136 (♂ & ♀). TYPE-LOC.: Malay Penin. (Leicester). TYPE: ♂ & ♀ in Brit. Mus.

Culex fidelis Dyar, 1920, Insec. Inscit. Mens. viii, p. 180 (♂ & ♀). TYPE-LOC.: Los Baños, Philippine Is., vii. 1915 (*Baker*). TYPE: 3 ♂♂ in U.S. Nat. Mus. *, also 2 allotype ♀♀, ix. & xi. 1917, same locality.

ADULT †.—Small dark species (wing about 2.5–3 mm.) characterized by peculiar form of ♂ palpi (fig. 84, b).


♀.—Head. narrow pale scales and upright yellow scales on vertex and nape, the upright scales numerous. Antenna,

* Edwards 1929 a, p. 4.

† Edwards 1913 b, p. 237 (syn.), 1922 d, p. 472 (syn.), 1929 a, p. 4; Barraud 1924 h, p. 1277; Borel 1926, p. 114.

palpi, and proboscis dark brown, palpi about $\frac{1}{2}$ length of proboscis. *Thorax* integument of mesonotum brown, a pair of submedian darker lines, a green tinge at sides; scales narrow and dark brown, lighter on front margin. Two rows of long black bristles from front margin of mesonotum to lateral lobes of scutellum, others over wing-roots, giving thorax a markedly bristly appearance. Scutellar scales narrow, dark brown. Pleuræ pale green or pale ochreous, *ppn* without scales, but with 3-4 bristles and some hairs, *apn* with very few pale scales anteriorly, a small patch of broad white scales on sternopleura, no lower mesepimeral bristle. *Wings* dark scaled. *Legs* unusually long, dark brown, with bronzy sheen, femora paler beneath, as usual,

Fig 84



Structural details (♂ hypopygium and palpi) of *Oulex*, subgenus *Neoculex*: a, *tenuipalpis*, b, *biviripalpis*


especially hind pair. *Abdomen* dorsum dark brown, with bronzy sheen; venter pale green, creamy on terminal segments

♂.—Very similar to ♀. Palpi (fig. 84, b) dark brown, about $\frac{1}{2}$ — $\frac{3}{4}$ length of proboscis (may appear shorter in dried specimens, as terminal segments often become distorted), long segment curved outwards at tip, with some short erect hairs externally and a few strong bristly hairs at tip, penultimate segment fairly long, curved (convexity above), with similar hairs, terminal segment short, with a fine strong bristle at tip. *Hypopygium* (fig. 84, b) subapical lobe of coxite with a leaflet and about 6 strong spines, 3 long and 3 shorter. Phallosome divided into two broadly conical lobes, with minute spines or teeth on apical part. Paraproct with spines at crown; no lateral arm

LARVA * (fig. 85, b)—Remarkable for the extreme length and narrowness of the siphon, which is 12–14 times as long as diameter at base, and for the reduction in size of the branched hairs on siphon

Antenna slightly curved, shaft widest at about middle, with fairly numerous small spinelets along its length; a many-branched tuft of subplumose hairs, on outer side, at about $\frac{3}{4}$ of length from base, 3 long bristles at tip, and 1 very short. *Frontal hairs* A, B, and C with subplumose branches, the number of which is, respectively, 8–10, 2, 2

Fig. 85



Larval structures of *Culex*, subgenus *Neoculex* ·
a, *tenuipalpis*, b, *brevipalpis*.

Hair *d* small, internal and anterior to B and C. Mouth-brush hairs simple; maxillæ very prominent. Pre-clypeal spines strong and of moderate length. Mentum with 8–10 teeth either side of unusually large central tooth. Some very long single hairs on *thorax*, and branched, barbed hairs, arising from tubercles, on meso- and metathorax laterally. *Abdomen*: long single and 2-branched hairs on I and II, those on following segments smaller, except for a very long single hair on either side of VII. Comb a triangular patch of about

* Brug 1924 a, p 440, Barraud 1924 m, p. 432, Borel 1926, p. 116; Senior-White 1927, p 71.

50 small teeth, each tooth with swollen fringed apex. Sub-siphonal tuft with 6-8 rather long subplumose branches. Branched hairs on *siphon* minute, usually 5 pairs, about evenly spaced, along length of tube beyond pecten; latter with 14-18 teeth, each tooth with lateral denticles along one side from base to apex. *Anal segment* enclosed in chitinous ring. Both *osc* and *isc* single and long, $\frac{1}{2}$ small, divided into several branches towards tip. Dorsal papillæ about length of segment, ventral pair shorter, but length rather variable. Fan rather short, about 12 hairs, each divided into a number of branches, attached to fan-plate.

HABITAT—Tree-holes, bamboos, once in open iron tank (*Barraud*)

DISTRIBUTION—Common over larger part of Indian region east of a line drawn from the western boundary of Nepal to Bombay. A number of other Indian mosquitoes have a very similar distribution.

Known also from MALAY PENINSULA, SIAM, COCHIN CHINA, and MALAY ARCHIPALAGO.

201. *Culex (Neoculex) tenuipalpis* Barraud, 1924.

Ind. Journ. Med. Res. xi, p. 1278 (♂ & ♀). TYPE-LOC. Sural, near Mungpoo, Darjeeling dist., 5,500', x 1922 (*Barraud*)
TYPE ♂ & ♀ in Brit. Mus.

ADULT.—Distinguished from *C. (N.) brevipalpis* by dark pleuræ and form of ♂ palpi.*

♀.—**Head** pale brown narrow and upright scales on vertex and nape. Antennæ, palpi, and proboscis dark brown. Palpi $\frac{1}{3}$ — $\frac{1}{2}$ length of proboscis. **Thorax**: mesonotal scales dark brown, bristles fairly numerous, especially over wing-roots, but less conspicuous than in *C. (N.) brevipalpis*. Scutellar scales narrow and brown. Pleuræ without scales, dark brown, except in region of coxæ and under wing-root, where colour is pale green, *ppn* with 4-6 bristles, 1 lower mesepimeral bristle. **Wings**: dark scaled. **Legs**: fairly long, tibiae and tarsi dark brown; fore and mid-femora pale posteriorly and ventrally; hind femur white on both surfaces, except dorsally towards extremity. **Abdomen**:

* [Species rather closely resembling *C. tenuipalpis* are *C. hayashii* Yam. (Japan and China), with ♂ palpi $\frac{1}{2}$ proboscis, and *C. sumatranus* Brug. (Sumatra and Hong Kong), with ♂ palpi only $\frac{1}{3}$ proboscis. Some species of the subgenus *Lophoceratomyia* are also very similar, especially *jenseni* de Meij. and *curtipalpis* Edw. All these differ in hypopygial structure and in the larva; from recent work by Dr R. B. Jackson in Hong Kong it would seem that there has been some confusion in associating the larvae and adults of some of these species, as he finds that a larva identical with that described by Edwards and Given as *curtipalpis* produces adults apparently indistinguishable from *sumatranus*, whereas Brug. attributed a totally different larva to *sumatranus*.]

dorsum dark brown, a fringe of pale hairs along apical margin of each tergite, and usually narrow basal pale bands. Venter brown, paler at base of each segment.

♂.—Very similar to ♀. Palpi about $\frac{1}{2}$ length of proboscis, thin and straight, without hair-tufts, terminal segment very small (fig. 84, a). Antennæ plumose. *Hypopygium* (fig. 84, a) subapical lobe of coxite with a number of rod-like spines, but no leaflet. Lobes of phallosome of slightly different shape to those of *brevipalpis*, and with larger spines or teeth.

LARVA * (fig. 85, a).—Recognized by the unusual form and arrangement of frontal hairs. *Antenna* comparatively long, dark at base and on apical $\frac{1}{2}$, a fairly large hair-tuft at about $\frac{1}{2}$ from base, small spinelets on more than basal $\frac{1}{2}$ of shaft, 1 short and 3 long bristles at tip. *Frontal hair* *A* of about 6 long branches, *B* of 2 branches, shorter than *A*, *C* single, very short, standing behind *B*, *d* small, single, or split towards tip, standing internal and anterior to *B*. Preclypeal spines dark and stout. Mentum subtriangular, about 7 teeth either side of central one. Some long single hairs on *thorax*, and branched hairs arising from tubercles on meso- and metathorax laterally, as usual. *Abdominal* segment I with tubercles laterally, carrying 2 long hairs, one single, one 2-branched, a similar 2-branched hair on II, following segments with finer 2-branched hairs, but those on VII very long. Subsiphonal tuft and the 4 associated hairs all fairly well developed. Comb of 40–45 rather long, fringed teeth in triangular patch. *Siphon* pale in colour, widest at base, tapering to tip and slightly curved, 6–7 times length of diameter at base, 5 pairs of hair-tufts along posterior border, about evenly spaced beyond pecten, latter of 9–10 long teeth, with numerous denticles along one side from base to apex, those on apical $\frac{1}{2}$ very small. Tracheæ within siphon very narrow. *Anal segment*, with its hairs and papillæ, as shown in fig. 85, a.

HABITAT.—Road-side pools.

DISTRIBUTION.—Known in INDIA only from type-locality, but also recorded from JAVA †

Subgenus **MOCHTHOGENES** Edwards, 1930.

Bull. Ent. Res. xxi, p. 305. Genotype, *O. malayi* (Leac.)


ADULT.—Essentially as given for *Neoculex*, but palpi quite short in ♂ as well as ♀, and there are rather numerous flat scales on the head in two species. Subapical lobe of coxite of ♂ hypopygium with several spines, rod-like processes,

* Barraud 1924 m, p. 433.

† Brug 1931, p. 38.

and one or more leaflets, the latter very large in two species in which also the style is of unusual form. The Indian species are all small and dark coloured, without ornamentation, and the ♀♀ are difficult to separate from those of some of the smaller species in other subgenera. Bucco-pharyngeal armature of ♀ as in *Neoculex* in some species (*C. iphis*, in which the teeth are numerous and long), but in other species the teeth are shorter (*C. khazani*)

Fig 86



♂ hypopygial structures of *Culex*, subgenus *Mochthogenes* (style, appendages of coxite, phallosome). a, *khazani*, b, *iphis*, c, *castrensis*; d, *plumalis*, e, *malayi*

LARVA.—As in *Neoculex* (so far as known).

DISTRIBUTION and BIONOMICS.—Nine species are at present known, of which one is confined to Africa, five occur in India, and the remainder in Malaya or islands to the east of the Malay Archipelago. So far as is known, the larvæ live in tree-holes, small rock-pools, and sometimes in ground-pools. Nothing appears to be known regarding habits of the adults.

Key to Adults

1. Integument of pleura with a dark spot occupying larger part of mesepimeron . . . *khazani*, p. 357.
 Integument of pleura without such spot (though in *malayi* and *castrensis* there is a dark area across upper part) . . . 2
2. Head mainly flat-scaled, style of ♂ hypopygium forked (fig. 86, e) . . . *malayi*, p. 358.
 Head with large area of flat scales on each side, continued in front as a border to eyes nearly to middle point, some narrow and upright scales on vertex and nape, style of ♂ hypopygium not forked . . . *pluvialis*, p. 350.
 Head mainly covered with narrow and upright scales, flat scales confined to a small area low down each side, style of ♂ hypopygium not forked . . . 3
3. Bristles on vertex pale, a dark area across upper part of pleura, very small species, style of ♂ hypopygium much swollen in middle, lobes of phallosome elongate and without distinct spines or teeth (fig. 86, c) . . . *castrensis*, p. 355.
 Bristles on vertex dark, pleura uniformly dark brown, larger species, style of ♂ hypopygium not swollen in middle, but with enlarged tip (fig. 86, b), lobes of phallosome wide and short, and with a row of small teeth . . . *iphis*, p. 354.

202. *Culex* (*Moethogenes*) *iphis* Barraud, 1924.

Ind. Journ. Med. Res. xi, p. 1279 (♂ & ♀) TYPE-LOC. Nilgiri Hills, South India, ix. 1915 (*Khazan Chand*) TYPE: ♂ & ♀ in Brit. Mus.

ADULT—Size rather larger than other species of this subgenus. Wing 3.4–3.8 mm.

♀—*Head* narrow dull white and dark upright scales on vertex. Palpi and proboscis dark brown. Length of proboscis 2.2–2.4 mm, of palpi about 0.35 mm. *Thorax*: mesonotal scales dark brown, lighter on front margin: a number of long dark bristles, especially over wing-roots. Scutellar scales narrow and brown. Pleurae dark brown, without scales or definite markings; one lower mesepimeral bristle. *Wings*: dark scaled. *Legs* dark brown; hind femur pale on basal $\frac{1}{3}$ or more. *Abdomen*: dark brown dorsally, paler ventrally.

♂.—Antennae only moderately plumose, appearing less hairy than in the other species. Proboscis 2.2–2.4 mm. long, palpi about 0.38 mm., apical segment longer than penultimate. Other details as in ♀. *Hypopygium* (fig. 86, b): subapical lobe of coxite with one leaflet and 3 strong curved

spines Style somewhat widened at tip Lobes of phallosome with only a single row of minute spines or teeth

LARVA.—Unknown

DISTRIBUTION —Known only from type-locality

203 *Culex (Mochthogenes) castrensis* Edwards, 1922

Ind Journ Med Res x, p 285 (♂ & ♀) TYPE-LOC · Castle Rock, North Kanara (*James*) TYPE ♂ & ♀ in Brit. Mus

Aedes nigrescens Theobald, 1907, M C iv. p 540 (in part) TYPE-LOC and TYPE as above

As I have not seen any specimens of this species, I quote Edwards's original description * —“ A small dark species, almost devoid of ornamentation Dorsal surface of head apparently covered with small, whitish, narrow, curved scales, perhaps some flat ones also, but these difficult to make out owing to the poor condition Vertical bristles whitish, but the orbital bristles dark as usual Palpi alike in the two sexes, slender, exceeding the clypeus by about twice its length Mesonotum with dark narrow scales, some white ones round the front margin Pleuræ damaged, but apparently with dark markings a dark stripe across the upper part, and a dark patch lower down, no conspicuous white scales Abdomen dark above, paler below Male hypopygium · lobe of side-piece with two long rods, hooked at tips, and several undifferentiated bristles, apparently no leaf Clasper entire, but much swollen in the middle Lobes of mesosome elongate, nearly straight, rounded at tips, without small spines † Tenth sternites with crown of spines, no basal arm

“ The specimens on which this description is based are those which were labelled by Theobald as the type ♂ and ♀ of his *Aedes nigrescens*, from Castle Rock, India (*James*) The name *nigrescens* must be considered as preoccupied by *Danielsia nigrescens* Theo, described earlier in the same volume, which is also a species of *Culex* Moreover, it is certain that Theobald's *Aedes nigrescens* was described from specimens of two distinct species The male which he figured, and on which his conception of the species was mainly based, is an example of *C. malayi* Leic. The remaining specimens, which provided the material for his description of the head and thorax, are different, as is clearly shown by the scaling of the head (*C. malayi* having the dorsal surface of the head almost covered with small flat scales) and by the structure of the hypopygium, which is as described above. It was certainly a remarkable coincidence, amply sufficient to excuse the error, that Theobald should receive in the same collection,

* See also Barraud 1924 h, p 1280.

† [Actually there are several small tubercles.]

from the same place, two Oriental *Culex* with short male palpi, when no such species was known previously "

♂ —*Hypopygium* as in fig 86, c

LARVA* —[As figured by Senior-White this has some resemblance to *C* (*M*) *khazani*, but differs in many respects, as follows —*Frontal hairs* *A*, *B*, and *C* all simple (i.e., not plumose), *A* 3-4-branched, *B* and *C* 2-branched, but much shorter than in *khazani*, *B* not nearly reaching front edge of clypeus, *C* much shorter even than *B*, and placed almost directly behind *B*. *Comb* with fewer, shorter, and broader teeth. *Siphon* quite straight, very slightly and evenly tapering, barely 6 times as long as diameter at base. *Pecten* with the distal teeth almost 3 times as long as those at base. *Siphonal tufts* at least twice as long as diameter of tube.]

DISTRIBUTION —The typical form is known only from NORTH KANARA (type-locality), and from Suduganga and Marble, CEYLON (*Senior-White*) †

204 *Culex* (*Mochthogenes*) *pluvialis* Barraud, 1924

Ind Journ Med Res xi, p 1281 (♂ & ♀) TYPE-LOC Kairua, North Kanara, ix 1921 (*Barraud*) TYPE ♂ & ♀ in Brit Mus.

ADULT —Very similar to *khazani* and *malayi*, but differs as follows. —A large area of flat scales on each side of head, the flat scales continued as a border to eyes nearly to middle in front, but less numerous than in *malayi* and lighter in colour; vertex with narrow curved and upright scales

* Senior-White 1927, pl x

† [Brug (1932a, p 82) has described a var *foliatus*, from Java, which differs from the type-form in having the ♂ coxite "adorned with six fairly broad leaves instead of two rods and four bristles", the larva, according to Brug, agree closely with Senior-White's description, except that the comb-scales are broader (he should have said narrower, the figures he gives are accidentally reversed). One isolated skin and two whole larvae from Brug's material are in the British Museum, those agree with Senior-White's figure in all the points noted above except the form of the comb-scales, the lateral hairs of abdominal segments I and II are not noticeably thickened, this being another striking distinction from *khazani*.

This variety of the species occurs in Hong Kong, where larvae and adults were collected by Dr R. B. Jackson in 1932-3. The Hong Kong adults are closely similar to the South Indian types, but as they are in better condition it is possible to see that there is a narrow border of small flat scales adjoining the eyes in front. ♂ *hypopygium* with style formed as in the type, but coxite bearing one large twisted leaf, two long, bent, and flattened rods, and four or five short leaflets with pointed tips (this seems to be the structure in Brug's type also, his figure and description being rather incorrect; the specimen not being stained it is not very easy to make out fine details), *lp* with numerous tubercles. The Hong Kong larvae are in all respects similar to those from Sumatra, except that frontal hair *B* is very slightly plumose (much less so than in *khazani*); they were found in a shaded pool in the bed of a stream, in company with *Uranotaenia annandalei*]

Length of proboscis, ♀ 1.3–1.4 mm, ♂ 1.5 mm, length of palpi, ♀ 0.21, ♂ 0.4 mm, palpi of ♂ longer in proportion to those of ♀ than in *khazani*. Pleuræ dark, without a definite black spot on mesepimeron. One or two lower mesepimeral bristles. Wing, ♀ 3 mm., ♂ 2.3 mm.

♂—*Hypopygium* (fig. 86, d) subapical lobe of coxite with 3 leaf-like processes and 3 long, hooked, or clubbed spines. Style simple, narrow at apex, but not noticeably widened in middle, with terminal appendage *lp* pointed, with fairly numerous minute teeth or spines.

LARVA—Unknown.

HABITAT—Small rock-pools in jungle.

DISTRIBUTION—Known only from type-locality.

205 *Culex (Mochthogenes) khazani* Edwards, 1922

Ind Journ Med Res x, p 286 (♂ & ♀). TYPE-LOC Pudupach, Malabar Coast, xi 1915 (*Khazan Chand*). TYPE ♂ and allotype ♀ in Brit Mus.

ADULT*—A small, dark mosquito (wing about 2.5 mm), distinguished from allied species by presence of a black spot occupying larger part of mesepimeron.

♀—*Head* vertex and nape covered with narrow and upright scales, mostly pale in colour, some white and dark flat scales at sides. Antennæ, palpi, and proboscis brownish-black, proboscis slightly longer than fore femur, palpi about $\frac{1}{2}$ length of proboscis, latter often with appearance of pseudo-joint, or slight kink, at about $\frac{2}{3}$ from base. *Thorax*. Integument of mesonotum and scales brownish-black, fairly numerous and rather long black bristles, an elliptical darker area at sides over wing-roots. Pleuræ without scales, pale ochreous or greenish, a dark area occupying larger part of mesepimeron, one lower mesepimeral bristle. *Wings* dark scaled. *Legs* brownish-black, fore and mid-femora pale beneath, hind femur conspicuously white, except at tip and dorsally on distal $\frac{1}{3}$. *Abdomen* tergites brownish-black, sternites brown or pale brown.

♂—Very similar to ♀. Antennæ plumose. Palpi only very slightly longer than in ♀. *Hypopygium* (fig. 86, a): form of style and of processes arising from subapical lobe of coxite distinctive. Phallosome similar to that of *C. (N) tenuisulcus*, but teeth or spines more numerous and more variable in size.

LARVA—Not known with certainty, but some which were preserved from a batch, from which only adults of this species were reared, all have the following characters:—Length about 5–6 mm including siphon. *Head* moderately large. *Antenna* fairly long, almost uniformly dark, shaft with

* Edwards 1922 b, p 102, Barraud 1924 b, p 1281.

small spicules, a well-developed fan-shaped tuft at about $\frac{3}{4}$ from base, branches strongly plumose. Two long subapical bristles ($\frac{2}{3}$ length of antenna) very near tip of shaft, one apical bristle about $\frac{1}{2}$ length of subapical, and one quite small. *Frontal hairs* *A*, *B*, and *C* with 6-8, 2, and 3 branches respectively, but *B* and *C* sometimes single, branches of last two strongly plumose and long (reaching to tips of preclypeal spines or beyond), *C* distinctly internal to *B*, *d* fine and simple, fairly long, slightly anterior and internal to *B*; *e* with 4-5 fine branches. Mouth-brushes fairly large, hairs simple. Preclypeal spines stout and conspicuously dark. Some long single and 2-branched hairs on prothorax, and long branched hairs on meso- and metathorax laterally, some of the branches being very long. *Abdominal* segment I with a very long and strong, barbed, 2-branched hair, and one similar single hair; II with one similar single or 2-branched hair. Hairs on following segments much finer and shorter, but a long 2-3-branched hair on each side of VI, much longer than any hair on VII. Comb of very many long narrow teeth in a patch. Subsiphonal tuft of 5-8 subplumose branches. *Siphon* long and narrow, about 10 times length of diameter at base, slightly tapering on basal $\frac{1}{3}$, then almost parallel-sided to tip, with a slight but perceptible double curve near base. Pecten of 8-12 long teeth similar in form to those of *tenuipalpis*, distal teeth not very much larger than those at base, usually 6 pairs of very small branched hairs (not longer than diameter of tube), evenly spaced, along siphon. *Anal segment* very similar to that of *tenuipalpis*, but papillae shorter, hardly as long as segment.

HABITAT—Tree-holes

DISTRIBUTION.—MALABAR COAST. Pudupadi*, type-locality, as given above. NORTH KANARA Kadra*, ix 1921 (*Barraud*)

Not known from elsewhere

206. *Culex (Mochthogenes) malayi* (Leicester), 1908

Cul. Malaya, p. 184 (*Aedes*) (♂ & ♀). TYPE-LOC. Kuala Lumpur, Malay Penin (*Leicester*) TYPE ♂ & ♀ in Brit. Mus.

Aedes nigrescens Theobald, 1907 (in part), M C iv, p. 540 (nomo preocc) (♀). TYPE-LOC. Castle Rock, North Kanara (*James*). TYPE ♀ in Brit. Mus.

Anoretomyia aedes Leicester, 1908, Cul. Malaya, p. 180 (♂). TYPE-LOC. Kuala Lumpur, Malay Penin (*Leicester*) TYPE ♂ in Brit. Mus.

ADULT †—Very similar to allied species, but differs in head-scaling and coloration of pleurae. In some specimens there are narrow basal pale bands on abdomen. Wing 2-3 mm.

† Edwards 1917, p. 228 (*Micraëdes*, syn.), 1922, p. 286 (syn.); Barraud 1924 h, p. 1282, Borsal 1926, p. 117

♀—*Head* mainly covered with small light-coloured flat scales, a small area of narrow scales in middle of vertex; upright scales scattered over vertex and nape, the last appearing dark. Antennæ, palpi and proboscis brownish-black, palpi about $\frac{1}{8}$ length of proboscis. *Thorax*. mesonotum covered with brown narrow scales, bristles not very numerous or long. Pleura dark on upper $\frac{1}{2}$, greenish below, the two areas usually fairly sharply defined, 4-6 *ppn* bristles, one lower mesepimeral bristle. *Wings* dark scaled. *Legs*: deep brown, fore and mid-femora lighter posteriorly, hind femur mainly pale, but narrowly brown dorsally from near base to knee. *Abdomen* brownish-black dorsally, some specimens with narrow basal pale bands.

♂—Palpi about $\frac{1}{8}$ length of proboscis, antennæ plumose. Other details as in ♀. *Hypopygium* (fig 86, c). differs from that of all allied species in form of style, which is divided into two arms. Subapical lobe of coxite with a very large leaflet and several other processes.

LARVA *—From figures and a brief description of this larva by Borel from Cochín China it appears to be very similar to those of *tenuipalpis* and *khazani*, differing from the latter in having 5 fairly well-developed pairs of tufted hairs on the siphon, and from the former in having a smaller number of teeth in the comb (about 20 instead of 40 or more), the teeth also being more pointed. Siphon from 5-6 times length of diameter at base, pecten of 9-10 teeth, resembling those of *tenuipalpis* in form, distal teeth apparently not very long; tufts longer than diameter of tube.

DISTRIBUTION—Fairly common from the PUNJAB (Lahore, Karnal) to ASSAM, BURMA, and ANDAMANS, and through PENINSULAR INDIA to MADRAS. Probably, owing to its small size, it is frequently overlooked.

Known also from MALAY PENINSULA and ARCHIPELAGO as far south-east as TIMOR, and from COCHIN CHINA and SOUTH CHINA (Hangchow, *Feng-Swen Li*)

Subgenus LOPHOCERATOMYIA Theobald, 1905.

Ann Mus Nat Hung 31, p. 93. Genotype, *L. fraudatrix* Theo.

Phalodendromyia Theobald, 1907, MC 14, p. 623. Genotype,

P. barkeri Theo.


Cynithomyia de Meijere, 1910, Ann Jard Bot Buitenzorg (2) 31, p. 921. Genotype, *C. jenseni* de Meij.

ADULT—Small to medium-sized mosquitoes, brownish-black or reddish-brown in colour. Vertex of head usually with numerous small flat scales in front towards eye-margins. Proboscis and palpi dark brown, without pale rings. Scutellar

* Borel 1926, p. 118.

scales narrow. Lower mesepimeral bristle usually present. Tarsi entirely dark. Wing-scales dark and usually very scanty, except towards tip of wing. Palpi of ♂ (in known Indian species) a little longer than proboscis, and slender. Antennae of ♂ with a matted tuft of hairs on segment 9 (the

Fig 87




Modified segments (6-10) of ♂ antennae of *Culex*, subgenus *Lophoceros*: a, sensorium, b, mammillifer, c, fraudatrix, d, minor, e, minutissimus, f, cinctellus, g, plantaginis, h, flavicornis, i, uniformis.

torus being counted as segment 1) *, and usually with other matted or twisted tufts on segments 6-9 or 10. On segment 6 there are, in a number of species, scales or thickened hairs,

* Strictly speaking, the torus is the second segment

and the number and length of these are of diagnostic importance. The torus may, or may not, have a blunt prominence on inner side. ♂ hypopygium very similar in structure

Fig 88



♂ hypopygial structures (tip of coxite with style, phallosome from side or with proctiger) of *Culex*, subgenus *Lophoceratomyia*: a, m, *fraudatrix*; b, *flavicornis*; c, l, *mammulafer*; d, *seniori*; e, j, *minor*; f, o, *plantaginis*; g, *cinctellus*; h, n, *uniformis*; i, k, p, *minutissimus*.

to species of the subgenera *Neoculex* and *Mochthogenes*. Bucco-pharyngeal armature of ♀ with a concave row of long teeth (fig 80, d). So far as investigations have been

made there do not appear to be marked differences in the number and arrangement of teeth in closely allied species.

LARVA—Those of most species do not show any marked differences from those of other subgenera, but that of *uniformis* is remarkable in having the integument of the body covered with small hairs

DISTRIBUTION and BIONOMICS *—Several species are fairly common over large tracts of the Indian region, but are most numerous in forested regions of heavy rainfall. About 25 species are known in all, and these are confined to the Oriental and Austroalasian regions. Ten occur in India, these mostly breed in tree-holes, but the larvae of some live in ground-pools, shallow wells, etc. The adults do not appear to attack man, but little is known regarding their habits.

Key to Adults.

♂♂.

- | | |
|---|-------------------------------|
| 1 Abdomen with transverse basal pale bands on dorsum | 2 |
| Abdomen without pale bands on dorsum | 5. |
| 2 Mesonotal scales dark brown or brownish-black, segment 7 of antenna without a bunch of crumpled scales | 3 |
| Mesonotal scales light brown or reddish-brown, segment 7 of antenna with a bunch of crumpled scales | 4. |
| 3 Segments 7, 10, and 11 of antenna with some fairly short, thickened hairs, style somewhat expanded on apical $\frac{1}{2}$ (fig 87, a). | <i>senioris</i> , p. 365. |
| Segments 7, 10, and 11 of antenna without modified thickened hairs, style tapering from base to apex (fig 87, e) | <i>minutissimus</i> , p. 363. |

* [The species of this subgenus may be classed in three groups, according to their breeding habits, and there are also structural distinctions between these groups, which are—(1) Those which breed only in ground-pools and have the torus of ♂ antenna simple, without blunt prominence on inner side, palpi of ♂ longer than proboscis, with last two segments usually distinctly hairy, proboscis of ♂ usually with a row of stiff bristles beneath at base. This group includes *minutissimus*, *senioris*, *cinctellus*, *rubrothoracis*, and *fraudatrix* among Indian species. (2) Those which breed mainly in rock-pools, tree-holes, or bamboos, and have the torus of ♂ antenna with a blunt prominence on inner side; palpi of ♂ as long as proboscis or longer, but with few or no hairs at tip, ♂ proboscis without stiff bristles at base beneath. This includes the remainder of the known Indian species. (3) Those which breed mainly or exclusively in pitcher-plants, have a blunt prominence on inner side of torus of ♂ antenna, and palpi of ♂ shorter than proboscis and bare. This includes six Oriental species *caerulea*, *curtipalpis*, *eminentia*, *hewitti*, *jenseni*, and *navahi*, none of these have yet been found in the Indian area, but some probably occur. For distinction between the pitcher-plant forms, vide Edwards 1928, p. 276, and Edwards & Given 1928, p. 353]

- 4 Mesonotal scales reddish-brown, basal segment (torus) of antenna without a blunt prominence . . . *cinctellus*, p 366
 Mesonotal scales light brown, torus of antenna with a blunt prominence . . . *plantaginis*, p 372.
- 5 Torus of antenna without prominence . . . 6.
 Torus of antenna with prominence . . . 7
- 6 Palp with a small finger-like process at base, proboscis with a transverse row of about 6 fairly long stiff bristles on underside at base, antenna with a tuft of 12 or more long scales on segment 6 (fig 87, c) . . . *fraudatrix*, p 368
 Palp without a finger-like process, but with two small dense projecting tufts of hair near base, proboscis with a transverse row of 10-12 stiff bristles on underside at base, antenna with about 3 long pointed scales on segment 6 . . . *rubithoracis*, p 367.
- 7 Antenna with a tuft of long bright yellow hairs on segment 6 . . . *flavicornis*, p 375
 Antenna with a tuft of dark brown hairs or scales on segment 6 . . . 6
8. Palp with a row of stiff bristles on outer side near base . . . *mammulifer*, p 374.
 Palp without a row of stiff bristles . . . 9
- 9 Palp longer than proboscis by more than length of apical segment, last two segments distinctly hairy, antenna with a tuft of scales on segment 6, some of which are broad and very long (fig 87, l) . . . *unaformis*, p 373.
 Palp longer than proboscis by less than length of apical segment, last two segments with few hairs, segment 6 of antenna with a tuft of narrow hair-like scales all about same length (fig 87, d) . . . *minor*, p. 370

207. *Culex (Lophoceratomyia) minutissimus* (Theobald), 1907.

M.C. iv, p 235 (*Culicomyia*) (♀) TYPE-LOC. Peradeniya, Ceylon (Green) TYPE ♀ in Brit Mus

Culicomyia nigerrima Theobald, 1910, M.C. v, p 233 (♀) TYPE-LOC Trincomalee, Ceylon, x 1907 (Green) TYPE ♀ in Brit Mus.

Melanocomon justapallidiceps Theobald, 1910, M.C. v, p 456 (♀). TYPE-LOC Trincomalee, Ceylon, x 1907 (Green) TYPE: ♀ in Brit Mus

ADULT *.—Small species with banded abdomen and comparatively simple ♂ antennae†. Wing 2 mm. (♂) to 3 mm (♀)

Head. scales mainly dark brown, some paler brown scales on vertex and whitish flat scales at sides. Antennae, clypeus, palpi, and proboscis dark brown. Antenna of ♂ without a prominence on inner side of torus, segments 6 and 7

* Edwards 1913 b, p. 235 (syn.), 1922 c, p. 280 (syn.), Barraud 1924, p 41

† [The only other Oriental species with similar characters is *C. infanibilis* Edw., of Hong Kong, which differs slightly in ♂ hypopygium.]

without specialized hair-tufts, a small tuft of scales on 8, and a rather larger similar tuft on 9. Palpi of ♀ about $\frac{1}{2}$ length of proboscis, those of ♂ very slightly longer than proboscis, terminal segments with few hairs, a transverse row of long stiff bristles on underside of proboscis at base, a small dense tuft of hairs near base of each palp. *Thorax*: mesonotal scales brownish-black, fairly numerous dark bristles. *Pleurae* light brown, darker in some specimens, one lower mesepimeral bristle. *Wings* dark scaled. *Legs*: dark brown, femora faintly paler posteriorly. *Abdomen*: dorsum brownish-black, with narrow ochreous basal bands, sternite brown or light brown.

♂—*Hypopygium* (fig 88, i, k, p) essential points of structure shown in figures, *lp* pointed, without small teeth or tubercles or large median process.

LARVA *—*Antenna* comparatively large, shaft widest at about middle, spinelets along most of the length, dark at base and beyond hair-tuft, latter at about $\frac{2}{3}$ from base. Two strong preapical spines arising some little distance from tip of shaft, 2 weaker apical spines, one about $\frac{1}{3}$ length of other. *Frontal hair A* of 8–10 subplumose branches; *B* and *C* each of 2 long branches, but one or both may be single. Hair *d* anterior and internal to *B* and *C*, fine, fairly long, single or split towards tip. Preclypeal spines strong, dark, pointed. Mouth-brushes well developed, hairs simple. Some very long, single, unbranched hairs on thorax as well as the usual lateral branched hairs. *Abdomen*: segment I with 2 long lateral hairs, one usually 3-branched, the other 2-branched, a similar 2-branched or single hair on segment II. following segments with fairly long but weaker lateral hairs. Subsiphonal hair of 8–10 subplumose branches. Comb of about 40 long narrow teeth in a triangular patch. *Siphon* long and narrow, 7–9 times length of diameter at base. A fairly wide dark ring at base, and another wider dark ring at about middle. Pecten of about 12 teeth, with large lateral denticles from base to apex along one side, usually 4 pairs of small 2-branched hairs, fairly evenly spaced along tube from end of pecten to apex. *Anal segment* enclosed in chitinous ring, numerous small spines towards posterior border dorsally; *lh* of 3 fine short branches, *usc* usually of 4 branches, one very long, three short, *osc* single and long. Fan only moderately developed, about 12 hairs arising from fan-plate, each split into a number of branches. Papillae from $\frac{1}{4}$ – $\frac{1}{2}$ length of anal segment, both pairs pointed.

HABITAT—Rock-springs, pools in ravines and river-beds,

* Senior-White 1927, pl x

artificial collections of water in coconut-shells, etc., shallow wells, stagnant water in shaded culverts, etc

DISTRIBUTION —Fairly common from the PUNJAB to ORISSA, and through PENINSULAR INDIA to CEYLON. There are no records from Bengal, Assam, or Burma. It is probably frequently overlooked owing to its small size. It has been recorded from the islands of CELEBES, PANTAR, and ALOR (*Brug*)

208 *Culex (Lophoceratomyia) seniori*, sp n

TYPE-LOC Calcutta, Garden Reach, x 1931 (*Senior-White*)

TYPE. ♂ in Brit Mus

ADULT —Related to *minutissimus*, but differs in having either modified hairs or scales on segments 7–11 of antenna (fig 87, a) and in structure of hypopygium (fig 88, d). Wing about 2 mm

♂ —*Head* dorsum covered with dark narrow and upright scales, a patch of bluish-grey scales at each side. Palpi and proboscis dark brown. Palpi longer than proboscis by rather less than length of apical segment, last two segments with fairly numerous outstanding hairs. A transverse row of 10 long stiff bristles at base of proboscis beneath and two small, dense, projecting tufts of hairs near base of each palp. Antennæ strongly plumose, without prominence on torus, segment 6 simple, 7 and 8 with some short thick hairs, 8 also with a number of short, wide dark scales, 9 with about 6 wide dark scales with long pointed tips, 10 and 11 with 5–6 moderately long thickened hairs. *Thorax* mesonotal scales dark brown, scutellum and scales appearing lighter. Postnotum dark in middle, paler at sides. Pleuræ brown, without scales. *Legs* dark brown, femora paler beneath, as usual. *Abdomen*: tergites brownish-black, with well-marked basal ochreous bands. *Hypopygium* (fig 88, d): coxite with a group of about 14 straight hairs on dorsal border, not forming a row, and continued from near base to subapical lobe, latter with 3 rod-like processes, all about same length, two slightly curved, with hooked tips, one straighter, with filamentous tip, also a leaflet and several other leaf-like or sword-like processes. Style slightly expanded along apical $\frac{1}{2}$, with a rather long hair projecting from inner side. *Ip* with long, smooth, pointed apex, projecting at right angles to basal part as in *C. (L.) fraudatrix*. Paraproct (when mounted flat) with smooth rounded projection at apex, a number of strong spines and 2–3 small strong hairs just below.

♀ and LARVA.—Unknown.

DISTRIBUTION —Known only from type-locality.

209. *Culex (Lophoceratomyia) cinetellus* Edwards, 1922.

Ind Journ Med Res x, pp 281 & 473, nom nov. for *L. taenata* Leic
Lophoceratomyia taenata Leicester (nec Wiedemann), 1908, Oul.
 Malaya, p 127 (♂ & ♀) TYPE-LOC Kuala Lumpur and Klang,
 Malay Penin (Leicester) TYPE ♂ & ♀ in Brit Mus

ADULT * —Distinguished by banded abdomen and antennal structure of ♂ Wing about 3 mm •

♀—*Head*. narrow and numerous upright dark brown scales on vertex and nape, dark brown scales flat on either side, and greyish-white flat scales lower down at sides Palpi and proboscis dark brown Palpi about $\frac{1}{2}$ or $\frac{1}{3}$ length of proboscis *Thorax* mesonotum deep reddish-brown, pleuræ pale brown *Legs* deep brown, femora paler beneath, as usual *Abdomen* tergites brownish-black, with well-marked basal transverse ochreous bands

♂.—Coloration and markings as in ♀ Palpi longer than proboscis by a little less than length of apical segment, last two segments with numerous outstanding hairs Length of proboscis about 1.8 mm, and of segments of palp, respectively, 1.4, 0.34, and 0.35 mm A transverse row of stiff bristles on underside of proboscis at base, and two small, dense, projecting hair-tufts near base of each palp, as in some other species Antenna (fig 87, f) without a prominence on torus, 5-6 rather large scales on segment 6, matted and twisted tufts on segments 7-9, about 4 very long scales on segment 10, also some rather short thick hairs on this segment, as well as on segment 11 *Hypopygium* (fig 88, g) ooxite with a collection of straight hairs near dorsal border, not forming a row, as in some other species, subapical lobe with 3 slightly curved rods, two with hooked tips and one with expanded apex (this expansion can usually only be clearly seen in stained preparations), also 2 leaflets, one rather wide Style broader on apical $\frac{1}{2}$ than on basal $\frac{1}{2}$ resembling *fraudatrix* and *minutissimus*.

LARVA † —The following description of what was thought to be the larva of this species is taken from Edwards and Given (1928) —“*Head* broader than long, but not so large as in *C. fraudatrix*, pale in colour Clypeal spines slender, bristle-like, curved Clypeal tufts *A*, *B*, and *C* each with about eight slightly plumose branches, *C* very obliquely behind *B*, and neither of them reaching appreciably beyond the front of the head *Antennæ* entirely pale, moderately slender, and scarcely curved, not more than two-thirds as long as the head, shaft with numerous small spicules. Tuft placed at three-fifths, composed of about 12 slightly

* Edwards 1922 d, p 473; Barraud 1924 j, p 42.

† Edwards & Given 1928, p 353.

plumose branches, which do not reach beyond tip of shaft. Subapical hairs slender, pale, placed very close to the tip and not more than half as long as the shaft. *Mentum* with a rather small median tooth, and about ten teeth on each side of this, two or three on each side being larger and more detached. *Thorax* with the lateral tufts large; prothoracic hairs only just reaching front of head, the innermost hair double. *Abdomen* much as in *C. fraudatrix*, subsiphonal tuft less heavily plumose, comb-teeth pale. *Anal segment* with the posterior margin of the plate finely spinose (about 10-12 short spines). Inner dorsal hair simple. Brush composed of 6-8 small tufts, which have at most 4 or 5 branches, the transverse bars not well developed. Gills thick, rather blunt-ended, twice as long as the plate. *Siphon* brownish, slightly tapering on basal half, then almost parallel-sided, index about 6-6.5. Four pairs of small latero-ventral tufts which are mostly two-branched, the branches simple and about as long as the diameter of the siphon. Pecten of about ten teeth, which are rather short, with a row of five or six denticles on one side, several of these being as long as the main tooth.

"The adults issuing from these larvæ were unfortunately lost, and their identity is therefore uncertain †, from naked-eye appearance they seemed to be *Lophoceratomyia* with banded abdomen, and were therefore regarded as *C. cinctellus*."

DISTRIBUTION.—MALABAR COAST. Pudukudi*, x 1915 (*Khazan Chand*).

Known also from MALAY PENINSULA*, BORNEO, SUMATRA, and JAVA.

210. *Culex (Lophoceratomyia) rubithoracis* Leicester, 1908.

Cul Malaya, p 119 (♂ & ♀). TYPE-LOC. Kuala Lumpur, Malay Penn. (Leicester). TYPE ♂ & ♀ in Brit. Mus.

ADULT ‡.—Very similar to *cinctellus*, but abdomen unbanded. Wing about 2.5 mm.

♀.—*Head*. vertex and nape covered with pale narrow scales and numerous brown upright scales, some flat brown scales laterally, and below these, at sides, a patch of bluish-grey flat scales, the last extending dorsally as a narrow border.

† [The slender, pale, preclypeal spines and simple *asc* of this larva distinguish it from all others of the subgenus *Lophoceratomyia* hitherto known, though both these features occur in all the known larvæ of Oriental species of the subgenus *Culicromyia*. In the spinose margin of the chitinous ring of the anal segment this larva resembles *C. viridiventer* and *C. shebbeareae*, but differs from both these in the position of the antennal tuft.]

‡ Edwards 1922 c, p. 287.

to eyes Palpi and proboscis dark brown *Thorax* mesonotum reddish-brown or yellowish-brown, with scanty covering of narrow brown scales Pleuræ pale brown, without scales. *Legs* dark brown, undersides of femora paler *Abdomen* dorsum entirely dark brown, venter paler

No very definite characters can be given which distinguish this from ♀♀ of several allied species

♂—General coloration as in ♀ Palpi longer than proboscis by about $\frac{1}{2}$ length of terminal segment, last two segments with numerous outstanding hairs A transverse row of 10–12 stiff bristles on underside of proboscis at base, and two small, dense, projecting tufts of hairs near base of each palpi. Antenna without prominence on torus About 3 long, narrow, pointed scales on segment 6, matted and twisted tufts on segments 7–9, 3 long scales on segment 10 which are expanded towards tips, but have hair-like terminations Some rather short thick hairs on segments 10 and 11 *Hypopygium* processes on subapical lobe of coxite very similar to those of *cinctellus*, but only 3 hairs near dorsal border of coxite in a row, instead of a large collection. Style expanded apically, but less so than in *cinctellus*

LARVA—Unknown

HABITAT—Ponds (*Leicester*)

DISTRIBUTION—(Not recorded from India previously.) UPPER BURMA Bhamo *, x 1931 (*Feegrade*), 1 ♂. BENGAL Calcutta, Matiabruz *, Garden Reach, x 1931 (*Senior-White*), 1 ♂ SOUTH INDIA Mudgere *, Kadur dist, Mysore, 1931 (*Anantaswami Rao*) 1 ♂

Known also from MALAY PENINSULA, SIAM *, BORNEO, and SOUTH CHINA

211 *Culex (Lophoceratomyia) fraudatrix* Theobald, 1905.

Ann Mus Nat Hung in, p. 94 (♂ & ♀). TYPE-LOC. Friedrich-Wilhelmshafen, and Stephansort, Astrolabe Bay, both New Guinea, 1900 (*Buró*) TYPE ♂ & ♀ in Nat. Mus Hung., Budapest

Lophoceratomyia variata Leicester, 1908, Cul. Malaya, p. 121 (♂ & ♀) TYPE-LOC. Kuala Lumpur, Malay Penin (*Leicester*). TYPE ♂ & ♀ in Brit Mus

(For further synonymy and varieties see Edwards 1932, p. 197.)

ADULT †—Readily distinguished from other species of the subgenus by the large scale-tuft on segment 6 of ♂ antenna. Wing 2–3 mm

♀—*Head* vertex and nape covered with dark brown narrow and upright scales, a very narrow pale border to eyes, widening

† Theobald 1907, p. 474, Edwards 1913 b, p. 235 (syn.), 1922 c, p. 281 (*fraudator*), 1924, p. 398, Barraud 1924 j, p. 41 (*fraudator*).

out at sides. Antennæ, clypeus, palpi, and proboscis dark brown. Palpi about $\frac{1}{2}$ length of proboscis. *Thorax* mesonotal scales dark reddish-brown. Pleuræ brown, without scales, one lower mesepimeral bristle. *Wings* dark scaled. *Legs* dark brown, fore and mid-femora paler posteriorly, hind pair pale on both surfaces, but dark dorsally. *Abdomen* dorsum dark brown, small basal lateral ochreous markings usually present, sternites pale brown. (In var *annulatus* Taylor, from Australia, there are pale basal bands on dorsum).

♂—Antennæ plumose, torus without a projection on inner side. Some long scales on segment 6, forming a dark tuft projecting outwards and slightly upwards, below these, projecting downwards, are some long hairs, which appear white from certain angles, smaller tufts of crumpled scales on segments 7 and 8, the usual matted tuft on 9, and a tuft of hairs and scales on 10 (fig 87, c). Palpi longer than proboscis by $1\frac{1}{2}$ times length of apical segment. Length of proboscis about 1.8 mm, long segment of palp about 1.7 mm, each of last two segments about 0.3 mm, the terminal segments with numerous outstanding hairs, a small finger-like process at base of each palp, and transverse row of about 6 rather long stiff bristles on underside of proboscis at base. Thorax, legs, and abdomen as in ♀, except that on the last the lateral pale markings are sometimes absent. *Hypopygium* (fig 88, a, m) structure similar to that of *minutissimus* in some respects, but processes on subapical lobe of coxite of different form, and style widened towards tip.

LARVA*.—The following description of larvæ of *fraudatrix* from Singapore is taken from Edwards and Given — “*Head* large, considerably broader than long, mainly pale in colour, but with several dark spots on posterior part of clypeus, and also darkened round occiput. Clypeal spines moderately stout and straight. Tuft *A* large, plumose, with 7–12 branches, *B* and *C* both double, very long and reaching far beyond front of head, *C* placed obliquely behind *B*, *d* trifid, rather longer than usual. Antennæ fully as long as the head, rather strongly curved; a dark ring at the base and another immediately beyond the tuft, leaving the tip pale, shaft with strong spicules except on the slender terminal portion. Tuft very large, with 25–30 plumose branches, placed beyond two-thirds. Preapical bristles placed a short distance before the tip, both very long, about two-thirds as long as the shaft of the antenna, one of the terminal bristles also long, but shorter than the preapical pair. *Mentum* with a moderately large median tooth and 10–12 small teeth on each side, the outer one or two sometimes a little larger

* Edwards & Given 1928, p. 351.

and more detached, but not conspicuously so. *Thorax* with the lateral tufts large, plumose, and set in small but distinct chitinous plates. Prothoracic hairs very long, extending forwards far beyond front of head. *Abdomen* with smooth skin, the lateral tufts mostly triple. Eighth segment with three rather large branched tufts; the upper pair 6-branched and plumose, the middle (subsiphonal) pair with 6-8 branches and very thickly plumose, the ends of the branches sometimes split into several twigs. Comb a patch of about 30 very small dark scales, not obviously fringed. *Anal segment* with the chitinous ring hardly longer than broad, almost smooth on the posterior margin, inner dorsal hair with two small branches from base; lateral hair small, with 3-4 branches. Brush with about 10-12 large many-branched tufts. *Siphon* long, brown, usually with a dark brown ring in the middle, index about 8-9; in shape slightly tapering on about the basal half, then almost parallel-sided, the tip very slightly widened. Three or four pairs of ventral or ventro-lateral tufts, which have usually 2-3 branches, the branches simple and hardly longer than the diameter of the siphon. Pecten of 10-12 rather slender teeth, which are fringed along one side with fine denticles, those at the base rather stronger, the last tooth at about one-third the length of the siphon.

HABITAT—Pot-holes, etc., chiefly in mangrove tidal area, but also away from coast."

DISTRIBUTION.—ASSAM Dibrugarh*, viii. 1922 (*Barraud*); Golaghat*, Sibsagar dist., xii 1924 (*Barraud*). ANDAMAN Is Mount Harriet*, ix 1911 (*Christophers*).

Also known from MALAY PENINSULA, PHILIPPINES, MALAY ARCHIPELAGO, AUSTRALIA, NEW GUINEA, and SOLOMON IS.

212. *Culex (Lophoceratomyia) minor* Leicester, 1908.

Cul Malays, p 126 (♂ & ♀) TYPE-LOC.: Kuala Lumpur, Malay Penn (*Leicester*) TYPE ♂ & ♀ in Brit Mus.†

var n *bengalensis* (see below). TYPE-LOC.: Nongphoh, Assam (*Barraud*) TYPE ♂ in Brit Mus

ADULT ‡—Very similar to *uniformis* and *mammitifer*, but distinguished from these in the ♂ sex as indicated in key.

† There is some doubt regarding Leicester's type-specimens, as none could be found in his collection which agreed exactly with his description. Edwards (1917) re-defined the species, and I have followed this in describing the type-form. The variety referred to above may very possibly prove to be a distinct species. I have seen a few other specimens which differ slightly from the type-form and from var. *bengalensis*, and it is possible that these may be *C (L) bernardi*, described by Borel (1926, pp 111-114) from Cochin China, but further material is required before a decision can be arrived at.

‡ Edwards 1917, p 227; Barraud 1924, p. 44; Borel 1926, p. 109.

No reliable differences can be given by which ♀♀ may be separated

♂—Palpi only very slightly longer than proboscis⁺, last two segments with very few hairs. Antenna with a small number of fairly short hair-like scales on segment 6, all about same length (fig 87, d); no long dark expanded scales as in *uniformis* and *mammilifer*, but very similar matted and twisted tufts on segments 7-9, and some rather short thickened hairs on one side of segment 10. Hypopygium (fig 88, e, j) coxite with the usual 3 rod-like processes on subapical lobe, all about same length, two with hooked tips, one ending in a thin upturned filament, also a leaflet, 2 curved bristles, and a curved sword-like process. A row of usually 7 recurved hairs, extending further up dorsal border of coxite than in *uniformis*. Style not elbowed, not expanded in middle, but gradually tapering. *lp*, in side view, less pointed than in *uniformis*, and with a smaller median process.

var n *bengalensis*—Specialized hairs on segment 6 of antenna all very long, not scale-like, and with the basal part thickened. Palpi longer than proboscis by about length of last segment. These differences are correlated with a slight difference in the hypopygium, viz., a row of usually only 5 recurved hairs situated (as in *uniformis*), low down on dorsal border of coxite.

LAERVA †—Very similar to that of *fraudatrix*, but quite distinct from that of *uniformis*, the body-integument being smooth and not covered with minute hairs. Some larvae thought to be of this species, collected in India, agree closely with the following notes taken from Edwards and Given:— 'Structurally similar to *C. fraudatrix*, differing chiefly as follows—Antennae without dark ring at base, the narrow apical portion slightly and uniformly darkened, the longer apical bristle almost or quite as long as the preapical. Clypeal tuft 4 with about 8 branches. Siphon uniformly pale brownish and rather longer; index about 9-10, the basal third tapering, the outer two-thirds parallel-sided and very slender. Pecten with about 16 teeth, similar in form to those of *C. fraudatrix*, but the basal denticles somewhat stronger. Gills rather pointed, over twice as long as the anal plate.

"HABITAT.—The Singapore specimens were found in rock-pools in company with *Aedes (Finlaya) saxicola* Edw. Other

* [In Indian specimens there is a small dense patch of minute hairs on each side of the proboscis near base, and a number of longer curved hairs laterally and on under surface of proboscis before middle. These hairs do not appear to be present in Malayan specimens.]

† Borel 1926, p. 110, Senior-White 1927, p. 71; Edwards & Given, 1928, p. 352.

specimens in the British Museum were taken in old fallen bamboos at Ulu Gombak, F.M.S., by Dr H P Hacker these differ from the rock-pool specimens only in the darker siphon and stronger basal denticles to the pecten teeth, the adults are quite similar Barraud records the larvae from bamboos, rock-pools and tree-holes "

DISTRIBUTION —♂♂ of the type-form have been examined by the author from —LOWER BURMA Tenasserim, Victoria Point, and Mergui, all 1922 (*Sharma*). UPPER BURMA: Akyab, viii 1924 (*Feegrade*)

[Further ♂♂, apparently of the type-form, are in the British Museum from NORTH KANARA, Karwar (*Cogill*), and BOMBAY DECCAN, Tavargatti, viii 1921, from larvae in tree-holes (*Barraud*)]

♂♂ of var *bengalensis* —ASSAM Shillong, ix 1917 (*Fletcher*) Nongpoh, vii 1922 (*Barraud*) NORTH BENGAL: Sukna, and Marianbarrie Tea Estate, viii 1928 (*Sobha Ram*)

213 *Culex (Lophoceratomyia) plantaginis* Barraud, 1924

Incl. Journ. Med. Res. xii, p. 46 (♂) TYPE-LOC Kota, near Kalka (Kalka-Simla road), Himalayan foothills TYPE ♂ in Brit. Mus.

ADULT —Closely allied to *C. (L.) minor*, differing in the banded abdomen. Wing (♂) about 2.6 mm.

♂.—*Head*: vertex and nape covered with narrow pale scales and darker upright scales, a patch of creamy scales at each side, continued as a border to eyes, in front. Antennae, palpi, and proboscis dark brown. Palpi longer than proboscis by a little more than length of apical segment, length of proboscis about 1.87 mm., of long segment of palp 1.6, penultimate segment 0.31, apical segment 0.32, the last two segments with a few outstanding hairs, more numerous on penultimate than on apical segment. Antenna very similar to that of *minor* (type-form). *Thorax*: scutum and scutellum covered with brown scales, lighter in colour than in *C. (L.) uniformis* and allied species. Pleurae dark brown, coxae lighter, 2-3 *ppn* bristles, one lower mesepimeral bristle. *Wings*: thinly clothed with dark scales. *Legs*: dark brown, lighter when seen from behind; femora paler beneath, as usual. *Abdomen*: tergites brownish-black, with narrow basal ochreous bands*. Sternites brown. *Hypopygium*:

* In some specimens, apparently of this species, the pale bands on the abdomen are absent, or invisible owing to shrinkage. After examining a number of ♂♂ of this form, and of the type-forms of this species and *minor*, I am inclined to think that possibly they may all be forms of one species showing variation in markings, in the form of the scales on 6th segment of antenna, and in hypopygial details. Further work upon this difficult group of species is required.

(fig 88, f, o) very similar to that of *minor* (type-form) but usually 8-9 hairs in the row along dorsal border of coxite. Of the 4 rod-like processes on the subapical lobe of coxite, the one nearest to the style is usually longer, more slender, and more curved.

LARVA.—Unknown

HABITAT.—Pools (*Barraud*), cement water-channel in heavy sal forest (*Senior-White*).

DISTRIBUTION.—WESTERN HIMALAYAN foothills. Koti *, type-locality, as given above. ORISSA Gua *, Singhbhum dist, xi 1929 (*Senior-White*). BOMBAY DECCAN Tavar-gatti *, Belgaum dist, viii 1921 (*Barraud*).

214 *Culex (Lophoceratomyia) uniformis* Theobald, 1905.

Journ Bomb Nat Hist Soc xvi, p 245 (♂ & ♀). TYPE-LOC. Peradeniya, Ceylon (*Green*). TYPE. ♂ & ♀ in Brit Mus

ADULT † —Very similar to *C (L) minor* and *mammilifer* in the general blackish-brown colour, and no definite characters apparently exist by which the ♀♀ may be distinguished with certainty, there are, however, structural differences in the ♂♂, as given in key. Wing 2.5-3.2 mm

♀ and ♂.—*Head*: bluish-white flat scales at sides, continued as a narrow border to eyes in front, remainder covered with narrow and upright brownish-black scales. Antennae, palpi, and proboscis brownish-black. Antenna of ♂ (fig 87, i) with a blunt prominence on inner side of torus, segment 6 with 7-8 scales of varying lengths, the tip of the longest reaching beyond the matted tuft on segment 9 (when the antenna is mounted flat); the usual matted and twisted tufts on segments 7-9, no scales on 10. Palpi of ♀ about $\frac{1}{4}$ length of proboscis (comparatively slightly longer than in allied species). Palpi of ♂ longer than proboscis by rather more than length of apical segment, last two segments distinctly hairy and rather longer than corresponding segments in *minor* and *mammilifer*. Length of proboscis 1.7 mm, long segment of palp 1.5, penultimate segment 0.31, apical 0.34 mm. No spines or processes at base of proboscis or palpi. *Thorax*: mesonotum covered with narrow brownish-black scales. Pleurae dark green or dark brown, without any conspicuous markings or scales, 3-4 *ppn* bristles, one lower mesepimeral bristle usually present. *Legs*: deep brown, femora paler beneath, as usual. *Abdomen*: tergites entirely covered with brownish-black scales, venter somewhat paler. ♂ *hypopygium* (fig. 88, h, n): coxite with 3 rod-like processes on subapical lobe, two with hooked tips and one shorter

† Barraud 1924 j, p 43

and straight, also a leaflet and 2 bristles, dorsal border with usually only 4 hairs in a row, low down and some distance below subapical lobe. Style slightly elbowed and very little expanded in middle, tapering from this part to tip *lp* with a number of small teeth and rather large pointed median process

LARVA *.—Differs from other known larvae of the subgenus in having the integument of the body thickly covered with very small bristle-like hairs. *Antenna* of moderate length, with fine spinelets on shaft, tuft at about $\frac{1}{4}$ from base. 2 long preapical spines and one much smaller *Frontal hairs* *A*, *B*, and *C* with subplumose branches, the number of branches being respectively 6-11, 3-6, and 2-4. Hair *d* unusually long and stout, reaching nearly to front of clypeus. Predypeal spines strong and dark, in several specimens with small accessory spines projecting from sides. Mentum with large central tooth and 8-10 smaller teeth either side. Some long, single, unbranched, as well as usual branched, hairs on thorax. *Abdomen* with fairly long 3-5-branched hairs. Comb of 35-45 narrow fringed teeth in triangular patch. Three branched hairs and two single hairs posterior to comb on segment VIII, the former with 3-6 branches. *Siphon* widest at base and slightly tapering to tip, about 8-9 times length of diameter at base. Four subposterior pairs of hairs, with 4-5 rather long fine branches, evenly spaced between pecten and apex of tube. Pecten of 13-18 teeth, each with 3-4 sharp lateral denticles. *Anal segment* with numerous long thin spines on and near posterior margin, between fan-plate and subdorsal hairs, *lh* fine and unbranched. About 12 fan-hairs of moderate length, each divided into a number of branches, *osc* single, *isc* of 2-3 branches of different lengths. Both pairs of papillae pointed and about length of obtusised part of segment

HABITAT.—Tree-holes and rock-pools

DISTRIBUTION.—Common from BOMBAY down WEST COAST to CEYLON. There are, so far, no records from other parts of the Indian region, or from other countries

215 *Culex (Lophoceratomyia) mammillifer* Leicester, 1908.

Cul Malaya, p. 128 (♂ & ♀) TYPE-LOC. Raub, near Kuala Lumpur, Malay Penn. (Leicester). TYPE. ♂ & ♀ in Brit. Mus.

Lophoceratomyia bicornuta Theobald, 1910, Rec. Ind. Mus. iv, p. 26 (♂) TYPE-LOC. Dawna Hills (base), near Kawkaireik, Lower Burma, in 1908 (Annandale) TYPE ♂ in Ind. Mus.

ADULT †.—Very similar to *C (L.) uniformis* and *minor*, but mesonotal scaling in both sexes deep reddish-brown rather

* Senior-White 1927, pl. x, Edwards 1928 a, p. 276.

† Edwards 1913 b, p. 236 (syn.), Barraud 1924 j, p. 43.

than brownish-black. There are also differences in the ♂♂ in structure of hypopygia and in scale-tufts on antennæ.

♂.—Antenna with blunt prominence on inner side of torus, segment 6 with about 9 conspicuous scales of varying lengths the longest reaching to matted tuft on segment 9 (when antenna is mounted flat), also some hairs with expanded bases and long fine points. In dried specimens this collection of scales is rather more conspicuous than in *uniformis*. The usual twisted and matted tufts on segments 7–9, and a tuft of fairly short hairs on 10, some slightly expanded on basal $\frac{1}{2}$. Palpi longer than proboscis by a little less than length of apical segment, the last two segments short and subequal and with few outstanding hairs, a row of small, dark, stiff bristles on outer side of each palp near base (not present in *uniformis* or *minor*). Length of proboscis about 1.8 mm, of long segment of palp 1.5, penultimate 0.22, apical 0.20 mm. *Hypopygium* (fig 88, c, l) parts of diagnostic importance shown in figures. The presence of a long curved spine on subapical lobe of coxite should be noted, also form of style and presence of hairs, forming a crest, near tip.

LARVA †—This has not been isolated in India. The following note is taken from Edwards —“This species was reared by Dr. A. T. Stanton at Ginting Simpah, Federated Malay States. One larval skin is in the British Museum collection. This resembles *C. minor* very closely, the siphon is a little longer and more slender still, index about 10.5 anal gills shorter, not much longer than the plate.”

HABITAT—Pools in jungle (*Leicester*), rock-pools (*Senior-White*).

DISTRIBUTION—BURMA: type-locality, as given above NORTH BENGAL. Sukna*, Darjeeling dist., 500', viii. 1928 (*Sobha Ram*). ANDAMAN Is. Mount Harriet*, ix. 1911 (*Christophers*), and vii. 1926 (*Covell*). MALABAR COAST: Pudukudi*, x. 1915 (*Khazan Chand*). Ceylon: Suduganga, Matale dist., x. 1919 (*Senior-White*).

Known also from MALAY PENINSULA and BORNEO.

216 *Culex (Lophoceratomyia) flavicornis* Barraud, 1924.

Ind Journ Med Res xii, p. 45 (♂ & ♀). TYPE-LOC. Nilgiri Hills, x. 1915 (*Khazan Chand*). TYPE: ♂ & ♀ in Brit Mus.

ADULT.—Rather larger than the other species. Wing (♂) 3.5 mm, (♀) 4.3 mm. Distinguished by unusual hairiness of ♂ palpi.

♀ and ♂.—*Head*: both the narrow and flat scales on vertex and sides creamy-white, upright scales and bristles dark. Proboscis and palpi very dark brown. Palpi of ♀ $\frac{1}{2}$ – $\frac{1}{4}$ length

† Edwards 1928 a, p. 276

of proboscis, those of ♂ longer than proboscis by a little less than length of rather long apical segment, proboscis about 2.4 mm. long, long segment of palp 2 mm., penultimate segment 0.27 and apical segment 0.62 mm., long segment with numerous hairs of moderate length along underside for whole length, last two segments with a number of short hairs. Antenna of ♂ (fig 87, h) with blunt prominence on inner side of torus, segment 6 with tuft of long yellow hairs, tufts of crumpled scales and twisted hairs on segments 7, 8, and 9, very similar to those of several other species, a small tuft of straight hairs on segment 10. No bristles or processes at base of proboscis or palpi. *Thorax* mesonotal scales dark chestnut-brown, scutellar scales very narrow and lighter brown, mesonotal and scutellar bristles dark. *Pleuræ* dark brown, without scales, 4 *ppn* bristles, one lower mesepimeral bristle (apparently sometimes absent). *Legs* deep brown, femora paler beneath, as usual. *Abdomen* dorsum dark brown, venter ochreous. ♂ *hypopygium* (fig 88, b) subapical lobe of coxite with a leaflet, 3 long curved or sinuous spines, and a thick clubbed process. Style expanded apically and square-ended.

LARVA —Unknown

DISTRIBUTION —SOUTH INDIA Nilgiri Hills *, type-locality, as given above. Kodaikanal *, Palm Hills, 7,000', ix. 1929 (Fletcher)

Subgenus CULICIOMYIA Theobald, 1907

- M C iv, p 227 Genotype, *C fragilis* Lull (as *C. inornata* Theo.)
Trichorhynchus Theobald, 1905 (nec Balbian), Journ. Bomb. Nat. Hist. Soc. xvi, p 240 Genotype, *C fragilis* Lull (as *T. fuscus* Theo.).
Neomelanesconon Theobald, 1907 (nec Nowstead), M C iv, p 514. Genotype, *C. nebulosus* Theo (as *N. rima* Theo., ♂)
Pectinopalpus Theobald, 1910, M C v, p 416 Genotype, *C. nebulosus* Theo (as *P. fuscus* Theo.)
Trichorhynchomyia Brunetti, 1912, Rec. Ind. Mus. iv, p 477, nom. nov. for *Trichorhynchus*


ADULT —Mosquitoes of moderate size. Head with flat scales at sides, and with a row of small flat scales along eye-margins, usually forming a continuous border. Other head-scales narrow and upright. Mesonotal scales narrow and dense, giving the thorax a characteristic smooth appearance in most species. Scutellar scales narrow. Tarsi and wing-scales dark. *Pleuræ* without scales, but with dark markings in those species in which abdomen is marked with pale bands. Usually one lower mesepimeral bristle. Palpi of ♂ longer than proboscis, the long segment with a row of peculiar translucent scales projecting downwards. Antenna of ♂ without scale-tufts. Coxite of ♂ hypopygium without scales;

usually with an apico-ventral hairy lobe, also a subapical lobe carrying several spines and usually a leaflet. Style usually with spiny crest. Phallosome divided into two pointed plates, with a row of small teeth. Paraprocts with spines or hairs at crown, and lateral arm variously developed.

Bucco-pharyngeal armature of ♀ with usually two rows of small sharp teeth.

LARVA — Very similar to those of subgenus *Culex*.

Fig 89.



♂ hypopygial structures (tip of coxite, proctiga, and phallosome) of *Culex*, subgenus *Culicomyia*: a, *paludothorax*; b, *shebbeari*; c, *viridiventer*; d, *bairdi* (phallosome not shown); e, *nigroaenatus*. av, apico-ventral lobe of coxite. Other lettering as on p. 4.

DISTRIBUTION and ECONOMICS — This subgenus is represented in most parts of India, several species being common. Of nine Oriental species, seven are found in India. The remaining five known species are confined to Africa. Breeding-habits vary with the species; the larvae of several live in tree-holes, of others in stream-pools, domestic collections of water, rice-fields, etc. The adult ♀♀ do not appear to attack man to any extent.

Key to Adults

- 1 Abdomen with transverse pale bands on dorsum, integument of pleuræ with dark markings 2
Abdomen entirely dark on dorsum, pleuræ uniform in colour 6
- 2 Pleuræ pale, with a distinct black spot on upper part of mesepimeron *nigropunctatus*, p. 383.
Pleuræ with a brown stripe across upper part, and sometimes another dark area on sternopleura, no distinct isolated black spot on mesepimeron 3
- 3 Mesonotal scales deep brown, palpi of ♂ with only a few hairs on last two segments 4
Mesonotal scales brown or fawn in colour, palpi of ♂ with numerous hairs on last two segments 5
- 4 Abdominal pale bands usually subtriangular, fork-cells in wing of ♀ long, 3 times length of stems, basal arm of ♂ paraproct very large and bag like (fig. 289, c) *mutiventris*, p. 378
Abdominal pale bands straight, fork-cells in wing of ♀ shorter, not more than $2\frac{1}{2}$ times length of stems, basal arm of ♂ paraproct quite small (fig. 89, b) *shebbareti*, p. 380
- 5 Mesonotal scales dark fawn in colour, a dark area on sternopleura, as well as dark stripe across upper part of pleuræ, style of ♂ with spiny crest (fig. 89, a) *pallidothorax*, p. 381.
Mesonotal scales brown, no distinct dark area on sternopleura, but a dark stripe across upper part of pleuræ, style of ♂ without obvious spiny crest (fig. 89, d) *barlyi*, p. 382
- 6 Plume series of wing-scales broader (about 5 times length of greatest width), penultimate segment of ♂ palpi only about twice as long as labella *fragilis*, p. 385
Plume series of wing-scales narrower (about 10 times length of greatest width), penultimate segment of ♂ palpi fully 3 times as long as labella *bahræ*, p. 386.

217 *Culex (Culicomyia) viridiventer* Giles, 1901.

Journ. Bomb. Nat. Hist. Soc. xxi, p. 609 (May) (♀). TYPE-LOC.: Nani Tal, U.P., 4,000' (*Giles*). TYPE: ♀ in Brit. Mus.

Culex angulatus Theobald, 1901 (November), M.C. ii, p. 324 (♀). TYPE-LOC. as above. TYPE: ♀ in Brit. Mus.

Culex longifurcatus Theobald, 1910 (nec Becker), Rec. Ind. Mus. iv, p. 19 (♂ & ♀). TYPE-LOC.: Dahawangahary Hills, near Bengal Frontier, Nepal, in 1908 (*Brachy*). TYPE: ♂ & ♀ in Ind. Mus.

Culex pseudolongifurcatus Theobald, 1910, M.C. v, p. 366, nom. nov. for *C. longifurcatus* (name prooon).

ADULT * —Size rather large. Wing 4–5 mm.

* Edwards 1913 b, p. 235 (syn.), Barraud 1924 a, p. 18, Borel 1926, p. 106.

♀—*Head*. a patch of flat white scales at each side usually visible from above, the flat scales continued dorsally as a narrow border to eyes, narrow pale ochreous and dark upright scales on vertex, upright scales more numerous on nape. Palpi and proboscis black, palpi about $\frac{1}{2}$ length of proboscis. *Thorax*: mesonotum covered with deep brown narrow scales, giving it a smooth, dull, velvety appearance, usually some pale scales either side of antescutellar space, bristles dark, most numerous over wing-roots. Scutellar scales narrow and pale ochreous. Pleurae usually pale green and without scales, a dark stripe from *ppn* to upper part of mesepimeron, and another dark area on sternopleura, 4–5 *ppn* bristles, one lower mesepimeral. *Wings* outstanding (plume) scales long and narrow, fork-cells long, each about 3 times length of stem. *Legs* black when seen from front or from above, sometimes with deep bluish lustre, pale brown when seen from behind, hind femur white except dorsally, and with pale knee-spot. a narrow pale ring also at tibio-tarsal joint. *Abdomen*: tergites black, with well-defined basal pale bands which are usually produced triangularly in middle, but this is not very marked in some specimens. Sternites pale green or pale ochreous.

♂—Palpi longer than proboscis by a little less than length of apical segment, long segment with a row of long, pointed, translucent scales projecting downwards, last two segments with few outstanding dark hairs. Proboscis about 2–3 mm. long, segments of palp of following lengths, respectively. 1.65, 0.34, and 0.48 mm. Fork-cells of wing about twice length of stems. Markings as in ♀ *Hypopygium* (fig 89, c). form of paraproct distinctive, the lateral arm being very much enlarged.

LARVA*.—*Antenna* long (about 0.8–0.9 mm.) and about $\frac{1}{2}$ length of head, shaft with fine spicules, most numerous on basal $\frac{1}{2}$, fan-shaped hair-tuft slightly nearer base than apex, preapical spines some little distance from tip of shaft, apical $\frac{1}{2}$ of shaft sometimes darkened, but not always. *Frontal hair* A with 5–7 branches, B and C with 2–3 long branches (in one specimen C is 4-branched), d fine, single, and fairly long. Preelypeal spines long and slender. Mouth-brushes moderately large, hairs simple. Mentum subtriangular, about 15 teeth either side of central one. *Thorax*: prothorax with some long single and 2-branched hairs dorsally. Lateral and subventral branched hairs on meso- and metathorax moderately developed. *Comb* of about 40 fairly long fringed teeth in a triangular patch.

* Borel 1926, p. 107.

Siphon about 2.3 mm long and about 6 times length of diameter at base, slightly tapering from near base to apex (not swollen at about middle as in *pallidothorax*). Pecten of 14-18 teeth, each with 5-7 fairly large lateral denticles along one side from base to apex. Four pairs of subposterior hairs along siphon between pecten and apex, each of 2-3 fairly long branches. *Anal segment* with some strong spines on posterior margin towards dorsum. Both *osc* and *isc* single and long, *lh* fine, single, moderately long. Fan small, 8-10 hairs, each divided into several fairly short branches, arising from fan-plate. Papillae very long and bluntly pointed, dorsal pair sometimes nearly 4 times length of anal segment.

HABITAT—Tree-holes, garden water-tanks, and butts.

DISTRIBUTION—Common in WESTERN HIMALAYAS from western boundary of Nepal to Kashmir, 5,000' to over 8,000'. I am in doubt as to records from other parts, and have not seen any typical specimens from Assam, South India or Ceylon. Examples from Ceylon previously identified as this species proved to be *C. (C.) shebbearei*.

218 *Culex (Culicomyia) shebbearei* Barraud, 1924.

Ind Journ Med Res xii, p 10 (♂ & ♀). TYPE-LOC. Kurseong, Darjeeling dist., E Himalayas, 18 1022 (Barraud). TYPE: ♂ & ♀ in Brit Mus.

ADULT.—Resembles *C. (C.) viridiventer* and *pallidothorax* fairly closely, but is rather smaller. Wing 3.3-3.6 mm. From the former it differs chiefly in structure of ♂ hypopygium, in venation of ♀ wings, and in shape of pale abdominal bands, which are not triangularly produced in middle. Fork-cells 2-2½ times length of stems. Basal arm of ♂ paraproct quite small compared with that of *viridiventer*. From *pallidothorax* this species differs in colour of head-scaling, the narrow scales on vertex being white or very light in colour instead of brown. Palpi of ♂ with only few outstanding hairs on last two segments. Mesonotal scales deep brown, as in *viridiventer*. Subapical lobe of coxite without a leaflet which is present in *pallidothorax*. Style with only a short crest of spines towards apex (fig. 89, b).

LARVA—Very similar to *viridiventer*. One skin definitely identified as this species by examination of hypopygium of ♂ which resulted apparently differs only in length of siphon, which is 1.4 mm. long. The siphonal hairs and majority of clypeal hairs are missing from this specimen.

HABITAT—Tree-holes (India), large pitcher-plant, *Nepenthes rajah* (Borneo).

DISTRIBUTION—EASTERN HIMALAYAS Kurseong *, type-

locality, as given above WESTERN HIMALAYAS. Simla *, ix 1915 (*Christophers*), and ix 1930 (*Barraud*).

One ♂ from Suduganga *, Matale dist., CEYLON, ix 1919 (*Senior-White*), appears to be this species

Also recorded from NORTH BORNEO (*Pendlebury*) †

219. *Culex (Culicomyia) pallidothorax* Theobald, 1905

Journ. Eoon. Biol. 1, p. 32 (*Culex*) (♂ & ♀) TYPE-LOC India (James) TYPE ♂ & ♀ in Brit. Mus.

Culex albopictura Theobald, 1907, M.C. iv, p. 456 (♀) TYPE-LOC India (*Christophers*) TYPE ♀ in Brit. Mus.

Culicomyia annulobdominalis Theobald, 1910, M.C. v, p. 236 (♂ & ♀). TYPE-LOC Peradeniya and Hakgala, Ceylon, 1907 (*Green*) TYPES non-existent

ADULT †.—Size moderate. Wing 3.5–4.5 mm.

♀.—Resembles *C. (C.) viridiventer* and *shebbearei* in general appearance, but differs as follows:—Head-scaling darker, the narrow scales brown instead of creamy. Scales covering mesonotum dark fawn-brown instead of deep dark brown. Anterior fork-cell of wing about $2\frac{1}{2}$ times length of stem. Hind femur without a distinct pale knee-spot dorsally. Pale basal bands on abdomen straight or slightly rounded on posterior margins.

♂.—Palpi longer than proboscis by about length of fairly long terminal segment, last two segments with numerous outstanding hairs. *Hypopygium* (fig. 89, a). differs from that of *viridiventer* in the much smaller basal arm of paraproct, and from *shebbearei* in the longer spiny crest on style and presence of a leaflet on subapical lobe of coxite.

LARVA §.—Resembles that of *viridiventer* in most details, but differs as follows:—*Siphon* somewhat swollen, widest at about $\frac{1}{3}$ of distance from base, and tapering from this part to apex, which is rather narrow; length about 5 times that of diameter at base, and about 3.6 times that of greatest width. Pecten of only about 5–8 teeth, usually commencing some little distance from base of siphon. Four pairs of subposterior hair-tufts, each of 4–5 fine short branches, the two most distal pairs often fairly close together and some distance from the two more proximal pairs, but position of individual tufts varies, as in other species.

HABITAT.—Tree-holes, bamboos, shallow wells, stream and rock-pools, swampy ground-pools, sometimes in foul water.

DISTRIBUTION.—Common in the Indian region, both in the hills and plains, from the PUNJAB to ASSAM and BURMA, and southwards to CEYLON.

† Edwards 1931, p. 28.

‡ Theobald 1907, p. 446, Edwards 1913 b, p. 235 (syn.); Barraud 1924 a, p. 20, Borel 1926, p. 104.

§ Borel 1926, p. 105, Senior-White 1927, p. 71.

Known also from MALAY PENINSULA, SIAM, COCHIN CHINA, SOUTH CHINA, and as far east as HALMAHEIRA ISLAND, east of Celebes.

220 *Culex (Culicomyia) bailyi*, sp. n.

TYPE-LOC. Virajpet, Coorg, South India, vi 1927 (*Baily*) TYPE.
♂ & ♀ in Brit. Mus.

ADULT.—Resembles *C. (C.) pallidothorax*, differing chiefly in ♂ hypopygium. Wing 3.3–3.6 mm.

♀—*Head*: vertex and nape covered with narrow yellowish and dark brownish upright scales, broad dark brown scales on either side towards front, and a patch of greyish-white flat scales each side laterally, those pale scales continued dorsally as a narrow border to eyes to middle point. Tori pale brown, clypeus, palpi, and proboscis dark brown. Palpi about $\frac{1}{2}$ length of proboscis. *Thorax*: mesonotum curved, with brown narrow scales, somewhat darker than in *pallidothorax*, but less dark than in *viridiventer* and *sheb-bearing*, bristles dark. Scutellar scales similar to those on mesonotum. Postnotum and pleurae pale greyish or greenish, except upper part of latter, where colour is darker and brownish from *ppn* to beneath wing-root. *Wings*: fork-cells about twice length of stems. *Legs*: dark brown, fore and mid-femora extensively pale posteriorly, hind femur pale on both sides, except dorsally. *Abdomen*: tergites brownish-black, with well-marked or pale ochreous bands on III–VII. VIII also pale scaled, posterior margins of pale bands almost straight.

♂.—Coloration and markings as in ♀. Palpi longer than proboscis by about length of last segment, last two segments with numerous outstanding dark hairs; long segment with about 8 long, pointed scales on apical $\frac{1}{2}$, directed downwards. *Hypopygium* (fig. 89, d) style wide on apical $\frac{1}{2}$, then tapering; terminal part narrowed for short distance; tip truncated, with terminal appendage as usual; no obvious spiny crest (differing from other species in this respect). Apico-ventral lobe of coxite not prominent, carrying a leaflet, a shorter straight sword-like process, and a strong bristle. Subapical lobe with two rod-like processes as usual; one very strong, tapering, with truncated tip, the other pointed; also a shorter, strong, bluntly-pointed spine and 4–5 fairly strong bristles. *tp* with one large tooth towards base and a row of much smaller teeth above, latter have a similar arrangement to those of *pallidothorax*, but appear to be smaller.

LARVA.—Unknown.

DISTRIBUTION.—Known only from type-locality.

221. *Culex (Culicomyia) nigropunctatus* Edwards, 1926*.

Bull. Ent. Res. **xvii**, p. 121 (nomen nov. for *annulata* Theob., preoccupied)

Culicomyia annulata Theobald (nec *Culex annulatus* Schrank), 1907, M.C. **iv**, p. 230 (♂ & ♀). TYPE-LOC. Kuching, Sarawak, in house (Barker). TYPE. ♂ & ♀ in Brit. Mus.

ADULT—A small brown species, which may be distinguished by presence of a very distinct velvety-black spot on mesepimeron. Wing about 3–3.6 mm.

♀—**Head**—a patch of flat white scales at sides, continued dorsally as a rather broad and irregular border to eyes nearly to middle point, vertex covered with narrow pale ochraceous scales and darker upright scales. Proboscis and palpi brown, latter about $\frac{1}{4}$ length of proboscis. **Thorax** mesonotal scales dark fawn-brown, scutellar scales lighter. **Pleurae** pale brown, with a very distinct velvety-black (not brown) spot on upper part of mesepimeron and two less distinct dark areas, one posterior to *ppn* and one in middle of sternopleura, one lower mesepimeral bristle. **Legs** brown anteriorly, hind femur pale except dorsally, all legs, when viewed from behind, pale brown or yellow. **Abdomen**: tergites brown, with narrow pale ochraceous basal bands, and with usually a thin line of pale scales along apical margins of tergites. Sternites pale ochraceous.

♂.—Palpi longer than proboscis by a little more than length of last segment, last two segments with a number of outstanding hairs, long segment with a row of very long, pointed, translucent scales on apical $\frac{1}{2}$, projecting downwards, a collection of moderately long hairs projecting downwards from middle of proboscis. Pale basal abdominal bands usually wider than in ♀, otherwise markings are very similar. *Hypopygium* (fig. 89, e) apico-ventral lobe of coxite with 2 large processes and 3 smaller bristles, subapical lobe with 2 rod-like spines, 2 smaller processes, and a leaflet. Style with spiny crest on apical $\frac{1}{2}$, continued nearly to tip. Basal arm of paraproct wide and fairly short.

* This species has been referred to previously as *Culex (Culicomyia) pullus*, originally described by Theobald as *Culex pullus* from New Guinea (Ann. Mus. Nat. Hung. **xii**, p. 87, 1905). This was later considered to be synonymous with *Culicomyia annulata* Theobald, but in 1926 Edwards considered that the Oriental form was distinct from that found in New Guinea, and introduced the name *nigropunctatus* for the former (the name *annulata* being preoccupied). The adults of *nigropunctatus* differ from *C. (C.) pullus*, as now restricted, in having a black (not brown) spot on the mesepimeron. I have seen adults of both sexes from the Philippines, and these appear to be identical with Indian specimens in markings and in structure of ♂ hypopygium.

[LARVA † — Differs from all other known larvæ of Indian Culicine mosquitoes in the false-jointed siphon-tube. *Antenna* of usual form, slightly and uniformly darkened. two hairs about $\frac{1}{2}$ length of antenna slightly removed from tip (about $\frac{3}{4}$ or $\frac{4}{5}$ of distance from tuft to tip). *Frontal hairs* *A*, *B*, and *C* with 6-8, 3-4, and 3 branches respectively, *B* and *C* reaching much beyond front edge of clypeus, *C* internal and only slightly posterior to *B*, *d* small and single or bifid. Pre-clypeal spines long, slender, curved, and pale. *Mentum* with 12-15 teeth on each side of median tooth. *Thorax* without special structures, meso- and metapleural tubercles without distinct spines, one of the long hairs in each of these groups with 6-8 branches. *Abdomen* with dorso-lateral hair on I and lateral hair on II 3-branched, lateral hairs on III-VI long and single. Comb of about 40 long, narrow, fringed teeth in a triangular patch. Dorso-lateral and subsiphonal tufts of VIII each of 6-8 plumose branches, subsiphonal tuft considerably the larger. *Siphon* 2-2.4 mm. long and about 9-10 times as long as its diameter at base, gradually tapering from base to tip. At about $\frac{2}{3}$ or $\frac{3}{4}$ length of tube from base is an unchitinised ring, wider anteriorly, narrow posteriorly, forming a false joint from which distal $\frac{1}{3}$ of siphon is readily bent forwards ‡. Pecten of 9-11 teeth, last few widely spaced, all similar in form, wide at base, with denticles extending along one side nearly to tip. Only three pairs of fine single or bifid hairs, all lateral in position, and scarcely longer than diameter of tube, one beyond end of pecten, one immediately before the false joint, and the third at middle of distal part of tube. *Anal segment* ringed as usual, posterior margin of ring with fine spicules on upper $\frac{1}{2}$. Both *osc* and *usc* long and single, *lh* also single, slightly longer than ring. Papillæ almost 3 times as long as ring, pointed.]

HABITAT — Rice-fields (*Barraud*), pools (*MacDougall*)

DISTRIBUTION — EAST BENGAL Chittagong* and Rangamati*, Chittagong Hill Tracts, viii 1922 (*Barraud*). ASSAM

† [A brief description of a larva supposed to be this species was given by Edwards (1926, p. 121). At the suggestion of the author the material in the British Museum has been re-examined, and it appears certain that the larvæ described as *nigropunctatus* were wrongly labelled; they are probably skins of *C. vishnu*. Other larval skins from Singapore were evidently correctly labelled, as they agree in all respects with a larva determined by Barraud as this species from Chittagong Hill Tracts, Eastern Bengal. The above description has been drawn up from these specimens.]

‡ [The extent of the area in which chitinisation is lacking is almost exactly the same in all skins examined (two from Singapore, two from Eastern Bengal, eight from Ceylon).]

Nongpoh *, Khasi Hills dist, and Dinapuri *, Naga Hills dist, viii 1922 (*Barraud*), Golaghat *, Sibsagar dist, x 1924 (*Barraud*) NORTH MADRAS Coast Vizagapatam *, ii 1928 (*Puri*) MYSORE. Mudgere *, Kadur dist (*Anantaswami Rao*) CEYLON. Colombo * (*James*), Galle Face Battery, Colombo, x 1909 (*MacDougall*)

Also known from MALAY PENINSULA, BORNEO, PHILIPPINES (see footnote on previous page), and CELEBES (Mamoedjoe, *Dr Eiber*)

222. *Culex (Culicomyia) fragilis* Ludlow, 1903.

- Journ N.Y. Ent Soc xi, p 142 (♂ & ♀) TYPE-LOC Oms. Samar, Philippine Is TYPE ♂ & ♀ in U.S. Nat Mus
Trichorhynchus fuscus Theobald, 1905, Journ. Bomb Nat Hist Soc xvi, p 242 (♀) TYPE-LOC Peradeniya, Ceylon, xii 1901 (*Green*) TYPE ♀ in Brit Mus
Culicomyia inornata Theobald, 1907, M.C. iv, p. 227 (♂ & ♀) TYPE-LOC Kuching, Sarawak, Borneo (*Barker*). TYPE ♂ & ♀ in Brit Mus.
Culicomyia ceylonica Theobald, 1907, M.C. iv, p 236 (♂ & ♀) TYPE-LOC Peradeniya and Maskahya, Ceylon (*Green*) TYPE ♂ & ♀ in Brit Mus
 ? *Culex graminis* Leicester, 1908, Cul. Malaya, p 158 (♂ & ♀) TYPE-LOC Kuala Lumpur, Malay Penin (*Leicester*) TYPES non-existent

ADULT †.—Differs from other Indian species, with the exception of *C. (C.) bahri*, in absence of pale bands on abdomen and of dark markings on pleuræ Outstanding (plume) scales on wings broader than in any other species Wing about 3–3.6 mm

♀.—Head and appendages similar to *vuulwenter* Thorax. mesonotal scales deep brown, pleuræ usually of a uniform greenish colour, without dark markings Wings. outstanding plume-scales about 5 times length of greatest width, instead of a more usual measurement of 10 times Legs dark brown, femora paler beneath, as usual, but without pale knee-spots Abdomen dorsum brownish-black, venter paler

♂.—Palpi longer than proboscis by about length of last segment; penultimate segment about twice length of labolla. last two segments with a moderate number of outstanding hairs Other details as in ♀ Hypopygium: very similar to that of *nigropunctatus*, but basal arm of proctiger narrower, spiny crest upon style rather longer, and apico-ventral lobe less prominent

LARVA.—Unknown

HABITAT.—Tree-holes (*Barraud*)

† Edwards 1914 b, p. 79 (syn.), 1920 a, p. 4, Barraud 1924 i, p. 21, Dyar & Shannon 1925, p. 85 (note on syn.).

DISTRIBUTION — NORTH KANARA Kadra *, ix 1921 (*Barraud*) ASSAM: Nongpoh *, Khasi Hills dist., vii. 1922 (*Barraud*) CEYLON type-localities, as given above, Colombo, 1913 (*James*)

Recorded also from MALAY PENINSULA, PHILIPPINES, and BORNEO

223. *Culex (Culiciomyia) bahri* Edwards, 1914

Bull. Ent. Res. v, p 79 (♂ & ♀) TYPE-LOC Badulla, Ceylon (*Bahr*). TYPE ♂ & ♀ in Brit Mus

ADULT † — Very similar to *C (C) fragilis*, but differs as indicated in key and in having fewer flat white scales along eye-margins, a more rounded coxite to ♂ hypopygium, and a shorter basal arm to proctiger. The following is a copy of the original description —

"Head clothed with black upright forked scales and yellowish narrow curved scales in the middle, flat scales at the sides, which are white below and purplish-black above, the white ones, however, extending upwards along the eye-margins towards the middle line, but they do not, as in other members of the genus, quite reach the middle. Basal joint of antennae blackish, without scales. Palpi and proboscis black-scaled. In the male the palpi are longer than the proboscis by a little more than the length of the last joint, the penultimate joint is only a little shorter than the terminal, and about three times as long as the labella, the projecting scales on the basal joints are much fewer, shorter and more difficult to see than in any other member of the genus. Thorax with the usual dull (matt) greyish-brown integument, clothed with small, narrow, dark brown scales, lighter on the scutellum. Plourae pale, unscaled, without any dark markings. Abdomen clothed with blackish-brown scales above, whitish ones below; no trace of white lateral spots at the bases of the segments, but a few paler scales at the apical corners. Male genitalia with spiny claspers and with the second plate of the harpagones long, pointed, and serrated below, as in all other species of the genus. Legs entirely dark-scaled except for the undersides of the femora and the lateral aspect of the hind pair. Wings with dark brown scales; the lateral veins towards the apex of the wing are linear. Fork-cells rather long, the upper one about twice as long as its stem, and with its base a little nearer the base of the wing than that of the lower.

"Ceylon Badulla (*Dr. P. H. Bahr*), a series bred from larvæ, 3 ♂ (including type), 4 ♀, in the British Museum; other

† *Barraud* 1924 i, p 21.

specimens in the London School of Tropical Medicine, 1 ♀, Hakgala, and 1 ♀, Peradeniya (*E E Green*)

"The specimens were all at first thought to be *C. fragilis* Ludlow, but there are abundant distinctions between the two, *C. bahr* having narrower lateral vein-scales, fewer flat white scales on the orbital margins, fewer outstanding scales, and a longer penultimate joint to the male palpi, and in the genitalia more rounded side-pieces and a shorter basal projection on the harpagones "

LARVA —Unknown *

DISTRIBUTION.—CEYLON Badulla, type-locality, as given above, Hakgala, and Peradeniya (*Green*).

Specimens apparently of this species are in the British Museum from JAVA (Djajasana, *Brug*).

Subgenus CULEX Linnæus, 1758

Syst. Nat ed x, p 602 Genotype, *C. pipiens* Linn

Leucomyia Theobald, 1907, MC iv, p 372 Genotype, *C. gelidus* Theo

Oculeomyia Theobald, 1907, MC iv, p 515 Genotype, *C. biternio-rhynchus* Giles (as *O. sarawaki* Theo)

Theobaldomyia Brunetti, 1912, Rec Ind Mus iv, p 462 (nom nov. for *Leucomyia* Theo)

(For further synonymy *vide* Edwards 1932, p 200)

ADULT —Moderate-sized to rather small mosquitoes Head with upright and narrow scales on vertex, flat scales at sides, but no border of flat scales round eye-margins in front. Scutellar scales narrow. Ornamentation rather varied. Palpi of ♂ always longer than proboscis, with last two segments upcurved and hairy, long segment without the erect translucent scales which characterize *Culicomyia*. Antenna of ♂ without scale-tufts. Coxite of ♂ hypopygium without scales and without an apico-ventral lobe, but with a subapical lobe which is usually more or less divided into two portions and bears some modified spines and usually a leaflet. Style without spiny crest. Phallosome divided into lateral portions which usually have a complicated structure and are provided with various teeth or lobes. Paraprocts with rather dense tufts of spines or hairs at crown, and usually with basal lateral arm.

Bucco-pharyngeal armature of ♀ with one row of teeth, which may be sharp or blunt.

LARVA.—With the characters of the genus; mouth-parts normal, not modified for predacity

* The figure given by Senior-White (1927, pl xi) probably represents an *Aedemorphus* larva in which he overlooked the single pair of very small hair-tufts.

DISTRIBUTION.—Species of this subgenus occur throughout the world; they may, however, be classed in two distinct groups, according to the presence or absence of the lower mesepimeral bristle, the species in which this bristle is lacking all have pale rings on the proboscis and tarsi, and this group is most extensively developed in the Oriental region, although many species occur in Australia and Africa. In the second group, where the lower mesepimeral bristle is present, the proboscis and tarsi are usually dark (as is the case in all Indian species), comparatively few species occur in India, several of those which are found having a wide distribution in other regions.

BIONOMICS.—All the Indian species of this subgenus breed in ground-pools, usually those of a more or less permanent nature. Several of them are among the commonest and most troublesome of Indian mosquitoes.

Key to Adults.

- | | |
|---|----------------------------------|
| 1 Proboscis and tarsi with pale rings (tarsal rings sometimes faint), no lower mesepimeral bristle . . . | 2 |
| Proboscis without pale ring (sometimes indistinctly ringed in ♂), tarsi entirely dark, lower mesepimeral bristle present. | 13 |
| 2 Wings with costa dark unless at tip | 3 |
| Wings with three pale spots on costa (including one at tip) | 12. |
| 3 A yellowish area at tip of wing, body and legs largely yellow | <i>epidesmus</i> , p. 389 |
| No yellowish area at wing-tip, body and legs not usually extensively yellow | 4 |
| 4 Wings speckled with pale scales, which are usually numerous | <i>bitaniorhynchus</i> , p. 391. |
| Wing-scales all dark | 5 |
| 5 Scales on anterior $\frac{2}{3}$ of mesonotum mainly or all white or pale ochreous | 6 |
| Mesonotal scales mainly dark or with indefinite pale mottling | 8 |
| 6 Abdominal tergites apically banded, mesonotal scales pale ochreous | <i>sinensis</i> , p. 394 |
| Abdominal tergites basally banded, mesonotal scales white | 7 |
| 7 Wings with broad scales on veins 1, 3, and 5, mid- and hind tibiae with pale lines . . . | <i>whitmorei</i> , p. 406 |
| Wings without unusually broad scales, tibiae not lined . . . | <i>gelidus</i> , p. 407 |
| 8. Abdominal tergites with distinct apical pale markings . . . | <i>cornutus</i> , p. 396 |
| Markings of tergites mainly or entirely basal | 9 |
| 9. Middle femur with a pale stripe in front . . . | <i>edwardsi</i> , p. 397. |
| Middle femur not striped . . . | 10. |
| 10. Femora speckled with pale scales, especially anterior surface of mid-femur . . . | <i>stirns</i> , p. 398 |

- Femora without any sprinkling of pale scales 11
- 11 Mesonotum with light and dark scales mixed in varying proportions, sometimes forming an indefinite pattern, but at least with light scales round front margin
Mesonotal scales uniformly dark brown [Barraud, p 403.
[Walker, p 402;
vishnui, p 400,
tritorhynchus, [p 404.
mimeticus, p 409
12. Pale spot at middle of wing involving only costa and subcosta mimulus, p 412.
Pale spot at middle of wing usually extending over vein 1 nigrinus, p 413
- 13 Pleura devoid of scales, proboscis all black, ♂ palpi entirely dark scaled
Pleura with patches of broad scales, proboscis pale beneath in middle, last two segments of ♂ palpi with a white line beneath, or at least with a white spot at the base of each 14.
- 14 Fore and mid-femora and all tibiae conspicuously striped in front 15
Fore and mid-femora all dark in front 16
- 15 Hind femur with a brown line beneath on distal ½ theileri, p 414.
Hind femur pale beneath from base to knee. vagans, p 417.
- 16 Integument of pleura uniformly coloured, abdominal tergites basally banded 17
Integument of pleura with bare blackish-brown areas situated immediately above and below a conspicuous patch of white scales in middle 19.
- 17 Two terminal segments, or at least penultimate segments, of ♂ palpi with white line beneath 18.
Terminal segments of ♂ palpi with white basal spots beneath, very small species hutchinsoni, p 424.
- 18 Mid- and hind tibiae with more or less obvious pale stripes on outer side, abdominal bands white; a patch of white scales behind prothoracic spiracle unvittatus, p. 418
Mid- and hind tibiae dark, except small pale spot at tip, abdominal bands creamy, no postspiracular scales fatigans, p 420
19. Abdomen unbanded fuscocephalus, p 425
Abdomen banded fuscicornis, p 426

224. *Culex (Culex) epidesmus* (Theobald), 1910.

Rec. Ind. Mus iv, p 22 (*Teniorhynchus*) (♀) TYPE-LOC Bhogaon, Purnea dist., Bihar, x. 1908 (Paiva). TYPE ♀ in Ind. Mus.

Teniorhynchus luteoabdominalis Theobald, 1910, Rec. Ind. Mus iv, p. 23 (♀). TYPE-LOC.: Kathihar, Purnea dist., Bihar, x. 1908 (*O. Paiva*) TYPE ♀ in Ind. Mus.


Grabhamia ochracea Theobald, 1905 (neo Theobald, 1901), Journ. Econ. Biol., p. 25 (♀). TYPE-LOC.: India (*Chrysophers*) TYPE two co-type ♀♀ in Brit. Mus.

ADULT*.—Readily distinguished by the predominant yellow colour. Wing 4–5 mm.

* Edwards 1913 b, p. 231 (syn.), Brunetti 1920, p. 133 (*Grabhamia*); Barraud 1924 e, p. 982

♀.—*Head* dorsal surface covered with narrow yellow and numerous upright yellow and brown scales, broader pale yellow scales low down at sides. Palpi mottled with yellow and brown scales. Proboscis with a yellow band or ring in middle, usually very wide and occupying nearly $\frac{2}{3}$ of the length, but sometimes narrower, a dark area at base, usually narrow, and another dark area between pale band and labella, latter yellowish. *Thorax* mesonotum covered

Fig 90



♂ hypopygial structures (proctiger and phallosome) of *Culex*: a, *epidesmus*, b, *bitaeniorhynchus* (with variation in form of phallosome); c, *sinensis*, d, *cornutus*. Lettering as on p 4.

with narrow yellow, brown, and white scales, the last forming indefinite patches in front of wing-roots. Scutellar scales narrow and yellow. *Wings*: scales mainly dark, but scattered yellow scales along costa and vein 1. A pale yellowish area, sometimes indistinct, at apex of wing, including costa and veins 2 1, 2.2, and sometimes 3. *Legs*: femora and tibiae yellowish, with numerous scattered brown scales, tarsi brown, with rather broad basal and narrower apical yellow

rings Occasionally the legs are almost entirely yellow
Abdomen. tergites with narrow basal and much wider apical golden-yellow bands and median dark brown bands Amount of yellow scaling variable, occasionally dorsum is entirely yellow

♂ — Differs from ♀ as follows — Palpi longer than proboscis by nearly length of last two segments, two broad yellow rings on long segment, apical $\frac{1}{2}$ of this segment and last two segments with outstanding brown and yellow hairs, apical $\frac{1}{2}$ of terminal segment pale, the last two segments strongly curved upwards Mesonotal scales often paler, and pale ring on proboscis narrower, than in ♀ *Hypopygium* (fig 90, a) *Ip* with large ventral cornu and lateral process, median process divided into several rather long teeth, as in *cornutus* Paraproct with one large wide basal arm and another quite small

LARVA — Has not been isolated or described.

DISTRIBUTION — Widely spread from the PUNJAB to ASSAM and southwards to CEYLON, but apparently never very abundant It is, perhaps, commonest in BIHAR

Not known from outside the Indian region

225 *Culex (Culex) biteniorhynchus* Giles, 1901.

Journ. Bomb. Nat Hist Soc xii (May), p. 607 TYPE-LOC Travancore (*James*) TYPE non-existent

Teniorhynchus ayes Giles, 1901, Entom. xxiv (July), p. 196 (♂) TYPE-LOC Madras (*Cornwall*). TYPE: ♂ in Brit Mus

Teniorhynchus tenax Theobald, 1901, M.C. ii (November), p. 198 TYPE-LOC. Perak, Malay Penin., xi. 1899 (*Wray*) TYPE ♀ in Brit Mus.

Culex infula Theobald, 1901, M.C. i (November), p. 370 (♀). TYPE-LOC.: Taiping, Perak, Malay Penin., xi. 1899 (*Wray*) TYPE ♀ in Brit. Mus.

Grabhamia ambiguus Theobald, 1903, M.C. iii, p. 248 (♂). TYPE-LOC. Quilon, Travancore (*James*) TYPE ♂ in Brit Mus.

Grabhamia tanirostris Theobald, 1907, M.C. iv, p. 209 (♀) TYPE-LOC Peradeniya, Ceylon (*Green*) TYPE ♀ in Brit Mus

Teniorhynchus tenax var. *ocellata* Theobald, 1907, M.C. iv, p. 488 (♀) TYPE-LOC.: Kuching, Sarawak, Borneo (*Barker*). TYPE: ♀ in Brit Mus

Oculeomyia sarawaki Theobald, 1907, M.C. iv, p. 515 (♀) TYPE-LOC. Sarawak, Borneo (*Barker*). TYPE ♀ in Brit Mus


Of the above names *ambiguus* and *tenax* are regarded as varieties.

ADULT*.—Recognized by the speckled wings, banded proboscis, and apical yellow bands on abdominal tergites. Wing about 5 mm.

* Edwards 1913 b, p. 231 (syn.), 1921 c, p. 337; 1922 c, p. 282, 1924, p. 392, Barraud 1924 a, p. 984; Brug 1924 b, p. 17; Borel 1926, p. 81; Martini 1931, p. 362.

♀ —*Head* vertex with narrow golden-yellow and numerous upright yellow and brown scales, a small darker patch on either side, and broad white scales laterally. Palpi black, with pale scales apically. Proboscis black, with a broad median creamy band and pale scaling towards tip. *Thorax*: mesonotum covered with narrow golden-brown scales from front to about level of wing-roots, remainder covered with

Fig 91



Larval structures of *Culex* a-d, *cornutus*; e-h, *bitaeniorhynchus*.

brownish-black scales, except for a few golden scales in front of scutellum, usually a pair of submedian, more or less distinct, dark spots in middle of golden-brown area. Scutellum with dark brown narrow scales at base of mid-lobe, golden scales at apex and on lateral lobes. *Wings* veins clothed with a mixture of fairly broad pale yellow and black scales, giving a speckled appearance. *Legs*. brownish-black, plentifully sprinkled with pale scales on femora, tibiae, and first tarsal segments. tarsi dark, with narrow apical and

basal pale rings. *Abdomen*. tergites with moderately broad apical yellow bands. otherwise black, or with a sprinkling of pale scales

♂.—Differs from ♀ as follows.—Palpi longer than proboscis by rather more than length of apical segment, two pale rings on long segment, narrow pale rings at bases of last two segments, tip of last segment broadly pale. Last two segments upturned and with numerous outstanding hairs, which are mainly dark, except at tip of apical segment. Proboscis with narrower pale band, slightly noarser apex than base. *Hypopygium* (fig 90, b) *lp* variable in form; ventral cornu with a point directed ventrally, external process always large, lateral process fairly large, median process ending in a few small blunt teeth. Basal arm of paraproct comparatively small. Leaflet on subapical lobe of coxite also rather small, a second leaflet sometimes present

var. *ambiguus*—Differs from type-form in having fewer pale scales on wings, and many of the wing-scales are narrower. *Abdomen* with median basal yellow patches and apical lateral yellow patches not always forming complete bands

var. *tenax*—Similar to *ambiguus* as regards wing-scaling, but abdominal tergites are without basal yellow patches, the first few segments being almost entirely dark dorsally, other segments with apical yellowish lateral patches

Numerous other variations in abdominal markings occur, in some of which the abdomen is almost entirely covered with yellowish scales dorsally*. In one form from Kashmir the proboscis is almost entirely pale in ♀.

LARVA † (fig. 91, e-h)—Recognized by form of mentum—triangular, pointed, and furnished with a large number of minute teeth—differing in this respect from all other known larvae of Indian species of the genus. Length about 7 mm, including siphon, which is about 2 mm. long. Colour usually bright green. *Head* comparatively small, thinly chitunised,

* [The closely allied species or variety *aurantaper* Edw. (= *domesticus* Leio., nec Germar) has apparently not been found in the Indian area, although it occurs in the Malay Peninsula and East Africa. This form somewhat resembles var. *tenax*, but has mesonotal scales almost all black; first few abdominal tergites mainly or all black, last few entirely orange. Specimens of this form might easily be mistaken for *C. (Lutria) fusca*, just as the typical form of *C. biteniorhynchus* might be confused with *C. (L.) raptor* or *vorax*.]

† Christophers 1906, p. 11 (*C. tenax*), Edwards 1912 b, p. 380 (*ager* var. *ethiopioides*); Ingrain & Macfie 1919, p. 65 (var. *ethiopioides* and pupa, *ager*); Barraud 1923 a, p. 936, Borel 1926, p. 82, Senior-White 1927, p. 71; Martini 1931, p. 363.

and pale. Antenna pale on basal $\frac{1}{2}$, darker apically, tuft rather nearer base than apex of shaft, latter with 6-7 minute spinelets on basal $\frac{1}{2}$. Subapical bristles very near tip of shaft. Some long single hairs on prothorax, lateral hairs of thorax and abdomen moderately developed. Comb of 5-6 large sharp teeth irregularly arranged, each tooth with a few hairs at base, but no lateral denticles or fringe. Siphon pale, comparatively long and narrow, 6-8 times length of diameter at base. Four pairs of small subposterior hair-tufts fairly evenly spaced along apical $\frac{1}{2}$ of tube. Pecten of only 3-6 small transparent teeth, difficult to detect. Anal segment, with its hairs and papillæ, shown in fig 91.

Larva of var *ambiguus*.—Differs from that of type-form as follows.—Head, antennæ, and siphon darker. Antennal shaft with about double the number of spicules. Siphon usually rather shorter. Pecten of 6-9 teeth, the most distal tooth usually much detached from others.

HABITAT.—Open weedy pools. The larva usually feeds amongst green algæ, which it resembles in colour.

DISTRIBUTION.—One of the commoner mosquitoes over the whole of the Indian region. Its range extends from AFRICA, through the Oriental region, to JAPAN and AUSTRALIA.

226. *Culex (Culex) sinensis* Theobald, 1903.

Culex gelidus var *sinensis*, M C III, p 180 (♀). TYPE-LOC. Shaohying, China (Cornford). TYPE ♀ in Brit Mus.

Culex sepositus Leicester, 1908, Cul Malaya, p 162 (♀). TYPE-LOC Kuala Lumpur, Malay Penin (Leicester). TYPE non-existent.

Tæniorhynchus tenax Leicester, 1908 (nec Theobald), Cul Malaya, p 167 (♀). TYPE-LOC Kuala Lumpur, Malay Penin (Leicester). TYPE: ♀ in Brit Mus.

ADULT*—Resembles *tæniorhynchus* in general appearance, but may be distinguished by the entirely dark scaled wings. Wing about 4 mm.

♀—*Head*: pale narrow and upright scales on vertex, a darker patch on either side, followed lower down by a patch of broad white scales. Palpi black, with a few pale scales apically. Proboscis black, with a creamy ring in middle occupying about $\frac{1}{2}$ of the length. *Thorax*: mesonotum covered with pale ochreous or greyish-white scales from the front, continued back nearly to level of wing-roots, area behind thus deep brown, except for a few pale scales in front.

* Theobald 1910 b, p 313 (*Leucomyza*), Edwards 1913 b, p 231 (syn.), Barraud 1924 a, p 986, Brug 1924 b, p 22, Martini 1931, p 383.

of scutellum. Some specimens with a pair of submedian dark spots in the pale area, and a pair of submedian narrow bare lines continued back a short distance from front margin of mesonotum. Scutellum with dark brown scales at base of mid-lobe, light brown scales at apex and on lateral lobes. *Wings* dark scaled, outstanding plume-scales long and narrow. *Legs* dark brown, with lighter speckling on femora and on tibiae of fore legs. Tarsi with narrow pale rings which are mainly basal, but include tips of segments. *Abdomen*. tergites brownish-black, with narrow ochreous apical bands, also sometimes narrow pale basal bands. In some specimens the bands are incomplete or absent, the dorsum being almost entirely dark.

♂.—Differs from ♀ as follows.—Palpi longer than proboscis by rather more than length of last segment, long segment with one narrow pale ring and a broader one nearer the apex, apical $\frac{1}{2}$ with dark hairs on outside, last two segments with narrow basal pale rings, and dense dark hairs, except at tip of last segment, which is rather broadly pale. Proboscis with a narrower pale ring than in ♀, a little nearer apex than base. *Hypopygium* (fig 90, c) *lp* with simple thumb-like ventral corn, lateral process small, no median process. Basal arm of paraproct fairly large.

LARVA.—From two isolated skins examined this appears to be indistinguishable from the larva of *cornutus*. The anal papillae, which have collapsed, are rather longer, being $2\frac{1}{2}$ times length of chitinous part of anal segment, *asc* (present in one specimen only) have 2 branches instead of 4. Antennae, frontal hairs, mentum, comb, and siphon resemble those of *cornutus*. Possibly small differences could be found by comparing a long series of larvae of the two species.

HABITAT.—Rice-fields and large weedy pools.

DISTRIBUTION.—Not common in India, though widely distributed throughout the Oriental region. CENTRAL PROVINCES: Sonakhan*, Raipur dist., vii 1928 (*Semnor-White*). BIHAR: Kierpur* and Katihar*, Purnea dist (*Paiva*), Pusa*, various dates. ORISSA: Bahghai*, near Puri, 1908 (*Annandale*), and viii 1911 (*Annandale & Graveley*). BENGAL: Diamond Harbour*, xii 1919 (*Christophers*); Chandernagore (*Borel*). Sukna*, (*Sobha Ram*). ASSAM: Nongpoh*, Khasi Hills dist., xi 1921 (*Christophers*), vii 1922 (*Barraud*); Dibrugarh*, viii. 1922 (*Barraud*), Nalbari*, Kamrup dist., i. 1920 (*Taylor*). BURMA: Bhamo*, x 1931 (*Feegrade*).

Known also from MALAY PENINSULA and ARCHIPELAGO, CHINA, FORMOSA, and JAPAN.

227 *Culex (Culex) cornutus* Edwards, 1922.

Ind Journ Med Res x, p 283 (♂ & ♀) TYPE-LOC Tavargatti, Belgaum dist, Bombay Deccan, viii, 1921 (*Barraud*) TYPE ♂ & ♀ in Brit Mus

ADULT † —Wing about 3.5 mm

♀ —*Head* clothed dorsally with brown narrow curved scales, pale flat scales laterally. Proboscis brownish-black, with sharply-defined pale ring occupying about the middle $\frac{1}{3}$. Palpi brownish-black, with some pale scales apically. *Thorax*: mesonotum covered with dark brown, rather coarse narrow scales, those on posterior part lighter, a few rather longer and wider white scales in a small patch a little in front of each wing-root. Narrow brown scales on *apn* and *ppn*. Pleurae uniformly brown. *Wings* veins clothed with dark, rather long, narrow scales. *Legs* dark brown, femora and tibiae evenly sprinkled with pale scales, tarsi with pale apical and basal rings which are mainly basal. *Abdomen*: tergites dark brown, with narrow apical pale bands, more or less separated into median and lateral spots, also narrow basal pale bands on most of segments.

♂ —Differs from ♀ as follows —Palpi longer than proboscis by rather more than length of apical segment, last two segments and apical $\frac{1}{3}$ of long segment densely hairy. Two pale rings on long segment, last two segments upturned and with narrow pale ring at base of each. Tip of last segment narrowly pale. Pale ring on proboscis narrower. *Hypopygium* (fig. 90, d) · *lp* with several pointed processes directed dorsally, much as in *epidesmus*. Basal arm of paraproct very wide and long.

LARVA ‡ (fig 91, a-d) —Closely similar to *sinensis*, differs from *biteniorhynchus* as follows:—Subapical bristles on antenna some distance from tip of shaft, almost midway between tuft and apex. Mentum with much larger teeth towards apex, those towards base on either side very small. Comb of 4-6 large sharp teeth arranged more or less in a row. Pecten of only 2-4 very small teeth placed close together. Siphon with 5-6 pairs of small subposterior hair-tufts. A slightly hooked spine at tip of siphon anteriorly.

HABITAT.—Ground-pools, pools in jungle, fallow rice-fields, etc.

DISTRIBUTION —BOMBAY PRESIDENCY · Tavargatti*, type-locality, as given above, Deccan, Belgaum*, viii 1921 (*Barraud*), Bombay Harbour, Trombay*, and Hog Island*, vii 1921 (*Barraud*), North Kanara, Karwar (*Cogill*). MADRAS PRESIDENCY Ennore*, coast near Madras, xii. 1919 (*Christophers*)

Not known from other countries

† *Barraud* 1924 e, p 988

‡ *Barraud* 1923 c, p 937

228. *Culex (Culex) edwardsi* Barraud, 1923

Ind. Journ Med Res xi, p 507 (♀). TYPE-LOC Shillong, Khasi Hills, Assam, vi 1922 (*Barraud*) TYPE ♀ in Brit Mus

ADULT *—Resembles *sitiens*, but differs from that, and other allied species, in having the femora and tibiae of mid- and hind legs marked with distinct longitudinal pale stripes. Wing (♀) about 4.2 mm

♀.—*Head* vertex and nape covered with yellowish-brown narrow and upright scales. Palpi dark, with a few pale scales apically. Proboscis black, with a well-defined pale band in middle, occupying about $\frac{1}{3}$ of the length, rather wider below than above. *Thorax* integument of mesonotum brownish-black, with dark brown and yellowish-brown narrow scales, latter chiefly towards lateral margins and in front of scutellum. Narrow pale brown scales on *apn* and *ppn* and scutellum. Integument of pleurae pale brown, with several indistinct darker areas, and patches of broad white scales. *Wings* dark scaled, outstanding scales rather narrow, base of posterior fork-collar slightly nearer base of wing than that of anterior. *Legs* all femora broadly pale posteriorly; fore pair dark anteriorly, with numerous pale scales, sometimes tending to form patches, at about the middle, mid-pair dark anteriorly, with a well-defined pale longitudinal stripe running the whole length; hind pair broadly pale on basal $\frac{1}{2}$ on outer side, except dorsally, apical $\frac{1}{2}$ dark, with a longitudinal pale stripe to tip. Mid- and hind tibiae dark brown, with a pale line on outer side from base nearly to tip, fore tibiae pale, but without a definite stripe. Tarsi dark brown, with narrow but distinct pale rings which are mainly basal, but include tips of segments as well as apices of tibiae, segment 1 of hind tarsi with a pale line on outer side from near base to near apex.

♂.—Palpi longer than proboscis by rather more than length of rather long apical segment; long segment with two pale rings, that nearer base narrower than that nearer apex, a fairly long row of small hairs projecting downwards from underside of long segment, as well as numerous long hairs projecting laterally from apical part of this segment, last two segments with basal pale rings, and numerous outstanding dark hairs, except at tip of last segment, where they are pale. Proboscis with fairly wide pale band in middle, and some fairly long hairs projecting downwards from underside at about level of basal border of pale band. Pale ochreous or nearly white basal abdominal bands rather wider than in ♀, otherwise markings are very similar. *Hypopygium* (fig 92, c): *lp* resembling *sitiens* and *cornutus*; median process with

* Barraud 1924 c, p. 995.

rather large teeth, external process pointed and caudally directed Paraproct with numerous spines at crown and a long pointed basal arm

LARVA —Unknown

DISTRIBUTION.—ASSAM Shillong *, as given above. CENTRAL INDIA Amol Nullah *, near Birsinghpur, Rewa State (*Senior-White*) SOUTH INDIA Mercara * and Virajpet *, Coorg, vi 1927 (*Baily*)

Not known from elsewhere

229 *Culex (Culex) sitiens* Wiedemann, 1828

Aussereur Zweifl Insec. 1, p 542. TYPE-LOC . Sumatra. TYPE : non-existent

Culex impellens Walker (nec Theobald), 1860, Proc. Linn. Soc Lond. iv, p 91 (♀). TYPE-LOC . Makassar, Celebes (*Wallace*). TYPE : ♀ in Brit Mus

Culex macroannulatus Theobald, 1901, M.C 1, p 353 (♂ & ♀) TYPE-LOC Quilon, Travancore, S India, in 1900 (*Jones*). TYPE ♂ & ♀ in Brit Mus

Culex nigricephala Leicester, 1908, Cul Malaya, p 149 (♂ & ♀) TYPE-LOC Batu Gajah, Malay Penin (*Daniels*). TYPE : ♂ & ♀ in Brit Mus

ADULT †—A rather larger and darker mosquito than *vishnui* and allied species. Wing about 4 mm


♀ and ♂.—General coloration brownish-black Head with narrow brown and upright black scales on vertex and nape, and broad pale scales low down at each side. Proboscis black, with a creamy band in middle, of about same width in the two sexes Palpi of ♀ black, with white scales apically; in ♂ with 2 pale rings on long segment, narrow basal pale rings to last two segments, tips rather broadly pale, otherwise outstanding hairs on last two segments are dark; a row of stiff translucent hairs projecting downwards from underside of long segment, as in *vishnui* Thorax . mesonotum covered with dark brown narrow scales, usually some paler scales towards lateral margins. Scutellar scales narrow and pale brown. Wings scales dark and moderately broad; fork-cells rather short, base of anterior slightly nearer wing-tip than that of posterior Legs . brownish-black, anterior surface of femora speckled with pale scales, most marked on mid-legs, mid- and hind tibiae with a pale line on the outside, not always very pronounced. Tarsi with distinct, but rather narrow, basal and apical pale rings. Abdomen : tergites brownish-black, with rather narrow basal white bands ♂ hypopygium (fig. 92, b) lp with a larger external

† Edwards 1913 b, p 232 (syn.); 1924, p 394; Barraud 1924 c, p 993, Brug 1924 b, p. 13, Borel 1926, p. 91

process than in *vishnu* and allies, median process with several teeth, one larger and longer than others

LARVA * (fig 93, a) —Antenna, mentum, siphon-tube, and anal segment as shown in figures. Distinguished from other known larvæ of this group by the comparatively short and wide siphon, which is only 2-2½ times length of diameter at base, and by the short anal papillæ †. There are usually

Fig 92



♂ hypopygial structures of *Culex*: a, *vishnu*; b, *stans*, c, *edwardsi*, d, *whitet*. Lettering as on p. 4.

6 pairs of tufted hairs on the siphon, one of these being lateral, the others near posterior border. The pecten-teeth have numerous denticles along one side.

HABITAT.—Brackish pools near sea-coast.

* Barraud 1924 m, p. 427; Cooling 1924, p. 36; Hill 1925, pl. viii; Borel 1926, p. 92; Buxton & Hopkins 1927, p. 79.

† [This larva is also distinguished rather sharply from all other Indian species of *Culex* by the remarkably thick preopyleal spines. Specimens in the British Museum from Fiji, South China, Bombay, and East Africa all show this feature.]

DISTRIBUTION—Common on sea-coast, and probably occurring all round Indian coast-line. Not usually found far inland.

The range extends from EAST AFRICA, throughout the whole of the ORIENTAL REGION, to AUSTRALIA and FIJI.

230 *Culex (Culex) vishnui* Theobald, 1901*

M.C. 1, p. 355 (♀ only). TYPE-LOC. Madras (Cornwall) TYPE ♀ in Brit Mus

Culex impellens Theobald, 1901 (nec Walker), M.C. 1, p. 362 (♀)

var *perplexus* (*Culex perplexus*) Leicester, 1908, p. 150 (♂ & ♀)
TYPE-LOC. Kuala Lumpur, Malay Penin (Leicester) TYPE ♂ & ♀ in Brit Mus

ADULT †—A very common small brown mosquito, with a pale band on the proboscis, distinguished from several closely allied species by structure of ♂ hypopygium, larval characters, and scaling of mesonotum and of ♂ palpi. Wing about 3-3.5 mm.


♀—*Head*. narrow and upright pale brown scales on vertex and nape (in some specimens scales are brown or dark brown). Palpi dark brown, with usually some pale scales at tips. Proboscis dark brown, with a fairly distinct pale band, rather nearer apex than base, sometimes small areas of pale scaling may be present between pale band and base. *Thorax*. mesonotum clothed with golden-brown, or pale brown, narrow scales, in some specimens there is a mixture of rather dark brown and pale scales, latter forming a more or less distinct pattern. *Wings*. dark scaled, outstanding series narrow, base of anterior fork-cell usually very slightly nearer base of wing than that of posterior. *Legs*. dark brown, tarsi with very narrow, often indistinct, pale rings, which are mainly basal, but involve apices of some segments. Fore and mid-femora dark brown on anterior surface, without any speckling of pale scales. Hind femur with pale scaling on outer side from base nearly to knee, and with sharply-contrasted dark scaling dorsally from near base to knee. Mid- and hind tibiae with more or less defined pale stripe on outer side for nearly whole length. *Abdomen*. tergites dark brown or black, with well-marked ochreous basal bands.

* *C. summorosus* Dyar, of the Philippine Islands, considered by Edwards to be synonymous with *vishnui*, appears to me to be distinct from examination of adults and larval skins sent to me from those islands. The adults agree very closely with *vishnui*, but the larval skins show definite structural differences. This form has so far not been met with in India.

† Edwards 1913 b, p. 233 (syn.), 1917, p. 225 (syn.), 1921 c, p. 339, Barraud 1924 s, p. 996, Brug 1924 b, p. 15, Borel 1926, p. 93, Martini 1931, p. 392.

♂—Palpi longer than proboscis by rather more than length of last segment, dark brown, with two pale rings on long segment, and a row of stiff translucent hairs projecting downwards from beneath, in addition to some long hairs projecting laterally. Pale rings at bases of last two segments,

Fig 93.



Larval structures of *Culex* (antenna, mentum, tail-end, enlarged pecten- and comb-teeth): a, *sitiens* (extra figure showing thickened pre-clypeal spines); b, *barraudi*; c, *vishnui*.

tip of last segment and hairs pale; otherwise outstanding hairs are dark. Bases of fork-cells of wing about level. *Hypopygium* (fig. 92, a): *lp* with median process divided into teeth directed caudally, one tooth much longer than

others and projecting some distance beyond plate Paraproct with basal and lateral arms

LARVA * (fig 93, c) head comparatively large *Antenna* of moderate length, with tuft at $\frac{2}{3}$ of length from base, and with preapical bristles slightly below tip Shaft of antenna dark at base and on part beyond tuft; fairly numerous spinelets along nearly whole length *Frontal hair* A conspicuously dark Teeth of mentum rather large and all about same size, 6-8 on either side of central one Comb of 6-7 large sharp teeth set in an irregular row (differing from the larvæ of *tritænorhynchus*, *barraudi*, and *whitei* in this respect) *Siphon* 4-6 times length of diameter at base, with 6-7 pairs of rather small tufted hairs along posterior border, and two small pairs of lateral hairs of 2-3 branches Pecten of 9-11 teeth, the distal teeth large and curved and about same size as comb-teeth, *sc* 3-branched, *lh* very small, with several branches Dorsal pair of anal papillæ about length of longest fan-hairs, ventral pair a little shorter

HABITAT—Ground-pools of almost every description, rice-fields, salt marshes, etc

DISTRIBUTION—One of the commonest of Indian mosquitoes From examination of ♂ hypopygia and of larvæ it has been ascertained that this species is present from the NORTH-WEST FRONTIER to ASSAM and BURMA, and through PENINSULAR INDIA to CEYLON. It is less common at high elevations, and in the Western Himalayas at altitudes of over 5,000' it is usually replaced by *barraudi*, and in parts of the Assam hills by *whitei*

Its range extends from MESOPOTAMIA to CHINA and JAPAN, and throughout the Oriental region as far south-east as NEW GUINEA

231 *Culex* (*Culex*) *whitei* Barraud, 1923

Ind. Journ. Med. Res. xi, p 508 (♂ & ♀). TYPE-LOC. Haflong, Cachar Hills, Assam, viii 1922 (*Barraud*) TYPE ♂ & ♀ in Brit. Mus.

ADULT †—Resembles *vishnui* and *barraudi* very closely, but differs from both in structure of ♂ hypopygium, from the former on larval characters and from one or both in the following points—Palpi of ♂ with a row of stiff translucent hairs projecting downwards and inwards, in addition to some long hairs at apex projecting laterally, as in *vishnui* and *sitiens* (there is no row of stiff hairs in *barraudi* and *tritænorhynchus*). Bases of anterior and posterior fork-cells

* Barraud 1923 c, p 938, Borel 1926, p. 95, Senior-White 1927, p 71, Martini 1931, p 392

† Barraud 1924 c, p 908

usually about level in both sexes, but, if not, that of the posterior is slightly nearer base of wing than that of anterior

♂—*Hypopygium* (fig 92, d) *lp* with more numerous and shorter teeth than in allied species (compare with figs 92, a-c, & 94, a, b)

LARVA.—From one incomplete larval skin, from which a ♂ of this species resulted, it can be seen that in most structural details the larva resembles *barraudi* and *triteno-rhynchus*, differing from *vishnui* in having numerous small teeth in the comb arranged in a triangular patch. Further material is required before a detailed description can be given

HABITAT.—Ground-pools, rice-fields, etc

DISTRIBUTION (as checked by author by examination of ♂♂)—ASSAM: Haflong, as given above, Shillong and Nongpoh, Khasi Hills (list, vi. & vii. 1922 (*Barraud*))

Also Chiangmai, SIAM, xii 1930 (*Sinton*)

232. *Culex (Culex) barraudi* Edwards, 1922 †.

Ind Journ Med Res x, p 284 (♂ & ♀). TYPE-LOC. Mahdopur, Punjab, iv 1921 (*Barraud*) TYPE: ♂ in Brit. Mus.

ADULT.—Resembles *vishnui* very closely, but differs in structure of ♂ hypopygium, in larval characters, and in absence of a row of stiff hairs on underside of long segment of palp of ♂. Base of posterior fork-cell of wing usually slightly nearer base of wing than that of anterior, though sometimes the bases may be about level.

♂—*Hypopygium* (fig. 94, a) *lp* with about 5 teeth, all about same length and evenly spread out. Basal arm of paraproct long and pointed

LARVA ‡ (fig. 93, b).—Differs from *vishnui* in several characters, chiefly in having numerous small teeth in the comb, arranged in a triangular patch. Resembles *triteno-rhynchus* very closely, differing as follows:—Teeth of mentum of more irregular sizes; siphon usually longer, *lh* of 2 fairly long fine branches; siphon with usually only 3 pairs of sub-posterior tufted hairs and one lateral hair.

HABITAT.—Ground-pools, marshes, etc.

DISTRIBUTION.—PUNJAB: Mahdopur*, type-locality, as given above, Pathankot* and Amritsar*, iv. & v. 1921 (*Barraud*); Kalka*, iv 1922 (*Barraud*). WESTERN HIMALAYAS: Kasauli*, 5,000', 1921–23 (*Christophers & Barraud*), Murree*, 7,000', 1922 (*Hill*). SOUTH INDIA: Karikal, Tanjore (*Borel*). CEYLON (*Senior-White*)

† *Barraud* 1924 e, p. 997.

‡ *Barraud* 1923 c, p 939; *Senior-White* 1927, p. 71.

* ♂♂ examined by Author.

233. *Culex (Culex) tritaeniorhynchus* Giles, 1901.

Journ Bomb Nat Hist Soc xiii, p 606 (May) TYPE-LOC - Travancore, South India (*James*). TYPE ♀ in Brit Mus


Culex vishnu Theobald, 1901, M.C 1, p 355 (♂ only) TYPE-LOC Quilon, Travancore (*James*) TYPE ♂ in Brit Mus

Culex annulus Theobald, 1901, M.C 1, p 358 (November) (♀) TYPE-LOC : Tai Po, Pokfulam, Hong Kong (*Rees*) TYPE ♀ in Brit Mus.

Culex sitrens Theobald, 1901 (nec Wiedemann), M.C 1, p 360 (described as *C sitrens* Wied from ♀♀ from Taiping, Perak, Malay Penin (*Wray*))

Culex buror Theobald, 1905, Ann Mus Nat Hung iii, p 82 (♂ & ♀) TYPE-LOC : Bombay, 1902 (*Biró*) TYPE ♂ & ♀ in Nat Mus. Hung, Buda Pest


Fig 94



♂ hypopygial structures (proctiger and phallosome) of *Culex* : a, *barraudi*, b, *tritaeniorhynchus*; c, *whitmorei*; d, *gelidus* (with style on smaller scale), e, *mimeticus*, f, *mimeticus* var. *mimuloides* (side view of phallosome only); g, *mimulus*.

ADULT *—Very similar in appearance to *vishnu*, *barraudi*, and *whitei*, but usually rather smaller, and differs from some or all of these three as follows—Mesonotum uniformly clothed with dark brown scales. Femora and tibiae dark brown, except undersides of femora, hind femur pale dusky from base to tip on outer side, slightly darker dorsally (in allied species the dark and light areas are usually more sharply defined), tibiae without any indication of pale stripes. Palpi of ♂ dark at tip, and without a row of stiff translucent hairs on underside of long segment. Pale band on proboscis of ♀ often extended towards base on underside, and occasionally on to sides and upper surface. Base of

Fig 95



Larval structures of *Culex tritaeniorhynchus* Giles: a-c, variety (?) from Karwar, d-f, typical form.

anterior fork-cell in wing of ♀ usually nearer base of wing than that of posterior cell. *Hypopygium* (fig. 94, b): *lp* of characteristic form (compare with figs. 92, a, d, & 94, a).

LARVA † (fig. 95)—Very similar to *barraudi* and *whitei*, differing from former in shape of mentum and in length of *lh* (compare figs. 93, b, & 95). There are usually five pairs of hair-tufts along the posterior border of siphon and

* Giles 1901 b, p. 192; Edwards 1913 b, p. 233 (syn.); 1917, p. 224; 1921 c, p. 339; Barraud 1924 c, p. 995; Brug 1924 b, p. 16; Kirkpatrick 1925, p. 111; Borel 1926, p. 96; Martini 1931, p. 387.

† Ingram & Macfie 1917 b, p. 149, and p. 86 (pupa); Edwards 1921 c, p. 339; Barraud 1923 c, p. 849; Kirkpatrick 1925, pp. 114, 115 (and pupa); Borel 1926, p. 96; Martini 1931, p. 388.

one lateral pair. Some variation has been noted, in fig 95 d-f represent the form regarded as typical, a-c another form found at Karwar.

HABITAT —As given for *vishnu*

DISTRIBUTION.—As widely spread in India as *vishnu*, but usually less common than that species

Range extends from the MEDITERRANEAN to CHINA and JAPAN, and from WEST AFRICA and EGYPT, through the ORIENTAL REGION, as far south-west as CELEBES

234 *Culex (Culex) whitmorei* (Giles), 1904

Journ Trop Med vii, p 367 (*Tæmorhynchus*) (♀). TYPE-LOC Philippine Is (*Whitmore*). TYPE ♀ in Brit Mus.

Tæmorhynchus argenteus Ludlow, 1905, Can Ent xxxvii, p 98 (♀) TYPE-LOC Luzon, Philippine Is (*Whitmore*) TYPE ♀ in US Nat Mus

Leucomyia plegipennis Theobald, 1907, M C iv, p 375 (♀). TYPE-LOC. Kobe, Japan (*Cornford*) TYPE ♀ in Brit Mus

Culex albus Leicester, 1908, Cul Malaya, p 148 (♀) TYPE-LOC. Kuala Lumpur, Malay Penn (*Leicester*) TYPE ♀ in Brit Mus

Culex loriscatus Leicester, 1908, Cul Malaya, p 151 (♀) TYPE-LOC Kuala Lumpur, Malay Penn (*Leicester*) TYPE ♀ in Brit Mus

ADULT * —A rather small mosquito with white scaling over larger part of mesonotum. Resembles *gelidus*, but differs as stated in key. Wing about 3.5 mm

♀ —**Head**. vertex and nape covered with narrow and upright white scales, darker scales towards sides, and broader white scales laterally. Palpi dark, with white scaling apically. Proboscis brownish-black, with a wide pale ring or band occupying the middle $\frac{1}{3}$. **Thorax**. mesonotum covered with white or greyish-white scales from front as far back as level of wing-roots, the pale area continued posteriorly in four lines of yellowish-white scales to scutellum, spaces between lines brownish-black. Scutellum clothed basally with dark scales, apically with pale scales. **Wings** median series of scales rather broad and dark, particularly on costa, subcosta, and veins 1, 3, and 5. **Legs**. dark brown; femora speckled with pale scales, mid- and hind tibiae more or less lined with pale scaling on outer side. Tarsi with well-marked basal pale rings, not involving the apices of segments. **Abdomen** tergites brownish-black, with narrow yellowish basal bands, usually triangularly produced in middle nearly to apical margin of each segment

♂. —Palpi longer than proboscis by rather more than length of apical segment. Long segment with a narrow pale ring near base and a wider ring nearer apex, last two segments

* Edwards 1913 b, p 232 (syn.), 1917, p 225 (syn.); Barraud 1924 c, p 988, Brug 1924 b, p 21, Martini 1931, p. 393.

with narrow basal pale rings and outstanding dark hairs, except at tip of last segment, where they are pale. Pale band on proboscis narrower than in ♀, and rather beyond middle. *Hypopygium* (fig. 94, c). *lp* with ventral cornu and large external process, median process with four subequal teeth, no lateral process. Basal arm of paraproct small.

LARVA * (fig. 96, g-k).—*Antenna* unusually long, shaft with spicules, tuft at $\frac{3}{4}$ from base, preapical bristles slightly below tip, one apical bristle much longer than other. *Mentum* with large regular teeth, 6-7 either side of central one. *Comb* of about 7 large sharp teeth set in an irregular row. *Siphon* unusually small and curved forwards, 4-5 times length of diameter at base, 5-6 pairs of 2-branched, fairly long, subplumose hairs along posterior border; 2 pairs of lateral 2-branched shorter hairs. *Pecten* of 8-14 teeth, with lateral denticles, a pair of fairly long bristles at apex of siphon, with hairs around their bases. *Anal segment* both *osc* and *asc* single and long, *lh* rather small and of several branches. Both pairs of papillae pointed and as long as siphon. Fan-hairs rather shorter than papillae.

DISTRIBUTION.—Although this was not recorded from India before about 1923, it is widely spread and common from the PUNJAB to ASSAM and BURMA, and through PENINSULAR INDIA to Ceylon. It has been found up to 7,500' in the Nilgiri Hills, SOUTH INDIA.

Known also from MALAY PENINSULA and ARCHIPELAGO, PHILIPPINES, WEST CHINA, and JAPAN.

235 *Culex (Culex) gelidus* Theobald, 1901.

M.C. II, p. 20 (♀). TYPE-LOC. Selangor, Malay Penm., x 1890 (Butler). TYPE: ♀ in Brit. Mus.

var. *cuneatus* Theobald, 1901, M.C. II, p. 22 (♀). TYPE-LOC. Taiping, Perak, Malay Penm., XII. 1899 (Wray), and Quilon, Travancore (James). TYPE. 2 co-type ♀♀ in Brit. Mus.

var. *bipunctatus* (of "*Leucomyia gelidus*") Theobald, 1907, M.C. IV, p. 374 (♂). TYPE-LOC: India (Aldrich), Sarawak, Borneo (Barker). TYPE. ♂ in Brit. Mus.

ADULT ♀ and ♂ †.—Very similar to *whitmorei* in size and general appearance, but differs as follows:—Anterior white area of mesonotum not produced in lines to scutellum, the space behind level of wing-roots being entirely brownish-black. Pale band on proboscis of ♀ usually narrower and rather further from base. Wings with scales in median series smaller and narrower. Pale bands on tarsi narrower.

* Barraud 1923 c, p. 941; Senior-White 1927, p. 71; Martini 1931, p. 394.


† Theobald 1907, p. 372 (*Leucomyia*); Barraud 1924 c, p. 990. Brug 1924 b, p. 19; Borel 1926, p. 88.

and involving apices as well as bases of segments. Femora not speckled and tibiae not lined with pale scales, uniformly dark in front and above.

♂.—*Hypopygium* (fig 94, d) ventral cornu of *lp* with a pointed tip directed dorsally, median process with 2 teeth pointing downwards, lateral process well developed, external process rounded. Basal arm of paraproct long. Style expanded in middle, abruptly narrowed at tip, a small terminal appendage.

var *cuneatus* —The commonest form, in which the basal pale abdominal bands are triangularly produced in the middle

Fig 96



Larval structures of *Culex*. a-f, *gelidus*, g-k, *whitmorei*
(c, j, comb-teeth, d, i, pecten-teeth)

var. *bipunctatus* —Resembles var *cuneatus* except that there is a pair of submedian dark spots in the white-scaled area of mesonotum.

LARVA * (fig 96, a-f).—Remarkably distinct from that of *whitmorei*. Antenna comparatively short; tuft at $\frac{2}{3}$ from base, subapical bristles very near tip of shaft; part of latter basal to tuft with small spinelets. Mentum with 9-10 rather small teeth either side of central one. Comb of about 35 small teeth in a triangular patch. Siphon widest

* Barraud 1924 m, p 428, Borel 1926, p 90.

towards middle and slightly tapering towards apex, 3-4 times length of diameter at base. Four pairs of 4-5-branched hairs along posterior margin, no lateral tufts. Pecten of 9-11 moderately large teeth, each with 4-5 sharply-pointed lateral denticles. *Anal segment* enclosed in smooth chitinous ring, *isc* 2-branched, *lh* single, moderately long. Dorsal papillæ about length of the rather short fan-hairs; ventral papillæ shorter.

HABITAT.—Ground-pools, usually those containing much weed, marshy tracts, etc.

DISTRIBUTION.—Common over the larger part of the Indian region from PUNJAB to ASSAM and BURMA, and southwards to CEYLON.

Also found in MALAY PENINSULA, many of the islands of the DUTCH EAST INDIES, and PHILIPPINES, SIAM, COCHIN CHINA, and FORMOSA.

236 *Culex (Culex) mimeticus* Noé, 1899.

Bull. Soc. Ent. Ital. xxxi, p. 240 (♀). TYPE-LOC. Basilicata, Italy. TYPE in Rome Univ. Mus.

var. *mimuloides* Barraud, 1924, Ind. Journ. Med. Res. xi, p. 902 (vide below). TYPE-LOC. Nilgiri Hills, S. India.

ADULT *—Distinguished from all other known species of Indian *Culex*, except *mimulus*, by the spotted wings, which resemble those of some species of *Anopheles*†. Wing about 4.5-5 mm.

♀.—**Head** vertex and nape covered with pale brown narrow scales and brown and black upright scales. Palpi black, narrowly pale apically. Proboscis black, with a well-defined median creamy ring. **Thorax**: mesonotum covered with bright brown narrow scales and with usually a mixture of lighter scales forming indefinite patches round front margin and (a pair) in middle; *apn*, *ppn*, and scutellum with narrow


* Edwards 1921 c, p. 337, Barraud 1924 e, p. 901, Borel 1926, p. 83; Martini 1931, p. 371.

† [In addition to *C. mimeticus* and *C. mimulus*, two other species of this group are known, both differing from either of the Indian species in the ♂ hypopygium and also in wing-markings. These are: (1) *C. diengensis* Brug., of Mt. Dieng, Java, which resembles *mimeticus* in that the middle costal spot involves only the costa and subcosta (not vein 1), but veins 3 and 5.1 are pale practically to the tips, and 5.2 has a small pale spot at tip, ♂ style remarkably broad; and (2) *C. orientalis* Edw., of Japan and China, with middle costal spot not only involving vein 1 (as in *mimulus*), but even spreading across vein 4; tip of 5.2 broadly pale, and often a pale area on 5 before the fork, also (in ♀) a small pale area at base of 1, ♂ coxite with tip narrowed and provided with a tuft of short hair.

For further notes on the species of this group, vide Appendix.]

pale brown scales. *Wings* marked with light and dark scales, the former forming spots, as follows —A small pale spot on costa and subcosta [but not involving vein 1] at about middle of wing, another larger spot immediately beyond subcostal junction, involving costa (sometimes tip of subcosta) and vein 1, and one near tip of wing on costa, vein 1, and 2 1. Vein 3 usually extensively pale, but length of pale area variable [tip always dark]. Small spots at bifurcations of veins 2 and 4. A pale area in middle of 5 1 [leaving tip

Fig 97



Larval structures of *Culex* a-c, *mimeticus*, d-g, *mimulus* (with variations in length of siphon-tube)

broadly dark], and broad pale area on basal $\frac{1}{2}$ of 6 [usually leaving a short dark area at base of vein, 5 and 5 2 entirely dark]. *Legs*: brownish-black, marked with pale scaling. Femora pale posteriorly and ventrally for nearly whole length from base, and with pale knee-spots. Fore and mid-tibiae striped with pale scaling on outer sides for whole length; hind tibia with similar stripe on middle $\frac{3}{4}$. Tarsal segment 1 on all legs indistinctly striped with pale scaling, narrow

pale rings over tibio-tarsal joint and over joints between tarsal segments. *Abdomen* tergites brownish-black, with narrow basal pale bands.

♂ —Palpi with one narrow and one broad pale ring on long segment, narrow basal pale rings to last two segments, and outstanding dark hairs, except at tip of last segment, which is narrowly pale. Proboscis with narrower pale ring than in ♀, and pale spots on wings usually less pronounced. *Hypopygium* (fig 94, e). *lp* with 3 straight teeth all about same length, and one smaller, directed outwards. Basal arm of paraproct well developed, but rather short.

var. *mimuloides* —Some specimens from the Nilgiri Hills are intermediate between typical *mimeticus* and *mimulus*, both in markings and in structure of ♂ hypopygium, but show variation in the former. In some specimens the wing-markings are very similar to the Himalayan form, in others the pale area on vein 3 is much reduced, and in some there is a small pale spot at tip of 5.2. Two ♀♀ have a pale spot in the middle of vein 4.2. *Hypopygium* (fig 94, f) *lp* usually with 5 stout teeth, three of which are curved and more caudally directed than in the Himalayan form, and basal arm of paraproct longer, being nearly length of plate. Larvæ have not yet been isolated.

LARVA *, Himalayan form † (fig 97, a-c) —*Antenna* dark towards apex, otherwise pale, small spinelets on basal $\frac{2}{3}$ of shaft, tuft at $\frac{2}{3}$ from base, subapical bristles some little distance below tip of shaft, but nearer to it than to the tuft (about $\frac{1}{3}$ – $\frac{2}{3}$ of the distance between tuft and apex). [Pre-clypeal spines stout and dark.] *Mentum* with 7–9 fairly large regular teeth either side of central one. *Comb* a triangular patch of about 35 teeth, each having a long point and lateral fringe. *Siphon* $4\frac{1}{2}$ –7 times length of diameter at base, with 8–10 hairs of 4–5 branches, which are twice length of diameter of siphon in middle and sometimes arranged in 4–5 pairs, arising between pecten and apex of tube, those nearest apex usually 2–3-branched. *Pecten* of 12–23 long narrow teeth, with hair-like lateral denticles. *Anal segment* with its hairs and papilla as shown in figure.

DISTRIBUTION.—Common along HIMALAYAS, to ASSAM hills, and highlands of BURMA and COCHIN CHINA. Also in hills of SOUTH INDIA and CEYLON.

* Christophers 1906, p. 9; Barraud 1928 c, p. 941; Ségué 1925 b, p. 16; Borel 1926, p. 83; Martini 1931, p. 373. Pupa:—Theodor 1924, p. 341.

† [Larval skins in the British Museum from Palestine and Hong Kong agree with Barraud's description and figures.]

Known also from MEDITERRANEAN REGION, CHINA, and FORMOSA. Like other hill-species, it is occasionally found in the plains of India.

237 *Culex (Culex) mimulus* Edwards, 1915.

Bull Ent Res v, p 284 (♂ & ♀) TYPE-LOCALITY Sarawak, Borneo (Sarawak Mus Coll) TYPE ♂ & ♀ in Brit Mus

ADULT * —Very similar to *mimeticus*, but rather smaller, [pale spot at middle of costa often larger, and in any case almost invariably involving vein 1], vein 3 usually entirely dark scaled, though occasionally there is a small [or large] pale area in the middle of that vein (especially in ♀) [Vein 5 (with 5 2) entirely dark, as in *mimeticus*] Some specimens of var *mimuloides* of *mimeticus* are intermediate, both in wing-markings and in structure of ♂ hypopygium, between typical specimens of the two species

♂ —*Hypopygium* (fig 94, g) · *lp* very similar to that of *vishnui* (compare figs 93, e, & 94, g), [with 5-6 curved teeth arranged like the spread fingers of a hand, and for the most part directed backwards, above the base of this group of teeth is a cluster of small denticles Basal arm of paraproct considerably longer than in *mimeticus*, style broader].

LARVA † (fig 97, d-g.) —Resembles *mimeticus* in general appearance, but differs as follows ‡ —Subapical bristles on antenna slightly nearer tip of shaft Teeth of mentum less regular in size Comb-teeth with more regular fringe, terminal point scarcely differentiated from fringe Siphon with 5-6 pairs of short branched hairs along posterior border, which are little, if any, longer than diameter of siphon in middle (instead of 8-10 unpaired and much longer branched hairs), and one pair of short lateral branched hairs. Pecten-teeth stouter, with stronger lateral denticles; *lh* with about 3 short branches. Siphon variable in length (as shown in fig. 97, g), 6-9 times length of diameter at base.

DISTRIBUTION —Fairly common from UNITED PROVINCES (Dehra Dun) to ASSAM and BURMA, and through PENINSULAR INDIA to CEYLON

Known also from the MALAYAN REGION to NORTH AUSTRALIA and HONG KONG

* Edwards 1924, p 394, Barraud 1924 c, p. 993; Brug 1924 b, p 11, Borel 1926, p 86

† Barraud 1923 c, p 942, Borel 1926, p 87; Senior-White 1927, p 71

‡ [Specimens in the British Museum from Ceylon and Hong Kong agree fairly closely with Barraud's description and figures. Those from the latter locality were obtained in 1933 by Dr. R. E. Jackson, together with *C. mimeticus* and a third form of this group, described as new in the Appendix.]

238. *Culex (Culex) nilgiriensis* Edwards, 1916.

Bull Ent Res vi, p 358 (♂ & ♀) TYPE-LOC Ootacamund,
Nilgiri Hills, South India, 7,500'. xu 1913 (Fletcher) TYPE.
♂ & ♀ in Brit Mus

ADULT *—The original description is as follows (the words in square brackets are mine) —

“Male and female.—*Head* with dark brown upright scales, creamy white flat and narrow curved ones at the sides and in the middle respectively, the whitish more numerous in proportion to the dark brown ones in the male than in the female, and the flat ones extending higher up, the flat ones tend to form a line round the upper part of the eye margins, but in neither sex do they actually reach the middle line. A row of dark bristles round the eyes, of which a pair on the vertex are much longer than the others, at least in the female. Palpi and proboscis entirely black scaled in both sexes, male palpi longer than the proboscis by their terminal joint, basal segment without any row of projecting scales, but with some pale shortish hairs and with a few long dark ones towards the tip; second and third joints with numerous dark long hairs, the second nearly five times as long as the labella, the third about one quarter longer than the second. *Thorax*: prothoracic lobes reddish brown, apparently without scales, scutum (mesonotum) reddish brown, slightly pruinose, clothed with reddish brown scales; scutellum pale, almost whitish, clothed with narrow pale brown scales, postnotum light reddish brown. Pleurae almost uniformly pale, unscaled. *Abdomen* blackish brown above, the segments with white basal bands of even width, broader in the male than in the female, venter almost uniformly pale. *Male genitalia*: side-pieces [coxites] of the usual *Culex* form without special modification, leaf-like appendage very narrow. Claspers moderate, smooth, gently curved, rather abruptly narrowed a short distance before the tip (as in *Culiciomyia*), a single fine hair arising from the point of narrowing. Unci [parameral plates] with pointed tips. Harpagones [lateral plates of the phallosome] apparently undivided, but provided with a rather long, pointed, downwardly extending projection, somewhat jagged on its lower edge (as in *Culiciomyia*). Harpes [paraprocts] short and broad, without basal projection, their tips

* Barraud 1924 g, p. 1272. [This species is still only known from the type ♂ and ♀. It is somewhat out of place in the subgenus *Culex*, as in spite of the absence of erect translucent scales on the ♂ palpi and of a subapical spiny crest on the ♂ style, it obviously has more affinity with the subgenus *Culiciomyia*, agreeing with the Indian species of that subgenus in form of phallosome and absence of scales on pleurae. It is quite distinct from any of the species included by Barraud in *Culiciomyia*, approaching most nearly to *C. bairdii*.]

broad and provided with shorter and less numerous spines than usual. *Legs* dark scaled, the under sides of the femora lighter, claws of the four anterior legs of the male each with a well-marked tooth. *Wings* rather thinly clothed with dark scales, those in the lateral series linear, few in number, except towards the tip of the wing. Fork-cells very long, yet scarcely twice as long as their stems, the upper with its base slightly nearer the apex of the wing than that of the lower. Cross-veins separated by about twice their own length. Halteres light brown.

"Length of body, 6 mm., male wing, 4.5 mm., female wing, 5 mm. Madras: Utakamand [Ootacamund], Nilgiri Hills, 7,500 ft., 24th-31st December, 1913 (*T. Barnbrigge Fletcher*)."

LARVA —Unknown

DISTRIBUTION —Known only from type-locality

239 *Culex (Culex) theileri* Theobald, 1903.

M.C. III, p. 187 (♂ & ♀) TYPE-LOC. Pretoria, Transvaal, S. Africa (*Theiler*) TYPE ♂ & ♀ in Brit. Mus.

Culex pettigrewi Theobald, 1910, Rec. Ind. Mus. IV, p. 15 (♀)

TYPE-LOC. Ukhrul, Manipur, 8,400', Assam, viii 1908 (*Pettigrew*)

TYPE ♀ in Ind. Mus.

Culex impuliformis Edwards (nec Theobald), 1912, Bull. Ent. Res. III, p. 31


ADULT * —A rather large brown species with conspicuously striped femora and tibiae, abdominal tergites with triangular yellowish basal bands and usually with two pale spots or scattered pale scaling in middle of each tergite in addition. Wing about 5 mm.

♀ —*Head* vertex and nape covered with narrow pale brown scales and numerous upright scales. Palpi black, with white scales apically. Proboscis brown, with median pale area beneath and at sides, but not forming a definite pale band. *Thorax*: mesonotum covered with narrow brown scales, a pair of submedian dark bare lines running back from the front, narrowing posteriorly, some narrow brown scales on upper part of *ppn* and on scutellum. Pleurae with a line of broad pale scales along posterior border of sternopleura, and one large or two small patches on mesopimeron. *Wings* dark scaled, c-v 4-5 only slightly nearer base of wing than 3-4. *Legs* fore femur yellowish dorsally for whole length, anterior surface with a median longitudinal

* Edwards 1911, p. 262, 1921 c, p. 330 (*impuliformis*); 1926 b, p. 637. Barraud 1924 g, p. 1266, Kirkpatrick 1925, p. 115 (*impuliformis*); Bedford 1928, p. 971, Martini 1931, p. 384.

dark brown stripe, a narrower white line ventrally, posterior surface pale, except for a narrow ventral dark brown line, mid-femur, seen from front, brownish-black, with a median longitudinal pale stripe, posteriorly almost entirely pale, hind femur dark dorsally from base nearly to knee, tip pale, basal $\frac{1}{3}$ or more of anterior surface otherwise pale, a brown line ventrally along apical $\frac{1}{3}$ or more. All tibiae with pale line anteriorly, entirely pale posteriorly. Tarsi without pale rings, but segment 1 usually faintly striped with pale scaling.

Fig 98



♂ hypopygial structures of *Culex* (proctiger and phallosome, lobe of coxite). a, *hutchinsoni*; b, *nigricornis*; c, *theileri*; d, *vagens*

Abdomen: tergites brownish-black, with triangular yellowish basal bands, wider in middle than at sides, usually a pair of pale spots or scattered pale scaling, or both, in middle of each tergite in addition.

♂.—Palpi longer than proboscis by only about length of last segment; long segment with narrow pale ring on basal $\frac{1}{3}$, and a patch of pale scales on outer side near apex, not forming a definite ring; tip of long segment and last two segments with outstanding dark hairs; penultimate

segment with some pale scales on outer side near base, last two segments with a line of white scales beneath. Proboscis brown, paler on apical $\frac{1}{2}$. Other markings as in ♀ *Hypopygium* (fig 98, c) appendage of subapical lobe of coxite, *lp*, and paraproct as shown.

LARVA * (fig. 99, a-f) — *Antenna* as shown in fig 99, a, dark brown on apical $\frac{1}{2}$ and at base, remainder pale. *Mentum* as in fig 99, b. *Comb* of about 30 teeth in a triangular patch, individual teeth comparatively long, with a terminal point and lateral fringe. *Siphon* widest at base and slightly tapering to tip, $4\frac{1}{2}$ – $5\frac{1}{2}$ times length of diameter at base, usually uniformly pale, but with dark ring at base, and occasionally basal $\frac{1}{2}$ is darkened. Usually 5 pairs of rather stout branched hairs along posterior border of tube, and one small pair of lateral 2-branched hairs. *Pecten* of 7–10 large curved teeth, those furthest from base more widely spaced than others, 3–4 lateral denticles at base of each tooth. *Anal segment* with its hair and papillæ as shown in fig 99, c.

HABITAT — Large ground-pools, stream-pools, and marshes.

DISTRIBUTION — Common in WESTERN HIMALAYAS, KASHMIR, PUNJAB, NORTH-WEST FRONTIER, and BALUCHISTAN. Less common to the east, but occurs in ASSAM hills and BURMA (Northern Shan States, *Senior-White*). It has been recorded from 9,500', in HINDU KUSH mountains (Wall).

It has a wide distribution in the Mediterranean region, extending to ATLANTIC ISLANDS westwards, and through MESOPOTAMIA to PERSIA eastwards. Also known from EAST and SOUTH AFRICA.

240. *Culex (Culex) vagans* Wiedemann, 1828.

Aussereurop. Zweifl. Ins. 1, p. 545 (♀). TYPE-LOC · Foochow, China. TYPE ♀ in Vienna Museum may be the type.

Culex tipuliformis Theobald (nec Edwards), 1901, M.C. II, p. 325.

(♀). TYPE-LOC. Bakloh, Western Himalayas (*Lindesay*).

TYPE ♀ in Brit. Mus.

Culex virgatipes Edwards, 1914, Bull. Ent. Res. v, p. 126 (♂ & ♀).

TYPE-LOC Hong Kong (*Macfarlane*). TYPE ♂ in Brit. Mus.

ADULT † — Resembles *thesleri* fairly closely, but usually rather smaller (wing 4–4.5 mm), hind femur entirely pale ventrally from base to knee on outer side; abdominal tergites

* Bedford 1919, p. 741 (*tipuliformis*), Edwards 1921 c, p. 340 (*tipuliformis*), 1926 b, p. 637; Ségué 1924, p. 40 (*tipuliformis*), Barraud 1924 m, p. 429, Kirkpatrick 1925, p. 117 (*tipuliformis*); de Meillon 1928, p. 321, Martini 1931, p. 385.


† Edwards 1921 c, p. 341 (*virgatipes*); 1922 c, p. 278 (*virgatipes*), 1922 d, p. 473, 1926 b, p. 640; Barraud 1924 g, p. 1269 (*tipuliformis*), Martini 1931, p. 391.

dark except for moderately wide creamy basal bands, which are not produced in middle, mesonotal scales rather more reddish-brown, c-v. 4-5 more separated from 3-4 than in *theileri*. Palpi of ♂ longer than proboscis by about $1\frac{1}{2}$ times length of terminal segment

The winter form of this species occurring in Western Himalayas is often darker both in adult and larval stages than is usual.

♂—*Hypopygium* (fig 98,d) *lp* differing in shape from that of *theileri*.

Fig 99.

Larval structures of *Culex*. a-f, *theileri*, g-l, *vagans*.

LARVA * (fig. 99, g-l).—Distinct from *theileri* in a number of characters (form of mentum, pecten-teeth, etc), but very similar to *fatigans*, and apparently indistinguishable from larva of *pipiens* (a species not yet found in India). From larva of *fatigans* it differs in the following details:—*Siphon* rather longer, about 5 times length of diameter at base (instead of about 4 times), number of pecten-teeth 12-15 (in *fatigans* usually 10 or 11, but varies between 8 and 12); *lh* of 2 fairly long fine branches (in *fatigans* usually single and rather shorter), comb-teeth rather wider and more completely fringed than in *fatigans*; mentum of slightly different

* Barraud 1924 m, p. 480 (*tipuliformis*).

form, as indicated in figures. Hairs on siphon of fewer branches.

HABITAT—Ground-pools, marshes, pools in beds of streams.

DISTRIBUTION—Common in WESTERN HIMALAYAS, KASHMIR, and NORTH-WEST FRONTIER, and extending to PUNJAB and DELHI in the cold weather. It appears to be less common in EASTERN HIMALAYAS, but occurs in SOUTHERN and EASTERN CHINA and SIBERIA (Vladivostok).

241 *Culex (Culex) univittatus* Theobald, 1901

M C II, p. 29 (♂ & ♀) TYPE-LOC Durban, East Africa, II, 1899
(Christophers) TYPE ♂ & ♀ in Brit Mus

Culex peregrinus Theobald, 1903, M C III, p. 100 (♂ & ♀) TYPE-LOC Sidon, Palestine (Cropper) TYPE ♂ & ♀ in Brit Mus

ADULT*—A small pale brown mosquito resembling *vishnui* in size and general coloration, but distinguished by absence of pale bands on proboscis and tarsi and by white, not ochreous, basal bands on abdomen. Wing about 2.7 (♂)—3.5 mm. (♀).

♀—*Head*: vertex with narrow pale brown scales, upright pale scales on nape, a patch of broad white scales low down each side. Palpi dark brown, with white scales at tips. Proboscis brown, without a pale band, but broadly and indistinctly pale in middle, beneath, and at sides. *Thorax*: mesonotum, scutellum, *apn*, and *ppn* covered with pale brown or pale golden narrow scales. Pleuræ with patches of broad white scales, [including a small but conspicuous postspiracular patch immediately behind prothoracic spiracle], one long lower mesepimeral bristle. *Wings* dark scaled. *Legs*: brown, but extensively pale posteriorly and beneath, femora with small pale knee-spots, hind femur usually entirely pale, except for brown scaling along dorsal border towards apex (in some Indian specimens, and in African specimens, the brown scaling is more extensive), mid- and hind tibiae with pale stripe on outer side, hind tibia with a rather conspicuous white marking at apex. Tarsi without pale rings at joints. *Abdomen*: tergites dark brown, with narrow basal white bands (not ochreous as in many other species) and wider lateral patches not usually visible in dorsal view. Sternites pale.


♂.—Palpi longer than proboscis by rather more than length of last segment, long segment light in colour and without any distinct pale rings, tip of that segment and last two segments with outstanding dark hairs, last two segments

* Edwards 1911, p. 262, 1912, p. 32; 1914 b, p. 67; 1921 c, p. 342 (*perexiguus*), 1922 b, p. 85, 1926 b, p. 642; Barraud 1924 g, p. 1263 (*perexiguus*), Kirkpatrick 1925, p. 125 (*perexiguus*); De Meillon 1928, p. 321.

with a line of white scales on underside *Hypopygium* (fig 100, a). ventral cornu of *lp* simple, but with a small spine-like process on ventral border, lateral process narrow, fairly long and pointed, median process large and pointed. Paraproct with long basal arm

LARVA * (fig 101, b) very similar to *tritæniorhynchus*, *barraudi*, and *fuscoccephalus*. Differs from the first-named

Fig. 100



♂ hypopygial structures of *Culex*. a, *univittatus*, b, *fatigans*, c, *fuscoccephalus*, d, *fuscitarsis*. Lettering as on p. 4.

as follows.—Antennal tuft at about $\frac{2}{3}$ from base, instead of at about $\frac{1}{3}$; 5 pairs of branched hairs on siphon, of which two are more or less lateral, instead of 6 pairs of which one is lateral; *lh* with shorter branches.

Differs from *barraudi* in having 5 pairs of branched hairs on

* Edwards 1921 c, p. 343; 1926 b, p. 648; Barraud 1924 m, p. 431; Séguy 1924, p. 38; Kirkpatrick 1925, p. 128; Galliard 1927, p. 99 (all as *peregrinus* except Edwards 1926). The larva figured as *O. univittatus* by Ingram and Macfie (1919, p. 68) is *Aedes (Aedimorphus) ochraceus* Theo.

siphon, of which two are lateral, instead of 4 pairs of which one is lateral, teeth of mentum all about same size and regular

Differs from *fuscocephalus* in number and arrangement of branched hairs on siphon, in length of siphon, which is about 7 times length of diameter at base (instead of $4\frac{1}{2}$ -5 in *fuscocephalus*), and in form of *lh*

HABITAT—Marshy pools, borrow-pits, stagnant drains and canals, shallow wells, less frequently in domestic collections of water and rice-fields

DISTRIBUTION—NORTH-WEST FRONTIER Shahgai*, Peshawar dist., vii 1929 BALUCHISTAN Quetta* (*Davys*) SIND Larkana* (*Barly*) PUNJAB. Amritsar and Mahdopur, iv 1921 (*Barraud*), Lahore*, ix 1923 (*Sinton*) UNITED PROVINCES · Saharanpur*, 1926 (*Sinton*)

Also known from MEDITERRANEAN REGION, AFRICA, and MADAGASCAR

242. *Culex (Culex) fatigans* Wiedemann, 1828.

Ausereurop Zweifl Ins 1, p 10 (♂ & ♀) TYPE-LOC. East Indies
TYPE Vienna Mus (?)

Culex cingulatus Doleschall, 1856, Nat Tijd v Ned-Ind x, p 405
TYPE-LOC Java TYPE non-existent.

Culex doleschallii Giles, 1900, Handbook, p 338, nom nov. for
C cingulatus Dol

Culex quasipipiens Theobald, 1901, M C ii, p 136 (♀) TYPE-
LOC Sambalpur, Orissa (*Murphy*) TYPE ♀ in Brit Mus.

Culex fouchouensis Theobald, 1901, M C ii, p. 137 (♂ & ♀) TYPE-
LOC : Fou Chow, China (*Rennie*) TYPE: ♂ & ♀ in Brit Mus

Culex reesi Theobald, 1901, M C ii, p. 145 (♂ & ♀) TYPE-LOC .
Hong Kong (*Rees*) TYPE. probably non-existent.

Culex sericeus Theobald, 1901, M C ii, p. 147 (♀) TYPE-LOC
Hong Kong (*Rees*) TYPE probably non-existent

Culex albolineatus Giles, 1901, Journ Bomb. Nat. Hist Soc xiii,
p 609 (♀) TYPE-LOC. Shahjahanpur, United Provinces, India,
x 1900 (*Giles*). TYPE ♀ in Brit Mus

Culex christophers Theobald, 1907, M C iv, p 453 (♂ & ♀). TYPE-
LOC. India (*Christophers*) TYPE ♂ & ♀ in Brit. Mus.

Culex minor Theobald, 1908, Rev Ind Mus ii, p 298 (♂ & ♀)
TYPE-LOC Lushai Hills, Assam (*Macleod*) (♂), Sylhet, Assam
(*Hall*) (♀). TYPE ♂ & ♀ in in Ind. Mus (examined by Author).


ADULT †—Usually recognizable by dark tarsi and (in ♀) somewhat rounded abdominal bands Wing about 3.5 (♂)-4.5 mm (♀).

♀—*Head* vertex and nape covered with narrow golden-brown scales and scattered upright scales, latter more numerous on nape, a patch of broad pale scales low down at each side. Palpi brown, about $\frac{1}{2}$ length of proboscis, latter dark brown,

† Theobald 1901 b, p 151, Edwards 1913 a, p 55; 1921 c, p. 345, 1924 b, p. 395, 1926 a, p 141, 1926 b, p. 645, Barraud 1924 g, p 1264, Brug 1924 b, p 9, Borel 1926, p 98, Martini 1931, p 364.

usually with an indistinct paler area about the middle, especially on underside and sides; labella pale brown. *Thorax*: mesonotum, scutellum, *apn*, and *ppn* covered with narrow golden-brown scales. From front of mesonotum two submedian dark bare lines run back a short distance, narrowing posteriorly. Pleurae brown, with two patches of broad pale scales on mesepimeron and two on sternopleura, [none on postspiracular area.] No definite dark areas. *Wings*: dark scaled, outstanding scales narrow. *Legs*: brown, dark brown,

Fig. 101



Larval structures of *Oulex*: a, *fatigans*; b, *univittatus*; c, *fuscicephalus*.

or nearly black when viewed from front or from above, paler, especially femora, when seen from behind or from beneath; hind femur dark brown on dorsal border on outer side, otherwise pale; femora with very small knee-spots, hardly visible, except on hind pair. Mid- and hind tibiae usually marked with a narrow yellowish ring at apex. Tarsi entirely dark. *Abdomen*: tergite I almost entirely covered with long yellow hairs, a patch of dark scales on apical border in middle; II-VII dark brown or black, with transverse

ochreous (not white) basal bands, a little wider in middle than laterally, lateral borders of VI and VII, and usually whole of VIII, pale scaled. Sternites entirely covered with pale scaling.

♂—Palpi longer than proboscis by rather less than length of terminal segment, long segment with narrow pale ring on basal $\frac{1}{2}$, and usually with a rather wide indefinite pale area near apex, apical $\frac{1}{3}$ of long segment and last two segments with outstanding dark hairs, a line of white scales on under surface of penultimate segment, a spot of white scales at base of terminal segment on underside. Pale abdominal bands usually wider than in ♀.

Variation—Specimens from Assam Hills (Shillong) have darker mesonotal scaling than usual and pleurae are very dark greenish-brown or black. In some Western Himalayan examples one or more basal pale bands on abdomen are triangular, and dark parts of tergites may be greyish-brown, with or without a pair of indistinct lighter spots. Giles's type of *C. albomeatus* has somewhat this type of colouring.

♂—*Hypopygium* (fig 100, b): *lp* very characteristic. Parapectus usually with a small lateral arm.

LARVA* (figs 101, a, & 102)—Head, antenna, mentum, siphon, and anal segment as shown in fig. 101, a. [Thoracic chaetotaxy† as shown in fig 102.] Comb of 35–40 small fringed teeth in a triangular patch. Siphon usually about 4 times length of diameter at base. Distinguished from *vagans*, which it closely resembles, as mentioned on p. 417.

HABITAT—Domestic collections of water and in such places as flooded open cement drains, flooded latrines, overflow water from houses, kitchens, etc. Also in ground-pools, ditches, and shallow wells. Rarely in tree-holes or bamboos;

* Christophers 1906, p. 7, Barraud 1924 m, p. 430, Berol 1926, p. 100, Senior-White 1927, p. 71, Buxton & Hopkins 1927, p. 83, Martini 1931, p. 365. *Pupa*: Christophers 1906, p. 7; Ingram & Macfie 1917 a, p. 84, Buxton & Hopkins 1927, p. 80. *Egg*: Buxton & Hopkins 1925, p. 295.


† [The thoracic chaetotaxy of *C. fatigans* is illustrated for comparison with *Tripteroides* (fig. 9) and *Aedes* (fig. 39). Several other species of *Culex* examined are closely similar, and the arrangement shown may be taken as typical of the genus. The notation adopted is the same as for the other genera. Noteworthy points are—*Prothorax*: the pair of plates surrounding bases of hairs 1–3 are more conspicuous than in *Aedes* or *Tripteroides*, and are much closer together than in *Aedes*; hair 8 is as well developed as 7, and its base is just visible from above; 9–12 are latero-ventral in origin, hair 14 absent. *Mesothorax*: arrangement much as in the other two genera; hairs 6–7 arising from a common plate, 8 large, 13 and 14 very small. *Metathorax*: arrangement much as in the other two genera; hairs 1–6 all small; 7 large, 8 very small, these two (as in *Aedes*) resembling in size and position hairs 8 and 14 of mesothorax, 11 apparently absent, 13 fairly well developed.]

not in jungle-pools, stream-pools, etc., if far from human habitations

DISTRIBUTION—One of the most abundant of Indian Culicini. Found in all parts of the Indian region, and is largely a domestic mosquito. It occurs up to 5,000' or more in the hills. In parts of the country it is most probably concerned in the transmission of filariasis.

It is common in the tropics and subtropics of both New and Old Worlds

Fig 102



Chætotaxy of larval thorax of *Culex fatigans*: left, dorsal
right, ventral.

243 *Culex (Culex) hutchinsoni* Barraud, 1924.

Ind. Journ. Med. Res. xi, p. 1261 (♂ & ♀) TYPE-LOC.: Nongpoh, Khasi Hills dist., Assam, vii. 1922 (*Barraud*). TYPE: ♂ & ♀ in Brit. Mus

ADULT.—A very small dark brown mosquito having some resemblance to *tritemorhynchus*, but proboscis not definitely banded, tarsi entirely dark, and ♂ hypopygium of very different form. Wing 2.6–3 mm.

♀—**Head**: very similar to that of *fuscocephalus*; a mixture of golden-brown and pale narrow-curved scales on vertex, with scattered upright scales, more numerous upright scales on nape; a patch of broad white scales low down each side. Palpi brown, with a few pale scales apically. Proboscis brown, paler ventrally, except at extreme base and tip, pale scales at sides, and a sprinkling of pale scales on middle $\frac{2}{3}$ of upper surface, not forming a definite band. **Thorax**: mesonotum, scutellum, *apn*, and *ppn* clothed with narrow

brown scales of a deep golden tint. Pleuræ apparently entirely dark, with one or two patches of white scales, one lower mesepimeral bristle. *Wings*: dark scaled, outstanding scales long and narrow. *Legs*: dark brown, with a pale sheen when seen from certain angles, fore and mid-pairs paler posteriorly, especially femora. Hind femur pale, except along dorsal border. Hind tibia with narrow pale ring at apex. Tarsi entirely dark. *Abdomen*: brownish-black, with narrow basal pale bands and wider lateral patches. Venter pale yellowish.

♂—Palpi longer than proboscis by a little more than length of terminal segment, deep brown except for a rather broad pale-scaled area on apical $\frac{1}{2}$ of long segment, last two segments and tip of long segment with outstanding dark brown hairs, a few light scales on ventral surface of penultimate segment, not forming a conspicuous line. Proboscis dark brown, with paler scaling forming an indefinite and inconspicuous band nearer apex than base. Otherwise ♂ resembles ♀, except for plumose antennæ. *Hypopygium*: ventral cornu of *lp* divided into three pointed projections, lateral process moderate, median process long and pointed, proctiger with long basal arm.

LARVA.—Unknown.

HABITAT.—Road-side pools.

DISTRIBUTION.—Known only from type-locality.

244. *Culex (Culex) fuseocephalus* Theobald, 1907.

M. C. iv, p. 420 (♀). TYPE-LOC: Peradeniya, Ceylon (*Green*).
TYPE ♀ in Brit. Mus.

Culex uniformis Leicester, 1908, *Cul. Malaya*, p. 159 (♀). TYPE-LOC: Batu Gajah, Malay Penin. (*Leicester*). TYPE ♀ in Brit. Mus.

Culex minimus Leicester, 1908, *Cul. Malaya*, p. 160 (♂ & ♀). TYPE-LOC: Kuala Lumpur, Malay Penin. (*Leicester*). TYPE: non-existent.

Culex taytayensis Banks, 1909, *Phil. Journ. Sci.* iv, p. 545 (♂ & ♀). TYPE-LOC: Rizal, Taytay, Philippine Is. TYPE: ♂ & ♀ no 11549 in Bureau of Sci., Manila.

Culex luteola Theobald, 1910, M. C. v, p. 378 (♀). TYPE-LOC: Peradeniya, Ceylon (*Green*). TYPE ♀ in Brit. Mus.

Culex inelegrans Dyar, 1920, *Insec. Insc. Mens.* viii, p. 179 (♂ & ♀). TYPE-LOC: Los Banos, Philippine Is., vi 1915 (*Baker*). TYPE ♂ no 23719 in U. S. Nat. Mus.

ADULT*.—Recognized by its general dull brown colour and unbanded abdomen, and by presence of two horizontal dark bands on pleuræ divided by a white-scaled band. Wing about 3 mm.

* Edwards 1913 b, p. 234 (syn.); Barraud 1924 g, p. 1270; Borel 1926, p. 101.

♀—*Head* vertex covered with a mixture of brown and white narrow scales and upright scales, latter more numerous on nape, a patch of broad white scales at each side. Head when viewed from front appears to be almost entirely covered with pale scales on a dark ground, from above vertex appears dark, with a conspicuous white area on each side. Proboscis brown, pale beneath, especially on apical $\frac{1}{2}$. Palpi brown. *Thorax* mesonotum uniformly clothed over largest part with dark brown scales, paler scales in front of scutellum, and some very light-coloured scales on front margin behind head. Scutellum, *apn*, and *ppn* with narrow pale brown scales. A broad line of white scales from *apn* across pleuræ to middle of mesepimeron; integument both above and below this line brownish-black and without scales, except for a patch of white scales around lower sternopleural bristles, one or two lower mesepimeral bristles. *Legs*: dark brown when seen from front or from above, paler beneath; hind femur mainly pale, with brown scales along dorsal border towards apex. Hind tibia with more or less defined pale stripe on outer side. Tarsi brown. *Abdomen*. dusky brown dorsally, posterior borders of tergites with pale hairs, giving a somewhat banded appearance; lateral margins of tergites with pale scales extending from base nearly to hind margin, not usually visible dorsally. In some specimens there are a few pale scales at base of several segments in middle, not forming definite transverse bands. Sternites entirely pale scaled.

♂—Palpi longer than proboscis by rather more than length of last segment; long segment with very narrow pale ring on basal $\frac{1}{2}$, and some pale scaling along outer side towards apex, last two segments and apical $\frac{1}{2}$ of long segment with outstanding dark hairs; penultimate segment with a line of white scales beneath. *Hypopygium* (fig. 100, c): *lp* with a number of teeth and moderately long median process. Paraproot with long basal arm, and an extension of the plate just above this.

LARVA * (fig. 101, c).—Head and antenna, mentum, siphon, etc., as shown in figures. Very similar to *barraudi*, *tritaniorhynchus*, and *univittatus*, differing from the first-named in shape of mentum, in having antennal tuft at $\frac{2}{3}$ of length of shaft from base (instead of $\frac{1}{3}$ – $\frac{2}{3}$), and in having usually 10 or 11 comparatively large pecten-teeth. Differs from the other two species mentioned above chiefly in number and arrangement of branched hairs on siphon, size of pecten-

* Barraud 1924 m, p. 432; Borel 1926, p. 103; Senior-White 1927, p. 71.

teeth, and in form of lateral hair of anal segment Siphon usually about 5 times length of diameter at base

HABITAT —Ground-pools, rice-fields, etc

DISTRIBUTION —Widely distributed in India from north-west PUNJAB (Jhelum) to ASSAM, BURMA, and ANDAMAN Is, and through PENINSULAR INDIA to CEYLON

Known also from MALAY PENINSULA, SIAM, COCHIN CHINA, and DUTCH EAST INDIES as far south-east as TIMOR, PHILIPPINE Is, and SOUTH CHINA

245 *Culex (Culex) fuscitarsis* Barraud, 1924.

Ind Journ Med Res xi, p 1272 (♂ & ♀). TYPE-LOC. Pachmarhi, Central Provinces, India, vii 1915 (*Dowson*) TYPE ♂ & ♀ in Brit Mus

ADULT —A small dusky-brown mosquito resembling *fuscicephalus*, and possibly only a variety of that species, differing in having pale ochreous basal bands on the abdomen (narrow in ♀, wider in ♂) There are also slight differences in the structure of ♂ hypopygium The head also appears darker on vertex and the lateral white areas are less evident Wing about 3 mm

♂ —*Hypopygium* (fig 100, d) paraproct without the pronounced expansion above basal arm which is present in *fuscicephalus*, otherwise structure is very similar

LARVA —Unknown

HABITAT —Ground-pools, river-pools, and swamps

DISTRIBUTION —CENTRAL PROVINCES. Pachmarhi*, typolocality, as given above, Padreganj*, Seoni dist., and Kukra Kapa*, Satpuras, vii. 1931 (*Sensor-White*). CENTRAL INDIA. Ghunghtui*, Rewa State, vii 1931 (*Sensor-White*). ORISSA. Ranchi*, viii. 1922 (*Fletcher*); Noamundi*, Singhbhum dist., viii 1929 (*Sensor-White*). DELHI PROVINCE: Delhi*, between old city and Jumna River, viii 1927 (*Sensor-White*). BOMBAY PRES. Poona* and Lonavla*, xi. 1928 (Dist -Surg. G I.P. Rlwy.) MADRAS PRES. NORTH Ambadola*, Vizagapatam Agency (*Sinton*)

Not known from elsewhere.

APPENDIX.

By F. W. EDWARDS, M.A., Sc D

I. Characters and Affinities of the Culicidæ	427
II. The Indian Dixina and Chaoborinæ	429
III. Further Descriptions of Indian Culicini	441

I. CHARACTERS AND AFFINITIES OF THE CULICIDÆ

The family Culicidæ is a well-circumscribed group of Nematocerous Diptera which may be defined as follows.—

Small, or smallish, slenderly-built flies, ranging from 3–10 mm. in length of body. Antennæ composed of 15–16 segments*, the first vestigial, forming a narrow ring, the second segment, or torus, more or less conspicuously enlarged. Ocelli absent. Thorax with pronotum divided in middle, mesonotum without any suture dividing the præscutum from the scutum, postnotum, or mesophragma, well developed, but metanotum reduced to a narrow strip. Abdomen with eight distinct segments, apart from the genital and anal segments. Legs long and slender; tibiæ without distinct spurs†. Wings with a uniform type of venation‡: vein Sc long and ending in costa; vein 2 forked; 3 simple; 4 forked, 5 forked, 7 absent or represented by an indefinite fold in the membrane; cross-veins 3–4 and 4–5 both present.

Pupa aquatic and free-swimming. Prothoracic respiratory horn present and provided with a single opening. Leg-sheaths coiled, so that they do not project beyond the short wing-cases. No strong spines on thorax and no rows of spines on abdomen.

Larva aquatic, with well-developed head-capsule and long antennæ. Only one pair of spiracles present, dorsally placed on abdominal segment IX§.

* Except in one species of *Chaoborus*, in which the antennæ are reduced.

† Very small tibial spurs are present in some Chaoborinæ, otherwise spurs are absent.

‡ The only genus to which this definition will not apply in its entirety is the New Zealand *Obolopoda*, which has a single long vein.

§ Spiracles absent or closed in some *Chaoborus*.

Some of the characters noted in the above definition are to be found equally in other groups, but a reference to the wing-venation will at once dispel any doubt as to whether a given insect belongs to the family Culicidæ or not, all members of this family, and none of any other, have veins 2 and 4 forked (2-branched), while vein 3 is simple. Certain small Tipulidæ (genera *Geranomyia*, *Toxorhina*, and others) are provided with a long proboscis, and are sometimes mistaken for mosquitoes, but apart from the fact that the proboscis is very differently constructed and incapable of use as a biting organ, these Tipulidæ may be recognized by their more complex venation, vein 7 being present. The majority of the Chironomidæ resemble the mosquitoes in having the antennæ plumose in the male sex, but have the venation less complex, vein 4 being simple and cross-vein 4-5 often absent. The Ceratopogonidæ and Psychodidæ are less likely to be confused with the Culicidæ owing to their smaller size. moreover, in these also the wing-venation is quite different.

The pupæ of some Chironomidæ (notably those of *Tanyptus* and related genera) are very similar to those of mosquitoes, having the same general form and habits, but differ in the structure of the respiratory horn, in having the paddles united basally (as is also the case in *Dixa* and *Corethrella*), and in abdominal chaetotaxy, segments I-VI being largely bare, VII and VIII with a row of stiff bristles along each side, and each paddle with two similar lateral hairs.

The only other aquatic Dipterous larvæ which might conceivably be mistaken for Culicidæ are those of certain Psychodidæ which possess a somewhat similar respiratory siphon, but in these the form of the body and the cuticular armature is quite different, apart from more fundamental differences in head-structure. The larvæ of Chironomidæ differ very obviously in the absence of spiracles and presence of prothoracic and terminal abdominal prolegs, those of the comparatively few aquatic Tipulidæ in the incomplete head-capsule, bare body, and terminal instead of dorsal spiracles.

From a consideration of all the characters of all stages of their life-history it now appears certain that the nearest relatives of the Culicidæ are the Chironomidæ, and that the Ceratopogonidæ and Simuliidæ, though rather less closely connected, belong to the same stock; the other families of Nematocera (Tipulidæ, Psychodidæ, Mycetophilidæ, etc.) have comparatively little in common with the Culicid group. Among the adult characters common to the four families of this group may be mentioned: (1) the reduction of the first antennal segment—not seen in other families, (2) absence of ocelli, (3) resemblances in structure of the thoracic pleuræ,

e. g., in the presence of a large membranous area behind the prothoracic spiracle. The difference in venation between the Culicidæ and the other three families, though great, is, perhaps, not so fundamental as might be supposed, as it is not difficult to show how the venation of the Chironomidæ might have been derived from that of the Culicidæ by a process of reduction and slight modification.

Another point in which the four families Culicidæ, Chironomidæ, Ceratopogonidæ, and Simuliidæ are alike is in having the legs of the pupa coiled and tucked under the wing-cases. In all other families of Nematocera the tarsal sheaths are straight and project well beyond the wing-cases. It would seem that this modification of the pupa was in the first instance developed to allow freedom of movement of the abdomen so that the pupa might swim actively, the fact that the modification has been retained in the inactive pupæ of Simuliidæ may, perhaps, be claimed as a further example of the irreversibility of evolution.

II.—THE INDIAN DIXINÆ AND CHAOBORINÆ.

In order to complete the account of the Indian Culicidæ it seems desirable to include a brief description of the subfamilies Dixinæ and Chaoborinæ. The members of these subfamilies are not blood-suckers, and have no economic importance—except that flies of the genus *Chaoborus* are used by some African natives as food, nevertheless, they are of great biological interest. As in the case of the mosquitoes (Culicidæ), all members of these groups are aquatic in their early stages. The larvæ of *Dixa* have some resemblances in structure and habits to those of *Anopheles*, and the two genera were at one time confused.

Subfamily DIXINÆ.

The genus *Dixa* was long regarded as constituting a distinct family of Nematocera, and was so treated by Brunetti in his volume on Diptera Nematocera in the 'Fauna of British India' series. It is, however, undoubtedly related to the Culicidæ, and is best included in the family Culicidæ, forming a separate subfamily. Only two genera of the subfamily are known, the second (*Neodixa*) may be disregarded here, as it is represented by a single known species from New Zealand.

The chief subfamily characters are the following:—

ADULT.—Eyes widely separated and rounded. Mouth-parts short, not adapted for biting; palpi (in both sexes)

incurved. Antennal flagellum of 14 segments (not 13 as in Culicinæ and Chaoborinæ), never plumose in ♂. *Wings* with venation similar in type to that of the other two subfamilies, but subcosta shorter, ending above or before base of sector, sector oblique, stem of upper fork strongly curved upwards. No scales, but only short hairs, on veins or in fringe. *Legs* without scales and not conspicuously hairy. *Hypopygium* of ♂ very varied in structure in the different species, but with coxite short and broad, style without a terminal spine, proctiger with distinct chitinisations (often elaborate), including a pair of hairy cerci or paraprocts, phallosome various, usually complicated.

LARVA—Differs from that of the other two subfamilies in the much more distinct segmentation of the thorax, which is little, if any, broader than the abdomen. Mouth-bristles present. Antennæ not prehensile, the spines at tip short and not movable. Fore gut communicating with mid-gut. Spiracular apparatus large, with functional spiracles but no siphon. Abdomen with a pair of ventral prolegs, surmounted by hooks, on each of the first two segments, and with pairs of ventral combs on segments VI and VII.

Genus *DIXA* Meigen, 1818

Syst. Besch. 1, p. 218. Genotype, *D. maculata*, Meig.

Brunetti recorded five Indian species of this genus*, one more has since been described by Senior-White, and several others are described below. All are rather small flies, with a wing-length of about 3-5 mm.

The genus is now divided into five subgenera, three of which do not occur in Europe or Asia, and, therefore, do not concern us here. The described Indian species all appear to belong to the subgenus *Dixa*, but one species of the subgenus *Paradixa* is represented in the British Museum by a specimen from Kashmir, described below as new. It is probable that many Indian species of both subgenera await discovery.

Key to Indian Species of Dixa

- | | | |
|---|--|--------------------------------------|
| 1 | Wings with elaborate markings, including two large brown areas in upper basal cell | [p. 431
<i>maculipennis</i> Brun, |
| | Wings less extensively marked | 2. |
| 2 | A brown or greyish spot (sometimes faint) in upper basal cell | 3. |
| | No trace of dark spot in upper basal cell | 6. |

* Brunetti brought his types to London in 1921, and I made some notes on them before they were returned to Calcutta, but some details of their structure remain undescribed.

- | | | |
|----|--|-----------------------------------|
| 3 | Wing-tip darkened, head yellow | <i>ochroleucata</i> Brun., |
| | Wing-tip not darkened, head dark | 4 [p 434] |
| 4 | Grey markings present between veins 2 and 4 towards tip of wing | 5 |
| | No such markings | sp. n.?, pp 433-4 |
| 5 | A small grey spot towards end of lower basal cell | <i>montana</i> Brun., p 433 |
| | No such spot | <i>trinitata</i> , sp. n., p 432 |
| 6 | Vertex and tori yellow, proboscis rather long | [p 434] |
| | Vertex and tori blackish, proboscis normal | <i>cylauca</i> Sonner-White, |
| | | 7 |
| 7 | Thorax almost uniformly blackish, a diffuse grey cloud over cross-veins | 8 |
| | Thorax with yellowish ground-colour, a small brown spot over cross-veins | 9 [p 435] |
| 8 | Upper fork-cell shorter than its stem | <i>chiropterus</i> , sp. n., |
| | Upper fork-cell longer than its stem | <i>platystyla</i> , sp. n., p 436 |
| 9 | First flagellar segment of antenna shorter, slightly thickened | 10 [p. 437. |
| | First flagellar segment longer, cylindrical | <i>kashmirensis</i> , sp. n., |
| 10 | Spot over cross-veins reaching well into marginal cell | <i>bistrata</i> Brun., p 435 |
| | Spot over cross-veins scarcely extending into marginal cell | <i>bifasciata</i> Brun., p 435 |

Subgenus *DIXA* Meigen, s. str

Genotype, *Dixa maculata* Meigen (Europe)

ADULT—Antennæ with first segment of flagellum slightly thickened towards middle, usually not more than about 4 times as long as broad. Hypopygium of ♂ with lobe at apex of coxite short or absent, style usually short and simple.

LARVA.—Antenna without hairs on inner side. Abdomen with a dorsal crown or rosette of stiff hairs on each of segments II-VII or III-VII. Apex of large lateral plate of spiracular apparatus without spur or tooth. Caudal projection rarely reaching beyond posterior border of spiracular apparatus.

So far as known, the larvæ of this subgenus live in running water.

1 *Dixa (Dixa) maculipennis* Brunetti, 1911.

Rec. Ind. Mus. iv, p. 286. Type-loc. Darjiling (*Brunetti*) and Matiana, Simla dist. (*Annandale*). Types: ♂ & ♀ in Ind. Mus.

ADULT.—Readily distinguished from other Indian species* by the conspicuously marmorated wings, the markings including two large dark brown spots completely crossing the upper basal cell and three crossing the marginal cell. Structural

* The Chinese *D. guttipennis* Thomson is very similar, but has the markings in the upper half of the wing relatively smaller and those in the lower half relatively larger.


characters as in *D. montana* and related species first flagellar segment moderately long but slightly thickened, subcostal vein ending slightly before base of sector, cerci of ♀ long and pointed ♂ not available for description

DISTRIBUTION—Known from type-localities and from Ritani, Muktesar, 13 v 1923 (*Fletcher*)

2 *Dixa* (*Dixa*) *trinotata*, sp. n

ADULT ♂—*Head* dark above, clypeus pale yellowish, quite short Antenna with scape dark brown (flagellum missing in type) Palpi dark *Thorax* mesonotum yellowish,

Fig 103



Wing of *Dixa trinotata*, sp. n. (a), coxite, with lobes, and style of ♂ hypopygium of *D. trinotata*, sp. n. (b), *D. christophersi*, sp. n. (c), and *D. platystyla*, sp. n. (d)

with three separate dark brown stripes, median stripe not distinctly divided, no darkening on shoulders in front of lateral stripes, scutellum dark at sides, postnotum dark, pleuræ mainly dark, pale on sutures, pruinosity very slight. No sternopleural hairs. *Legs* yellowish, including all coxæ: femora and tibiæ narrowly but very distinctly black at tips. First hind tarsal segment with a pair of long hairs beneath near base *Wings* (fig 103, a) with three conspicuous brown

spots, one extending from vein 1 over the cross-vein, one in middle of upper basal cell, and one across middle of vein 5, in addition, with faint grey markings, as follows: a patch connecting veins 3 and 4, a broad border to vein 5 2, and a small area on hind border of wing at base. Subcosta ending distinctly before base of sector. Vein 4 with numerous small hairs beneath on its basal section, the hairs extending more than half-way from cross-vein to base of wing. *Hypopygium* (fig 103, b) tergite simple. Cerci finger-like. Coxite small, with large basal lobe and finger-like apical lobe. Style long, slender, curved, with a short, pointed, basal arm.

DISTRIBUTION.—ASSAM: Shillong, 20 x 1920 (*R. Senior-White*), 1 ♂, "in shade at Crinoline Falls," type in Brit Mus, previously determined as *D. montana*.

In hypopygial structure this species appears to be intermediate between the subgenera *Dixa* and *Paradixa*, it also differs from most species of *Dixa* and resembles *Paradixa* in the absence of sternopleural hairs.

3 *Dixa (Dixa) montana* Brunetti, 1911.

Rec. Ind. Mus iv, p 265. TYPE-LOC.—Simla, 7,000' (*Annamdale*).
TYPE ♂ & ♀ in Ind. Mus.

ADULT.—Resembles *D. trinotata*, differing slightly in wing-markings; an irregular pale grey band extends from base of fork of vein 2 to base of fork of 4, an additional small grey spot present towards end of lower basal cell, and another similar spot opposite this below vein 5. Thorax rather darker, with some darkening on shoulders. In a ♀ in the British Museum the first antennal segment is rather long for a member of this subgenus, but slightly swollen towards base, ovipositor rather long, pointed. Hypopygium of ♂ type with style much shorter and broader than in *D. trinotata*.

LARVA (fig. 104, e)*.—As described by Baini Prashad from specimens from Kasauli this appears to be typical of the subgenus *Dixa*, but its specific identity requires confirmation, as there are, no doubt, several closely allied Indian species of this group.

DISTRIBUTION.—Simla district (*Annamdale*); Murree, PUNJAB, 1922 (*C. A. Gull*).

4 *Dixa (Dixa)* sp.

ADULT ♀.—Resembles *D. trinotata* and *D. montana*, but differs slightly from both in wing-markings, and is probably specifically distinct. The three main spots of *D. trinotata* are distinguishable but very faint; spot over cross-vein completely

* Baini Prashad 1918, p. 156.

filling area between sector and vein 1, a small faint grey cloud over base of fork of vein 2, but none between veins 3 and 4, and none over 5.2. Scutellum uniformly darkened. Antennae, wing-venation, and ovipositor as in *D. trinotata*.

DISTRIBUTION.—PUNJAB Kasauli, 5000', 6 iii 1921 (*Barraud*), 1 ♀ in Brit. Mus

5 *Dixa (Dixa)* sp

ADULT ♀.—Much resembles the last, but wing-markings even more reduced, spot over cross-veins smaller and occupying only a small portion of the space between vein 1 and the sector

DISTRIBUTION.—KASHMIR Aran, 8,000', 12 ix 1930 (*Christophers*), 2 ♀♀ in Brit Mus

6. *Dixa (Dixa) ochrilineata* Brunetti, 1911

Rec Ind Mus iv, p 267 TYPE-LOC Kurseong, Darjiling dist. (Ammandale). TYPE ♀ in Ind Mus

ADULT ♂.—*Head* light yellow, including proboscis and tori. *Thorax*. yellow; mesonotum with three rather narrow and well-separated brown stripes, lateral stripes curved towards and over the sides anteriorly. *Legs*. yellowish, tips of femora black. *Wings* with a spot in upper basal cell as in *D. montana* and related species, but differing in having the whole tip darkened, a dark brown streak extending from vein 1 over the cross-veins and continued, though more faintly, over vein 5.2. Ovipositor small, inconspicuous, pale yellow.

DISTRIBUTION.—Known only from type-locality

7. *Dixa (Dixa) zeylanica* Senior-White, 1924.

Spolia Zeylanica xii, p 381 TYPE-LOC. Suduganga, Matale, Ceylon (*Senior-White*) TYPES ♂ & ♀ in Brit. Mus

ADULT ♀.—Differs from most, if not all, other Indian species of the genus in length of proboscis, which is as long as remainder of head, clypeus nearly twice as long as its breadth at the base, palpi inserted well beyond base of proboscis*. *Head*. pale yellowish above. Tori pale yellow, contrasting with the black flagellum; first flagellar segment rather short and slightly thickened. *Thorax*. largely yellowish; mesonotum with three dark stripes, median lighter in front; pleurae with irregular dark markings. No sternopleural hairs.

* The Sumatran *D. binotata* Edw. has a similar proboscis and is in some respects not unlike *D. zeylanica*

Legs: pale brownish, tips of femora rather conspicuously black, femora with a paler pre-apical ring, bases of tibiae and first tarsal segments narrowly paler. *Wings* with a rather large and conspicuous dark cloud around cross-veins, and another across middle of vein 5, a brownish border to vein 5 2, no dark spot in upper basal cell. Subcosta ending well before base of radial sector

♂—Resembles ♀ Hind tarsus at base beneath with one hair longer than the rest, but not conspicuously so. *Hypopygium* a pair of short, bare, pointed, and curved hooks arising from tergite (or from paraproct?), coxite short, with small pubescent apical lobe, style longer than coxite, slightly widened and truncate at tip, with a small, sharp tooth near tip on inner side

DISTRIBUTION.—Known from type-locality in CEYLON, and from Gua, Singbhum dist, ORISSA, 1,500', 5. XI. 1929, ♂ reared from larva in forest-stream (*Senior-White*).

8 *Dixa (Dixa) bistriata* Brunetti, 1911.

Rec. Ind Mus. iv, p 268. TYPE-LOC. Darjiling (*Brunetti*).
TYPE: ♂ in Ind. Mus.

ADULT ♂.—*Head*: dark above First flagellar segment about 5 times as long as broad, but slightly thickened in middle. *Thorax* with yellowish ground-colour, mesonotum with the usual three dark stripes *Legs*: pale, tips of femora narrowly black *Wings* nearly clear, with a small dark brown spot over cross-veins, extending well up into marginal cell, but not reaching vein 1, a brown cloud below middle of vein 5. *Hypopygium*: not examined in detail; style appears rather long, curved, and simple.

DISTRIBUTION.—Known only from type-locality

9 *Dixa (Dixa) bifasciata* Brunetti, 1911.

Rec. Ind Mus. iv, p 269 TYPE-LOC.: Phagu, 9,000', Simla dist (*Annamdale*). TYPE: ♀ in Ind. Mus.

ADULT ♀.—Resembles *D. bistriata*, but first flagellar segment shorter, spot over cross-veins smaller and not reaching far into marginal cell; a more conspicuous dark cloud below basal $\frac{1}{2}$ of vein 5.

DISTRIBUTION.—Known only from type-locality.

10. *Dixa (Dixa) christophersi*, sp. n.

ADULT ♀.—*Head*: entirely blackish, including antennae and palpi. Clypeus and labium short. First flagellar segment somewhat swollen, but over 4 times as long as the greatest

breadth *Thorax*: almost uniformly blackish, with greyish dusting on mesonotum, the appearance varying with incidence of light, seen from in front three broad stripes appear almost without dust, the middle stripe divided by a grey line. Hairs all short and black. Scutellum, postnotum, and pleuræ all dark. A few sternopleural hairs present. *Legs*: rather dark brownish, femora lighter towards base, mid- and hind coxæ pale. *Wings* almost clear, but with a large grey area over cross-veins, not extending up to vein 1 or down as far as vein 5.2. Subcosta ending opposite base of sector. Fork-cells short, upper considerably shorter than its stem. Vein 4 completely bare as far as the cross-vein. *Abdomen* black. Ovipositor with cerci rather long and narrow, last sternite with a quantity of pale yellow hair on posterior margin.

♂—Resembles ♀ in colouring. First flagellar segment of antenna rather shorter. First segment of hind tarsus simple, without long hairs at base beneath. *Hypopygium* (fig. 103, c): small, dark. Coxite small, with small basal lobe but no definite apical lobe. Style short and rather broad, pointed at tip. Tergite simple.

DISTRIBUTION.—KASHMIR Nara Nag, 7,500', ix. 1930 (*Christophers*), 5 ♂, 2 ♀, including types, in Brit. Mus. TIBET · Yatung (*Lt-Col F. M. Bailey*), 1 ♀ in Brit. Mus.

Dixa (*Dixa*) *platystyla*, sp. n.

ADULT ♂—Very similar to *D. christophersi*, differing as follows.—*Thorax*: less dark, and with less dusting on surface, no line of grey dust visible in front view. *Wings* with grey spot in middle rather smaller, scarcely extending into marginal cell. Fork-cells longer, upper distinctly longer than its stem. Vein 4 with rather numerous short hairs before the cross-vein. *Hypopygium* (fig. 103, d) quite different, paraproct with a bare finger-like process laterally, coxite with a short, but distinct, apical lobe, style more flattened, as seen in side view broadest close to base and rounded at tip (not well shown in figure).

DISTRIBUTION.—KASHMIR Nara Nag, 7,500', 20. ix. 1930 (*Christophers*), type and one other ♂ in Brit. Mus.

Subgenus *PARADIXA* Tonnair, 1924.

Rec. Canterbury Mus. n. p. 223. Genotype, *Dixa neozelandica* Tonn (New Zealand).

ADULT.—Antennæ rather longer and more slender than in subgenus *Dixa*, first segment of flagellum cylindrical, not in the least thickened in middle, usually much more than 5 times as long as broad. *Hypopygium* of ♂ with lobe at apex of coxite long, style also usually long and often forked.

LARVA.—Antenna usually with a group of hairs on inner side towards tip. No dorsal rosettes on abdomen. Apex of large lateral plates of spiracular apparatus with a pointed spur. Caudal projection extending well beyond apex of lateral plates of spiracular apparatus.

The larvæ of most, or all, species of this subgenus live in stagnant water.

Dixa (Paradixa) kashmirensis, sp. n.

ADULT ♀.—*Head*. entirely dark above. Clypeus quite short, dark at base, yellowish at tip. Antennæ all dark, first flagellar segment quite 8 times as long as broad. *Thorax* with yellowish ground-colour, slightly dusted with grey. Mesonotum with three broad dark brown stripes, median stripe not distinctly divided, area between stripes and on shoulders in front of lateral stripes somewhat darkened, hairs dark. Scutellum uniformly light brownish. Pleuræ largely dark brown, including *ppn*, an ill-defined pale stripe across middle, lower part of sternopleura also pale. No sternopleural hairs. *Legs*: light brownish, coxæ all yellowish, tips of femora and tibiæ very narrowly and indistinctly darkened. *Wings*. almost clear, but with a small brown spot over c.-v. 3-4 and basal deflection of 3 (this latter being longer than in most Indian species of the genus). Subcosta ending above base of sector, which is slightly longer than in *D. montana* and related species, upper fork about as long as its stem, which is somewhat less curved than usual. Vein 4 with some fine hairs. *Abdomen*: blackish, cerci short and inconspicuous, quite unlike those of most of the Indian species of the subgenus *Dixa*.

DISTRIBUTION.—KASHMIR. Aran, 8,000', 12. ix. 1930 (*Christophers*), type ♀ in Brit. Mus.

Subfamily CHAOBORINÆ.

This subfamily comprises the "phantom-gnats," so-called from the remarkable transparency of the larvæ. Six genera are known, of which only two have hitherto been found within the limits of the Indian region. *Chaoborus* (three species) and *Corethrella* (one species).

The main characters of the subfamily are as follows:—

ADULT.—Mouth-parts short, not adapted for biting; palpi (in both sexes) 4-segmented, curved inwards in repose (not directed rigidly forwards as in Culicinae). Antennal flagellum with 13 segments (as in Culicinae). *Wings* with venation similar to that of Culicinae, and with a fringe of flattened scales

as in that subfamily, sometimes also with definite scales on the veins. *Legs* very hairy, but devoid of scales (differing in this respect from *Culicinae*, in which the legs are always covered with scales). *Hypopygium* of ♂ with large coxite and long, simple, terminal style, no definite chitinisation of proctiger; usually no definite phallosome, but a pair of chitinated processes at base of coxites are regarded as parameres.

LARVA—Very varied in form in the different genera, but in all cases with stout, movable antennae which are provided with 3-5 strong movable spines at the tip, the antennae being used to catch living prey. Mouth-brushes absent, or represented by a group of 10 bristles. Fore gut not communicating with mid-gut and often eversible.

Genus **CORETHRELLA** Coquillett, 1902

Journ N Y Ent. Soc x, p 181. Genotype, *C. brachyleyi* Coq.

Ramcia Annandale, 1911, *Spolia Zeylanica*, vii, p. 187.

ADULT—Very small flies (wing-length 2-3 mm), superficially resembling some *Ceratopogonidae* (such as *Forcipomyia*), but with mosquito-like wing-venation, distinguished from other genera of the subfamily by the relative shortness of vein 1, which ends far before the wing-tip and not far from the tip of the subcosta. Clypeus short and nearly bare. Eyes large, approximated above. Thorax short and rounded. Antennae of ♂ long and densely plumose.

PUPA—Short and broad, capable of very little movement, and rather resembling the pupa of some *Ceratopogonidae*. Paddles represented by a pair of pointed, immovable processes.

LARVA (fig. 104, a).—Head large and broad, with a remarkable row of spines on each side, extending rather obliquely from above downwards. Antennae placed close together in front of head, folding outwards, with three apical spines. Clypeus with four hairs on front margin. Thorax and abdomen provided with long lateral hairs. Segment IX with a short flattened "siphon," which presumably has no respiratory function, as tracheae are not visible within it.

HABITS—Little is known as to the feeding habits of the adults or larvae, but the latter are no doubt predaceous. The habitats of the larvae are varied, the single Indian species hitherto known lives in swamps, but other species are known to inhabit water in tree-holes, leaf-bases, and pitcher-plants, and it is probable that some of these occur in the Indian region.


DISTRIBUTION—*Corethrella* appears to have its headquarters in South America, where a fairly large number of species is known to occur, but the range extends throughout the tropics. Judging from the small number of specimens which exist in collections, the species are never very common.

1 *Corethrella inepta* Annandale, 1911

Spolia Zeylanica, vii, p 189 (*Ramcia*). TYPE-LOC Diyatalawa, 4,300', Ceylon (*Major MacDougal*) TYPE ♂ in Brit Mus

ADULT.—Yellowish species, with a single dark band extending across middle of wing, most distinct on costa. Femora entirely pale, tibiae dark apically, front tarsi somewhat darkened, middle and hind tarsi entirely pale.

Fig 104.



Larvæ of Chaobornæ and Dixinæ: a, *Corethrella calathicola* Edw. (a Malayan species breeding in pitcher-plants); b, antenna of same; c, *Corethrella inepta* Ann., head; d, antenna of same (after Annandale); e, *Dixia montana* Brun.; f, antenna of same (after Baini Prashed); g, *Chaoborus asiaticus* Giles (after Baini Prashed).

LARVA.—Distinguished from some other larvæ of the genus by shape of head (fig. 104, e), which is pointed in front. Antennæ more slender than in some.

DISTRIBUTION.—Known only from type-locality.

Genus **CHAOBORUS** Lichtenstein, 1800.

Wiedemann's Arch Zool 1, p 174 Genotype, *C antisepticus* Licht. (?=*crystallinus* De G.)

Corethra, Meigen, 1803, Illiger's Mag. u. p. 260. Genotype, *C lateralis* Meig. (as *culiciformis* De G.)

Sayomyia Coquillett, 1903, Can. Ent. xxxv, p. 190 Genotype, *C. punctipennis* Say

ADULT —Wing-venation almost as in *Culicinae*, with vein 1 ending near wing-tip and well beyond fork of vein 2. Clypeus rather long and densely hairy. Eyes widely separated above. Thorax longer than broad. Antennæ of ♂ plumose, but rather short. First segment of hind tarsi shorter than second.

PUPA —Abdomen long and narrow. Paddles large, broad, movable, often folded in a fan-like manner. Respiratory horn bladder-like, with minute terminal opening.

LARVA (fig 104, g) —Head much compressed, narrowed in front, without spines on sides, but with 8-10 flattened bristles in middle line beneath, between base of antennæ and labrum. Antennæ folding downwards, with five movable spines at tip. Few conspicuous long hairs on body. No siphon, and no trace of spiracles or spiracular apparatus. A pair of air-bladders in thorax and another in segment VII of abdomen.

HABITS.—The adults are sometimes found in great numbers over and around lakes and large ponds, so far as known they do not feed. The larvæ occupy a horizontal position in the water, remaining at a considerable depth, and feeding upon small Crustacea and other aquatic animals, sometimes on mosquito larvæ. The larva of an undetermined Indian species has been noted by Annandale as living in the interstices of a freshwater sponge.

DISTRIBUTION —More or less cosmopolitan.

The three Indian species belong to the subgenus *Sayomyia*, which is distinguished from typical *Chaoborus* by the absence of pulvilli and some small details of venation. They are all small species (wing 2.5-3 mm.), with almost uniformly pale legs. Many other species of the subgenus possess numerous narrow dark rings on the tibiae, and one or more of these should occur in India. Species of the holarctic subgenus *Chaoborus* are probably to be found in Kashmir.

2. *Chaoborus* (*Sayomyia*) *asiaticus* Giles, 1901.

Entom. xxxiv, p 196 (*Corethra*) TYPE-LOC.: Shalijahanpur, N.W.P., India, 3 x 1900 (Giles) TYPE: ♀ in Brit. Mus.

ADULT* —Clypeus and tori dark. Mesonotum with three dark stripes on integument, but whole surface covered with dense greyish dust. Scutellum with black median line.

* Edwards 1930 a, p. 534.

Wing-veins and the hair clothing them uniformly pale. *Hypopygium* of ♂ with parameres short, enlarged at tip.

LARVA (fig 104, g) *—Apparently differs from those of the European species in the smaller size of the air-bladders in the thorax and abdomen.

DISTRIBUTION.—Widely spread in the Indian region
PUNJAB Karnal (*Barraud*) BENGAL Calcutta (*Annan-
dale*), Madhupur (*Parva*) BIHAR Pusa (*Sharma*) SOUTH
INDIA Coimbatore (*Fletcher*) CEYLON Colombo (*Parva*)

Not certainly known from elsewhere. The Indian species was at one time regarded as the same as the Philippine *C. manilensis*, but is probably distinct.

3. *Chaoborus* (*Sayomyia*) *annandalei* Edwards, 1930

Ann & Mag Nat Hist (10) vi, p 535 TYPE-LOC Mudon,
Anhoret dist, Tonassorum (*Annanale*) TYPE ♂ in Brit Mus

ADULT—Clypeus and tori pale. *Thorax* with scarcely any grey dusting. Scutellum with dark median line faintly indicated. *Wing-veins* all pale, as in *C. asiaticus*. Parameres curved and scarcely swollen apically.

LARVA—Unknown.

DISTRIBUTION—Known only from type-locality

4. *Chaoborus* (*Sayomyia*) *indicus* Giles, 1905

Journ Trop. Med. vii, p 59 (*Curethra*) TYPE-LOC., Seondara,
N.W.P., India (*Giles*) TYPE, ♀ in Brit. Mus.

ADULT ♀—Clypeus black. *Thorax* as seen from in front largely greyish, with indications of three dark lines, as seen from side dark brownish, with lateral margin narrowly yellowish, scutellum with a black median line. *Wings*: pale, except for veins 3, 5, and 5 2, which are dark brown and clothed with blackish hair.

♂ and **LARVA**.—Unknown.

DISTRIBUTION.—Known from type-locality and from SUMATRA

III. FURTHER DESCRIPTIONS OF INDIAN CULICINI.

The following notes and descriptions are based chiefly on some material submitted to me by Capt. Barraud after his volume went to press. Among this material are several strikingly distinct species which the Author was unable to place; as, however, it was clearly desirable that they should be described in this volume, he asked me to report upon them.

* Bani Prasad 1918, p. 154 (*manilensis*).

Genus **ÆDES** MeigSubgenus **OCHLEROTATUS** L. A.64 **Ædes (Ochlerotatus) pullatus** (Coquillett)

A single ♀ collected by Mr T. B. Fletcher at the same time and place as the ♂ described on p. 152 agrees with the ♂ in having broad scales on *apn* and *ppn*, it also agrees fairly well with the ♀ of *pullatus* as described by Peus (1933). According to Peus, the distribution of scales on the pleura affords valuable means of distinguishing some of the closely allied species of the subgenus *Ochlerotatus* in which the tarsi are dark, in *A. pullatus* the meron (hypopleura) is devoid of scales, there is a conspicuous patch of scales immediately below the prothoracic spiracle, and the scales on the mesepimeron extend to, or almost to, the lower margin of this sclerite. These features, together with the absence of scattered pale scales on the costa and on the anterior surfaces of the fore and mid-femora, will separate *A. pullatus* from all other species of the *communis* group in the European fauna.

The specimens referred to on p. 148 as collected by Col. L. G. Browse appear, on re-examination, to be probably *A. pullatus* rather than *A. cataphylla*, as the costa and femora are dark. The scales on *ppn* are not so broad as in the Gulmarg specimens, but the pleural scaling is the same. Several lower mesepimeral bristles are present.

DISTRIBUTION.—KASHMIR: Gulmarg, 5,000' (T. B. Fletcher); Deosai Plain, Shingo River, Chota Deosai, 13,500', and Tsurri, Indus Valley, Baltistan, 7,700', vii. 1923 (L. G. Browse).

Subgenus **FINLAYA** Theo.84 **Aedes (Finlaya) greeni** (Theo.)

This must be regarded as a variety of *A. (F.) aureostriatus* Doleschall, originally recorded from Amboina*. Dr. S. L. Brug has recently obtained in Ceram adults and larvae of a species which is almost certainly the true *A. aureostriatus*, in these specimens the external ornamentation is rather closely similar to *A. (F.) greeni* var. *kanaranus*, except that the abdominal tergites lack basal pale bands, a feature which is often unreliable as a specific distinction. The hypopygium of a ♂ from Ceram appears identical with that of one from Ceylon, in both the hairs at the tip of the basal lobe of the coxite are not simple (as wrongly shown in fig. 37, k, p. 175, of this volume), but are modified into flattened blades. The larval skin of the specimen from Ceram is damaged, but shows no noteworthy differences from the larva of *A. greeni* as described by Barraud.

* *Culex aureostriatus* Doleschall, 1857, Nat. Tijds. Ned.-Ind. xiv, p. 386.

Subgenus *AÊDIMORPHUS* Theo128a *Aedes* (*Aedimorphus*) *punctifemore* (Ludlow), 1921.

Military Surgeon, xlix, December (*Stegomyia*) (♀). TYPE-LOC
 Fort Win McKinley, Rizal, P I TYPE ♀ in U.S. Nat Mus,
 Washington

ADULT*—Very distinct from other members of the subgenus by the black thorax, with spots of white scales arranged somewhat as in *A. (Stegomyia) vittatus*, tarsi unbanded. Wing about 4 mm.

♀—*Head* dorsum covered with flat black scales in middle, a patch of flat white scales towards each side, and many dark upright scales. Palpi dark, with a tuft of pale hairs at tip. Proboscis of moderate length, mainly light brownish, with tip black. Clypeus black, bare. Basal segment of antenna with some flat white scales. *Thorax* with integument black, somewhat shiny, mesonotum clothed for the most part with small, narrow black scales and with small, flat, silvery-white scales arranged chiefly in seven small spots, distributed as follows: a pair of spots close to front margin; another pair about middle of mesonotum, directly behind first pair, a median spot at anterior end of ante-scutellar bare space, and a spot at each side of this space immediately in front of the scutellum. Scutellar scales flat and white. Scales on *apn* white, few in number, those on upper part of *ppn* narrow and black. *Wings*: dark scaled. *Legs*: black, including all tarsi, femora and tibiae with some scattered silvery-white scales tending to form small spots; front tibia with a silvery-white spot at tip, less distinct white tips to other tibiae and to femora. *Abdomen* black, tergites with basal lateral white spots.

♂.—Unknown in India. *Hypopygium*, as described by Dyar, similar in type to that of other members of the subgenus.

LARVA.—Unknown.

DISTRIBUTION—Two females taken at Gaya, BIHAR, 1928 (*V T Korke*). Otherwise known only from Rizal, PHILIPPINE ISLANDS

Subgenus *CANCRÆDES* Edw.

The species described on p. 444 is referred to this subgenus with considerable hesitation, but, according to the present classification, it cannot well be placed in any other group. In ornamentation it is, perhaps, more like *A. (Aedes) yusafi* than any other Indian species, but can hardly be placed in the subgenus *Aedes* on account of the broad scutellar scales.

* Dyar and Shannon, 1924, p. 75 (*A. (Theobaldii)*). Dyar, 1925, p. 217.

170a *Aedes* (*Cancraëdes* ?) *kanarensis*, sp nTYPE-LOC Yellapur, N Kanara, 6 x 1921 (*P. J. Barraud*).

TYPE 5 ♀♀ co-types in Brit Mus

ADULT —A medium-sized blackish species, well characterized by the white longitudinal stripe on pleuræ, single white ring on hind tarsi, and tufted abdominal sternites, as in some species of the subgenus *Finlaya*. Wing about 3.5 mm.

♀—*Head* scales on dorsal surface all broad, flat, and close-lying, almost all black, but a few white ones around eye-margins, and a small white patch low down on each side. a very few upright scales on nape. Proboscis about length of fore femur, slender, dark. Palpi about $\frac{1}{2}$ length of proboscis, dark. Clypeus blackish, bare. Tori dark brownish, bare. Eyes only very narrowly separated. *Thorax*: integument entirely blackish, mesonotum clothed for the most part with narrow, straight, blackish-brown scales, but with a rather broad border of narrow white scales extending round front margin and back to wing-base on each side. No dorso-central bristles, and, so far as can be seen, no prescutellars. Scutellum densely covered with broad, flat, blackish scales. Pleuræ largely bare of scales, but with a conspicuous white stripe extending from *apm* to above base of hind coxa, *apm* covered with flat white scales, *ppm* largely bare, with a few narrow blackish scales above and broader ones below. About 4-6 posterior pronotal and a similar number of postspiracular bristles, latter pale in colour. The small lateral sclerite of the mesonotum (in front of wing-base) clothed with flat white scales, forming part of the white border. *Wings* dark scaled, scales dense, those in plume series rather broadly ligulate. Venation normal, fork-cells about as long as their stems, bases about level. *Legs* mainly blackish, coxæ yellow, clothed in front with white scales, undersides of femora yellowish, hind femora almost entirely yellowish on the basal $\frac{1}{2}$, pale knee-spots present but quite inconspicuous. First segment of mid- and hind tarsi with a white ring at base, all tarsi otherwise entirely dark. Claws simple. Hind tibia as long as front pair, first hind tarsal segment noticeably shorter than tibia. *Abdomen* scaling of dorsal surface entirely black, with slight bluish lustre, tergites with small lateral basal white spots not visible from above. Sternites largely whitish, with black scales on hind borders, those on sternites VI and VII (and to a less extent on V) long and suberect, giving a tufted appearance. Segment VIII in all specimens entirely concealed within VII.


♂ and LARVA —Unknown.

DISTRIBUTION —Malabar coast only; type-locality, as given above, also x 1915 (*Khazan Chand*)

Subgenus **INDUSIUS**, nov.

This new subgenus is proposed for the species of which the male sex is described on p. 269. Having now examined the two specimens referred to, I believe there can be little doubt that they are males of *A. pulverulentus* Edw. they agree in chaetotaxy with the genus *Aedes* (presence of squamal fringe, absence of spiracular and presence of postspiracular bristles, etc.), they are somewhat smaller than the female *A. pulverulentus* (as might be expected), although very largely denuded of scales, one specimen shows similar scaling of the

Fig. 105



Structural details of *Aedes (Indusius) pulverulentus* Edw., ♂: a, head from above; b, one segment of antennal flagellum; c, d, end of abdomen from side and from beneath (diagrammatic); e, process arising from proctiger or base of coxite; f, style from side; g, front tarsus; h, whole front leg.

head to the female *A. pulverulentus* (large area of flat whitish scales in middle, flat dark scales at sides); and, finally, they were taken in the same area as the female *A. pulverulentus*.

The new subgenus is sufficiently distinguished by the remarkable form of the male palpi, front tarsi, and hypopygium, which in many respects resemble those of the New Zealand genus *Opifex*, unfortunately no definite features can be

adduced by which the female may be distinguished from other subgenera of *Aedes*

The following details can be made out in the two damaged males available —

Head eyes rather widely separated above, and distinctly separated below. Orbital bristles normal, few in number; no fine hairs mixed with the scales on dorsal surface of head (this feature distinguishing *Indusius* from *Opifer*). *Proboscis* rather short, distinctly swollen on distal $\frac{1}{2}$, but narrowed again at tip, provided with long hairs ventro-laterally for almost its whole length. *Palpi* about as long as proboscis, clothed with dark scales, but with few or no hairs, swollen apically, with an incomplete joint before middle, distal portion entire. *Antennae* unusually long in proportion to proboscis, with the plumes reduced, only about 8 hairs in each whorl.

Legs femora rather thick, especially front pair; front tibia also somewhat thickened. Tarsi all unusually short, front and middle pairs shorter than their tibiae, first hind tarsal segment scarcely more than $\frac{1}{2}$ as long as tibia (in ♀ it is longer). Front tarsus with a double row of microscopic spinules along under surface of segment 1, segments 3 and 4 remarkably short, with spines at tip beneath, claws large, equal, and simple. Middle tarsus longer, unmodified, claws slightly unequal, the larger with a strong tooth near base. Empodium large and feathery, as on front legs.

Hypopygium large, coxites large, somewhat conical, without lobes or appendages other than the style, which is subterminal in position, broad but irregular in shape, without appendage; a pair of long bent rods apparently arise from proctiger (or possibly from base of coxite); structure of phallosome not ascertainable in the dissected specimen (tip of abdomen in other specimen missing).

Genus **PARAËDES**, gen. n.

This new genus is proposed for two Indian species which, though very different in ornamentation, and probably not at all closely related, agree in possessing the following characters —

Margin of squama quite bare. Membrane of wing with distinct microtrichia. Pulvilli absent. Anterior pronotal lobes small and well separated. Several posterior pronotal and a few postspracular bristles present, but no spiracular and no lower mesepimeral. Wing-venation normal; vein 6 extending to well beyond the base of the fork of vein 5. Wing-scales normal, not emarginate at tips.

This combination of characters makes it impossible to include these species in any genus of Culicini at present known.

The presence of postspiracular bristles suggests some affinity with the *Aedes* group, but the bare squama, now regarded as a feature of generic importance, prevents their inclusion in *Aedes*. The normal wing-venation, with moderately long upper fork-cell, separates the species from *Uranotarma* or *Zeugnomysia*, the normal wing-scales from *Hodgesia*, the absence of spiracular bristles from *Topomyia* or *Harpagomyia*, and the presence of postspiracular bristles from any of these genera. The structure of the hypopygium of the genotype (described below) is very peculiar, and quite unlike that of any other known species.

Early stages unknown.

Genotype, *P. barraudi*, sp. n.

170 b. *Paraedes barraudi*, sp. n.

TYPE-LOC Virajpet, Coorg, S. India, vi 1927 (*J. D. Bailly*).


TYPE: 2 ♂♂ co-types in Brit. Mus.

ADULT.—A small dark-coloured species without special ornamentation of body or legs, and with short palpi in ♂. Wing about 2.5 mm.

♂.—*Head*: scales on dorsal surface nearly all flat and black, a few narrow whitish scales on nape and a row of similar scales round eye-margins; a small patch of flat white scales low down at sides. Orbital bristles fairly numerous, dark. Proboscis about length of fore femur, slender and dark. Palpi very short, about $\frac{1}{10}$ length of proboscis and projecting beyond clypeus by scarcely the length of the latter. Clypeus and tori dark, bare. Antennae slightly shorter than proboscis, plumes moderately developed, evenly spread round shaft. *Thorax*: blackish, clothed dorsally for the most part with narrow dark brown scales, but with a few white scales on front margin of mesonotum, pairs of small indefinite spots of white scales on disc, and a few such scales on each side of the ante-scutellar bare space; scutellar scales narrow and white. Some flat white scales on *apn* and a few on lower part of *ppn*; some narrow white scales on upper part of *ppn*. Numerous long and rather stout dark bristles on mesonotum. Pleurae in both specimens damaged by pins, but chaetotaxy, as noted in generic diagnosis, can be made out. *Wings*: dark scaled, fork-cells somewhat shorter than their stems, base of *af* slightly nearer base of wing than that of *pf*. *Legs*: dark scaled, undersides of femora pale, inconspicuous white knee-spots present. Claws all small and equal, those of fore and mid-tarsi each with a small basal tooth. First hind tarsal segment slightly shorter than tibia. *Abdomen*: dark above, tergites with small lateral white areas extending from base nearly to hind margin of segments. *Hypopygium* (fig. 106): ♂: better

developed than usual, with a pair of rounded and almost bare lobes placed close together. Coxite nearly cylindrical, more than twice as long as broad, with a tuft of hairs at the somewhat truncate tip, no apical or subapical lobe, *bl* large, arising from base and reaching to tip of coxite, forked at mid-length, inner branch with bluntly rounded tip, outer more pointed, and bearing short bristles at tip, *bl* otherwise bare. A pair of well-developed harpagones present, in the form of long curved rods, as long as coxite, with a pair of spines at tip. Paraprocts represented by a pair of short, bare curved hooks.

Fig. 106.



Hypopygium of ♂ in tergal (left) and sternal view (right) of *Paraëdes barraudi*, gen. et sp. n. Lettering as on p. 4.

Phallosome small, divided into lateral plates. Style broad, without appendage, but with a slender finger-like process on external margin well before middle.

♀ and LARVA —Unknown

DISTRIBUTION —Known only from type-locality.

170c *Paraëdes* (?) *argyrurus*, sp. n.

TYPE-LOC. Nongpoh, Assam, 14 vi. 1921 and in 1927 (*Christophers*). TYPE 2 co-type ♀♀ in Brit. Mus.

ADULT.—Rather small species, with reddish thorax, conspicuous silvery markings on pleuræ and sides of abdomen, and four white rings on hind tarsi. Wing about 2.5–3 mm.

♀.—*Head* scales on dorsal surface all broad and flat, mostly black, but with a broad oblique stripe of white on each side and a small white area in front between the eyes, which are not very widely separated, scales on under surface of head yellowish, a few dark upright scales on nape. Upper pair of orbital bristles strong, others weaker but fairly numerous. Proboscis about length of fore femur, slender, recurved, dark. Palpi about $\frac{1}{2}$ length of proboscis, entirely dark. Clypeus of normal shape, dark, bare. Tori yellowish, bare. *Thorax* integument uniformly reddish-ochraceous; mesonotum clothed rather densely with small narrow brown scales, no lighter scales, even above wing-roots or on front margin, two pairs of dorso-central bristles on disc, in addition to several in front of scutellum. Scutellum largely bare, but with a small patch of flat dark brown scales on each lobe. Pleurae with four patches of flat silvery-white scales, one (the largest) forming a longitudinal stripe immediately below mesonotal border, one on upper part of mesopimeron, one in middle of sternopleura, and one on lower part of sternopleura immediately above base of mid-coxa; *apm* clothed with flat silvery-white scales, *ppm* devoid of scales, as is also the postspiracular area. About 4-5 posterior pronotal and 3-4 postspiracular bristles, few sternopleural. *Wings* dark scaled, scales moderately dense, those in plume series narrow. Venation normal, fork-cells about as long as their stems, bases level. Knob of halteres with black scales at base, silvery at tip. *Legs* dark scaled, undersides of femora yellow, hind femora largely yellow, but dark above to base, all femora (but not tibiae) with small white knee-spots, all tarsi with narrow white rings at bases of each of the first three segments, hind tarsi also with a broad white ring on fourth segment, leaving only the tip narrowly black, fifth segment black. Claws simple. *Abdomen* dorsal surface mainly dark scaled, with narrow and rather indefinite yellowish bands at bases of tergites II-VII, tergite VIII with a conspicuous patch of silvery scales; tergites I-VII each with a lateral subbasal patch of metallic silvery scales; venter mainly yellow. Cerci small and hidden within segment VIII.

♂ and LARVA.—Unknown.

DISTRIBUTION.—Known only from type-locality.

This species rather suggests some members of the subgenus *Stegomyia*, from all of which, however, it differs in the entire absence of white markings on the mesonotum, as well as in the completely bare squama.

Genus **CULEX** L.Subgenus **CULEX**, s. str.

Since this volume went to press Capt Barraud has submitted for study an extensive series of specimens of *Culex* of the *mimeticus* group, chiefly from the Nilgiri Hills and other localities in South India. Further material has also been received from Dr. R. B. Jackson from Hong Kong and from other correspondents in China. The results of the study of this material, in conjunction with that already in the British Museum collection, are given here. It was found that four (instead of three) distinct forms occur in India; another distinct form occurring in Hong Kong is described to complete the account. All these forms are here allotted specific rank, as the distinctions observed appear quite definite, but it is possible that *mimuloides* and the two new forms may be merely varieties of *mimeticus*.

When distinguishing *C. mimulus* from *C. mimeticus* in 1915 I laid most stress on the dark-scaled third vein. This, however, is not constant, the amount of pale scaling on the third vein, also the actual and relative size of the pale costal spots, is subject to a good deal of individual variation in all forms of this group. The best distinctions between the various forms now appear to be those given in the key below.

Key to Species or Varieties of C. mimeticus group.

♀♀.

- | | |
|--|--------------------------------|
| 1 First pale costal spot (at middle of wing) extending only on to subcosta . . . | 2. |
| First pale costal spot extending over vein 1 . . . | 6 |
| 2 Bases of fork-cells dark (sometimes a small pale spot at base of upper fork only); tergite VII broadly pale apically . . . | [p. 452.] |
| Bases of both fork-cells pale, tergite VII narrowly pale apically . . . | <i>fusCIFurcatus</i> , sp. n., |
| 3 Veins 3 and 5.1 pale practically to the tips (Java only) . . . | 3. |
| Veins 3 and 5.1 very distinctly dark-tipped . . . | <i>dsengensis</i> Brug. |
| 4 Fork-cells with their bases nearly level; vein 3 largely pale . . . | 4. |
| Base of <i>pf</i> distinctly proximal to that of <i>af</i> , vein 3 with a smaller pale area or none . . . | 5. |
| 5 Mesonotal scales uniformly reddish-brown Mesonotum with distinct mottling of light and dark scales . . . | [p. 451.] |
| <i>mimuloides</i> Barr., | |
| <i>jacksoni</i> , sp. n., p. 452. | |
| 6 First pale costal spot not extending beyond vein 1, vein 5.2 dark to the tip . . . | <i>mimeticus</i> Noé, p. 451. |
| First pale costal spot extending over vein 4; vein 5.2 broadly pale at tip . . . | <i>mimulus</i> Edw. |
| | <i>orientalis</i> Edw. |

The distinctions noted in this key will apply in the main to both sexes, but, as usual in mosquitoes, the ornamentation of the ♂ is often less well marked than that of the ♀, and specimens of the former sex are best distinguished on hypopygial characters

236 *Culex (Culex) mimeticus* Noé.

ADULT ♀—Mesonotum mottled with yellowish and brownish scales, the lighter scales tending to form three spots on anterior $\frac{1}{2}$, one near front margin and a pair in middle. *Wings* with bases of forks more or less level, or *af* slightly proximal to *pf*, length of forks rather variable, but *af* usually about 3 times as long as its stem, a spot of pale scales always present at base of each fork. Vein 3 almost always extensively pale in middle, leaving base and tip rather narrowly dark, pale spot at middle of costa variable in width, but never involving vein 1, vein 5.2 usually devoid of pale scales at tip. *Legs* with practically the same ornamentation as in other species of the group, hind femora largely dark on anterior surface except towards base, pale beneath almost to tip. *Abdomen* with basal white bands to tergites which do not spread out laterally as seen from above, tergite VII with a narrow apical pale band.

♂—Ornamentation similar. *Hypopygium* as described on p. 411.

LARVA as described on p. 411.

DISTRIBUTION.—In India this species appears to occur chiefly in the north. Specimens are in the British Museum from Bakloh, N.W.P. (*Landesay*); Nara Nag, KASHMIR (*Christophers*), Shillong and Cherrapunji, ASSAM (*Senior-White*), and Pusa, BIHAR (*Sharma*). No specimens have been seen from the hills of South India, but two atypical ♀♀ from CEYLON appear to belong here rather than to *C. mimuloides*.

More or less typical specimens have been examined from MACEDONIA, PALESTINE, TIBET (Yatung), NORTH CHINA (Pekin), SOUTH CHINA (Nanking District, Hangchow; Kiangsi, Chekiang), HONG KONG, and FORMOSA. So far as can be judged from the description, the Algerian *C. pseudo-mimeticus* Sergent does not differ appreciably from the typical form of this species.

236 a. *Culex (Culex) mimuloides* Barr.

ADULT ♀.—*Thorax* as in *C. mimeticus*. *Wings* with fork-cells very long, base of *pf* very noticeably proximal to that of *af*, both forks with distinct spot of pale scales at base. Scales often appearing shorter and less dense, and wings rather

narrower than in *C. mimeticus*. Vein 3 usually with a short pale area in middle, sometimes more extended, but occasionally reduced or absent. Vein 5 2 usually with a few white scales at extreme tip. Pale spot in middle of costa as in *C. mimeticus*, never involving vein 1. Legs almost as in *C. mimeticus*, but hind femora usually rather extensively dark beneath towards tip, and often with a pale line on anterior surface extending almost to tip between the dark dorsal and ventral areas. Abdomen as in *C. mimeticus*.

♂ Ornamentation similar, but wing-markings often less distinct, position of forks as in ♀ *Hypopygium* as described on p 411.

LARVA not yet isolated.

DISTRIBUTION.—Confined, so far as known, to South India. NILGIRI HILLS Ootacamund (*Fletcher, Puri*); Coonoor (*Puri*), various unrecorded localities (*Khazan Chand*), PALNI HILLS Kodaikanal (*Patton*).

Culex (Culex) jacksoni, sp. n.

ADULT ♀—Closely resembles *C. mimeticus* in almost all respects, but mesonotal scales almost uniformly reddish-brown, without trace of pale spots on anterior $\frac{1}{2}$, a few pale scales present around margin and in front of scutellum.

♂—Thoracic scaling as in ♀. *Hypopygium* closely resembling that of *C. mimeticus*; no obvious differences discoverable in the single specimen available.

LARVA.—Quite distinct from the typical form of *C. mimeticus*, and more resembling, though equally distinct from, *C. mimulus*. Antenna as in *mimulus*, with subapical bristles fairly near tip, and some spinelets beyond tuft. Mentum with fairly regular teeth. Comb-teeth as in *mimeticus*, ending in a long and strong spine. Siphon short, scarcely 5×1 , pecten of about 15 teeth, extending nearly $\frac{1}{2}$ length of tube, teeth rather short and with strong denticles, as in *mimulus*, only three pairs of ventral tufts, which are little, if any, longer than diameter of tube, one small lateral tuft near end of tube. One of the two isolated skins (♀) shows 2 short thick spines on distal $\frac{1}{2}$ of siphon-tube on each side, the other (♂) has no distinct spines, but some thickenings of the chitin. Anal segment with *sc* 2-3-branched, *lh* quite short and 2-branched.

DISTRIBUTION.—HONG KONG. Described from 1 ♂, 1 ♀, with isolated larval skins, from Shonson Hill, 18. iii. 1932 and 16 ii 1933 (*Dr R. B. Jackson*). Types in Brit. Mus.

236 b. *Culex (Culex) fuscifurcatus*, sp. n.

ADULT ♀.—Mesonotum somewhat less mottled and more reddish-tinged than in *C. mimeticus*. Wings with forks of moderate length, bases about level and usually entirely

without pale scales, though a few such scales may be present at base of *af*. Vein 3 extensively pale in middle, dark at base and tip. No white scales at tip of vein 5 2. *Legs* as in *C. mimeticus*. *Abdomen* with basal pale bands tending to spread out along sides of tergites, tergite VII with a broad apical pale band.

♂—Ornamentation as in ♀, in one specimen the lateral extension of the basal bands of the tergites is even more marked. *Hypopygium* with coxite, style, and phallosome almost as in *mimeticus*, but differing strikingly in the complete absence of the basal arm of the paraproct.

LARVA.—Unknown

DISTRIBUTION.—Known only from NILGIRI HILLS. Ootacamund, 7,500', XII. 1913, 3 ♂♂, including type, in Brit. Mus. (*T. B. Fletcher*), 1 ♂, 3 ♀♀, locality unspecified, x. 1915 (*Khazan Chand*).

ALPHABETICAL INDEX.

All names printed with a small initial letter indicate a species or variety.
All names printed in bold type are valid names of Indian Culicidae, names of genera or groups have a capital initial letter
All names printed in italics are synonyms
Thickened numerals refer to the page on which the "Systematic" treatment of the insect is given.

- abditus**, 279, 282.
acaudatus, 8
achates, 309
Aedeomyia, 131.
Aedes, 3, 28, 32, 33, 119, 124, 176, 182, 216, 300, 311, 422, keys, 136-143 Subgenus, 2, 4, 136, 137, 138, 277, 297, 443; keys, 278-280.
aedes, 358, 358
Aedimorphus, 119, 137, 138, 147, 148, 248, 270, 271, 278, 387, 443, key, 247-249
Aedomysia, 2, 31, 33, 181, 135, 157
egypti, 1, 5, 6, 140, 218, 219, 220, 221, 225, 227
anea, 38
affinis, 37, 38, 42, 43, 44, 45
africana Theo (*Diceromyia*), 271.
africanus Theo (*Tanorhynchus*), 130.
ager, 391, 393.
agrestis, 280, 293.
Agrostomyia, 277.
albipes Edw., 12, 14, 15, 16, 21
albipes Leio, 97, 98, 101, 102.
albipes Theo, 225
albipennis, 225
albocinctus, 60, 62, 63, 64.
albocinctus, 142, 154, 156, 172.
albolateralis, 140, 154, 205, 208, 210, 212.
albolineatus Giles, 420, 422.
albolineatus Theo, 137, 141, 217, 220, 243.
alboniveus, 140, 210, 211, 212.
albopictus, 5, 6, 141, 220, 233, 235, 237, 238, 239, 240, 241, 242.
albopleura, 381
albopunctata, 245.
alboscutellatus, 240, 248, 249, 250, 251, 252.
albotæniatus, 153, 156, 174, 176.
albus, 406.
Allotheobaldia, 80
alorensis, 240.
alternans, 144.
ambiguus, 391, 393, 394.
amboinensis, 24
ampra, 54.
andamanensis Barr., 97, 98, 101, 102.
andamanensis Edw., 279, 280, 290, 291
andrewsi, 240.
angulatus, 378.
Anisochaleomyia, 56.
annandalei Barr., 60, 62, 75, 356.
annandalei Edw., 441.
annandalei Theo., 189, 220, 227, 228, 229, 230, 281
annulata Schrk., 86, 93.
annulata Theo., 383.
annulatus Tayl., 369.
annuliferus, 272.
annulifera, 126, 127, 128, 129, 130, 131.
annulipennis, 321, 327.
annulipes, 136.
annulipes, 213, 215.

- annulitarsis**, 320, 321, 324, 325, 327, 328, 331
annuloabdominalis, 381
annulus, 404
Anopheles, 16, 32, 59, 94, 132, 409, 429
anopheioides, 97, 98, 101, 102, 103, 104
 antiseptious, 440
apicalis Adams, 347
apicalis Theo. (*Brevihynchus*), 321
apicalis Theo. (*Desvoides*), 319
apicalis Theo. (*Lecestera*), 321
aranoides, 35, 37, 38, 42, 44, 45
arboricollis, 103
argenteomaculata, 232
argenteotarsis, 24
argenteoventralis, 46
argenteus Ludi, 406
argenteus Poir., 221
argentinotus, 250
argyrotarsis, 70
argyrurus, 448.
Armigeres, 31, 32, 33, 135, 219, 310, 324 Subgenus, 313, 320, key, 313
asiaticus Edw., 150, 151.
asiaticus Giles, 440, 441
assamensis, 143, 151, 157, 163, 164, 165, 166, 169.
atra, 60, 62, 72.
atritarsis, 221, 222, 223
atrius, 280, 288, 293.
aurantapex, 393
aurantus, 144, 145
aurea, 100
aureolineatus, 313, 319.
aureopunctus, 344
aureostratus Leic., 144
aureostratus Dol., 442
aureoventer, 45, 46.
aureiflavus, 25
aurontens, 155, 202.

bahri, 378, 385, 386.
ballyi Barr (*Culex*), 378, 382, 413.
ballyi Barr (*Hodgesia*), 53, 55.
bambusa, 40
banksi, 180
Banksinella, 136, 137, 269.
barraudi Edw. (*Oulex*), 337, 389, 402, 403, 405, 419, 425
barraudi Edw. (*Paraides*), 447.
Barrandius, 334, 335, 346.
bengalensis, 370, 371, 372.
berlandi, 151.
bernardi, 370
bicolor, 61, 81, 83.
bicornuta, 374.

bifasciata, 431, 485.
bigoti, 338
bimaculata, 61, 78.
bimaculata, 65
binotata, 434
blpunctatus, 407, 408.
biroi, 404
bistrata, 431, 485.
bltæniorhynchus, 335, 387, 388, 391, 393, 394, 396
Blanchardomyia, 313
Bolbodeomyia, 399
 bonnem, 128.
brakoleyi, 438
brevicellulus, 120
brevipalpis Giles, 334, 337, 348, 351, 352
brevipalpis Theo., 8
Brevihynchus, 320
brugi, 257
butleri, 277, 279, 280, 296.

cacharanus, 156, 166.
cæus, 139, 246, 247, 248, 250, 257.
calopus, 221
campestris, 60, 62, 68.
cancer, 72
Canaraides, 137, 138, 297, 443.
canerlecomes, 297, 298.
caspius, 140, 148.
castrensis, 330, 354, 355, 356.
cataphylla, 148, 442
catasctica, 132, 134
Catataxomyia, 217
cautus, 279, 280, 288.
Ceratopogonidae, 428, 429
ceylonensis, 34, 42
ceylonica Theo. (*Ouhciomyia*), 385.
ceylonica Theo. (*Uranotenia*), 72.
ceylonicus Edw., 270, 288.
Chatomyia, 320
chamberlaini, 105, 107, 108, 116, 289
chandi, 301, 304, 308.
 Chaobornæ, characters, 437.
Chaoborus, 427, 429, 437, 440.
chemulpoensis, 225
Chironomidae, 428, 429.
christianus, 232
christophersi Barr, 60, 64.
christophersi Edw. (*Aedes*), 141, 154, 156, 185, 195, 198, 202.
christophersi Edw. (*Dixa*), 431, 435, 436.
christophersi Theo., 420.
Christophersomyia, 137, 138, 212; key, 213
Chrysocnops, 119.

- chrysogona*, 122
chrysolineatus, 142, 155, 185, 187, 188, 189, 190, 191
chrysoscuta, 267, 268
cinctellus, 338, 362, 363, 366, 307, 368
cinereus, 134, 216
cingulatus Dol., 420
cingulatus Leic., 320, 328, 329, 330
clavatus, 279, 294.
clavipalpis, 105, 108, 100
cæruleocephala Leic., 34, 40
cæruleocephala Theo., 44.
cærulescens, 362
cogilli, 156, 164, 165, 166
Colonomys, 34
comatus, 280, 292.
communis, 442
completus, 113
complex, 299, 301, 302, 308
concolor, 338, 341, 342
confirmatus, 147.
conopas, 121
Conomyia, 105
Coquillettia, 32, 118, 119, 124, 125, 135, key, 120
Corethra, 440
Corethrella, 428, 437, 438.
cornutus, 335, 388, 391, 395, 396, 397
coveilli, 301, 302, 304, 306, 307
craggi, 220, 228, 229, 230
crassipes Theo., 310
crassipes v. d. Wulp, 120, 122, 319.
crystallinus, 440
Culex, 2, 4, 5, 28, 30, 32, 33, 86, 119, 145, 183, 332, 334, 338, 399, 450, keys, 334-338. Subgenus, 338, 387, 450, keys, 388, 450.
Culicidæ, characters, affinities, 427-429
culiciformis, 440
Culicini, 6, 28, keys, 31-33
culleinus, 249, 252.
Culicomyia, 334, 335, 367, 376, 387, 413, key, 378.
Culineta, 86
cuneatus, 407, 408.
curtipalpis, 351, 362.
curtipes, 298.
Cyathomyia, 350.

Danielsia, 153.
Dasyomyia, 105.
decanus, 143, 157, 161, 163, 164, 165, 166
Dendroskusea, 271.
dentatus, 321, 331.

desmotes, 138, 218, 219, 222, 223, 225.
Desoudeia, 313
Desoudeia, 313
dibrugarchensis, 317, 318, 319
Diceromyia, 137, 138, 246, 271, key, 272.
diengensis, 409, 450.
digitatus, 320, 321, 330.
discrepans, 309, 310
Diseaso, relation of Culicids to, 5-6, 423
dissimilis, 143, 153, 155, 203, 204.
divina, 296
Dixa, 430, key, 430 Subgenus, 430, 431, 433, 436
dofleini, 38, 45.
doleschalli, 420
dolichocephalus, 320
domestica Theo., 246.
domesticus Lolo., 393
durhami, 314, 317, 318, 319

caulitha, 346
leucex, 246
edwardsi Barr. (Aedes), 220, 232.
edwardsi Barr. (Culex), 388, 397.
edwardsi Barr. (Megarhinus), 14, 15, 23, 24
edwardsi Barr. (Tripteroides), 34, 38, 41, 300
edwardsi Barr. (Uranotaenia), 57, 60, 65.
Ekrinomyia, 144
elegans, 113
elsiae, 130, 154, 157, 177, 180, 181, 183, 184.
emmaeana, 362.
epidesmus, 388, 389, 396.
erithrosops, 253.
ethiopicus, 393
Etorleptomyia, 105
Etorleptomyia, 105, 108.

falcipes, 74.
fasciatus, 210, 221.
fatigans, 1, 2, 6, 252, 338, 340, 345, 380, 417, 420, 422.
feegradei, 143, 156, 163, 164, 166
Ficalbia Leic., 216
Ficalbia Theo., 32, 33, 58, 105, 135; keys, 107-108.
fideis, 348.
Finlaya, 2, 4, 5, 137, 138, 148, 153, 219, 444; keys, 155-157, 163.
Finlayia, 153.
fisheri, 256

- flavicornis*, 363, 375.
flavicosta, 97, 98, 100, 103, 104.
flavithorax, 97, 98, 100, 103, 104, 105.
flavopictus, 141, 220, 237, 239.
flavus, 300, 311, 312, 313, 319, 320, 321, 325, 326
fluviatilis, 191
foliatus, 350
formosensis, 142, 155, 189, 190
fouchowensis, 420
fragilis Leic., 281
fragilis Ludl., 376, 378, 385, 386, 387
fraudatrix, 337, 359, 362, 363, 365, 366, 367, 368, 371
funerea, 116, 139, 301, 302, 307
fusca Leic (*Fleaiibia*), 108, 110, 115.
fusca Leic (*Skeatomyia*), 34, 42
fusca Leic (*Uranotenia*), 83
fusca Theo (*Desvoides*), 314
fuscus, 338, 340, 341, 343, 344, 393
fusolfurcatus, 450, 452.
fusoltarsis, 389, 426.
fuscocephalus, 338, 389, 410, 420, 423, 424, 426
fuscopteron, 120
fuscum, 347
fuscus Theo (*Pectinopalpus*), 376
fuscus Theo (*Trichorhynchus*), 376, 385

gardneri, 233
gebeleimensis, 242
Gestonomyia, 246
gellidus, 337, 338, 394, 406, 407
genurestris, 47, 48, 51, 52
gibbini, 121
gigantulus, 22
gilesi, 24
gilli, 141, 154, 156, 173, 185, 195, 196, 198
glaphyoptera, 97
Grahamia, 330
gracilis, 225
graminus, 385
grata, 8, 24
gravelyi, 14, 15, 16, 20.
greeni Theo (*Aedes*), 141, 154, 157, 184, 185, 196, 198, 202, 442.
greeni Theo (*Helzmannia*), 301, 304, 308.
greigi, 191
gubernatoris, 143, 153, 154, 157, 159, 162, 163, 165, 166, 168, 169, 170; key to group, 163.
guttipennis, 431

Hæmagogus, 3, 31, 32, 33, 135, 309.
hæmorrhoidalis, 8
hallfaxi, 340, 344.
Harpagomyia, 2, 32, 33, 46, 47, 48, 52, 135, 447
harveyi, 142, 155, 188, 190
hatsensis, 296
hayashii, 351
hebes, 61, 81.
hebrideus, 240.
Helzmannia, 2, 3, 31, 33, 116, 135, 138, 299, 300, 309, keys, 300-302
hewitti, 362
himalayana, 199
himalayensis, 141, 300, 301, 302, 306
hirsutipleura, 279, 280, 291, 292, 293
Hodgesia, 32, 33, 52, 58, 135, 447, key, 53
horishensis, 227
Howardina, 184, 242
Huleceteomyia, 153
hutchinsoni, 389, 428.
hybrida, 33, 40, 107, 108, 111, 112, 114, 115

ibis, 213, 215.
imitator, 232
immsericors, 24
impellens Theo, 400
impellens Wlk., 398.
imprimens, 257
Inchoatus, 321, 328, 330, 331, 332.
indecorabilis, 281.
Indiana, 126, 127, 128, 130.
Indica Edw., 88, 92, 93.
Indica Theo (*Helzmannia*), 300, 301, 302, 308, 304
indica Theo (*Neomaoleaya*), 290
Indiens Barr., 34, 37, 39, 41.
Indicus Giles, 441.
Indicus Theo (*Aedes*), 139, 277, 279, 280, 282, 283.
Indusius, 268, 445, 446.
inelegans, 424
inepta, 439.
infantulus, 363.
infula, 391.
Ingramia, 118.
inornata Theo (*Culicomyia*), 376, 385
inornata Theo. (*Squamomyia*), 34, 42.
inquinatus, 157, 169.
intermedia, 108, 109.
iphis, 353, 354.
Iyengar, 141, 272, 273, 274.

- Jacksoni**, 450, 452.
Jacobsoni, 48, 51.
Jamesi, 247, 249, 250, 251, 252
Jamena, 338
japonicus, 186, 187
Jensoni, 351, 359, 302
joloensis, 314
jugraensis, 180, 191
justapallidiceps, 363

kanaranus, 157, 184, 185, 442
kanarensis, 444.
karwari, 204.
kashmirensis, 431, 437.
kempli, 14, 15, 18.
khariana, 190, 191
khazani Edw. (Aedes), 143, 150, 168,
109
khazani Edw (Culex), 337, 353, 354,
355, 357, 359.
kochi, 158.
koreicus, 187
kotiensis, 143, 159, 161.
kuchingensis, 313, 314, 316, 317,
318

laniger, 147.
lateralis Ludi, 72
lateralis Mg., 440
leicesteri Edw (Aedes), 291
leicesteri Edw. (Uranotaenia), 83.
leicesteri Theo, 18, 19
Leicesteria, 2, 31, 313, 319, 320;
key, 320.
Leicesteriomyia, 320
lepchana, 174
Lepidotomyia, 153, 240
Leslieomyia, 246
Leucomyia, 387
lineatopennis, 100, 140, 269.
littoralis, 264
longiareolata, 86, 88.
longifurcatus, 378
longipalpis Leic., 320, 321, 328,
329.
longipalpis v. d. Wulp, 0, 126, 128,
129, 131.
longipes, 348.
longirostris Leic. (Aedes), 142, 216,
298.
longirostris Leic. (Uranotaenia), 60,
62, 71.
Lophoceratomyia, 2, 334, 335, 351,
359, 367; key, 362.
lophoventralis, 153, 156, 167.
loricatus, 406

Iowisi, 248, 250, 251
luolensis, 221, 222.
ludlowi, 111
Ludlowia, 105
lugubris, 280, 294.
luridus, 341
lutea, 153
luteoabdominalis, 389
luteola Edw, 61, 80, 81
luteola Theo., 424
luteolateralis, 269
lutescens, 80
Lutzia, 30, 145, 333, 334, 335, 338,
340, key, 340
luzonensis, 105, 108, 110, 112, 113,
110

macdougalli, 143, 157, 177.
macfarlanei Edw (Aedes), 180, 181.
macfarlanei Edw (Uranotaenia), 60,
62, 70.
mcgregori, 102
macropus, 348
maculata Mg, 430, 431
maculata Theo. (Orthopodomyia),
97, 98, 101.
maculata Theo (Pecomyia), 246,
260
maculata Theo (Pseudograhamia),
246, 258.
maculatus Theo (Anopheles), 286
maculipennis, 430, 431.
maculipes, 97, 98, 101.
maculipleura, 61, 81, 82.
magna Theo. (Aedes), 153, 159, 161.
magnificus, 21
magnus Theo (Armigeres), 320,
321, 324.
Malaya, 47
Malaya, 47
malayi Leic (Culex), 334, 336, 352,
354, 355, 359, 358, 359
malayi Leic. (Hodgesia), 53, 54.
malayi Theo, 313
malfeyti, 118
mammillifer, 337, 363, 370, 371, 373,
374.
manicatus, 16
manilensis, 441.
Mansonia, 32, 33, 112, 115, 118,
135.
Mansonloldes, 4, 32, 118, 123, 135;
key, 126.
margarsen, 290
maxima, 61, 77.
mediolineata Theo. (Ficofibia), 105,
113.
mediolineatus Theo. (Aedes), 249,
248.

- medlopunctatus*, 139, 220 280.
Megarhinini, 7, 8.
Megarhinus, 4, 8, 13, 30, 132, keys, 14-16
meronephada, 217, 232
metallica Leic (Ficalbia), 105, 109
metallica Leic. (Helzmannia), 301, 304, 306, 307.
metallicus Leic (Megahmus), 20
micans, 65
microannulatus, 308
miropterus, 141, 271, 272, 275, 277
mikranus, 156, 157.
mimeticus, 337, 389, 409, 411, 412, 450, 451, 452, 453, key to group, 450.
Mimomyia, 32, 105, 107, 108
mimuloides, 400, 411, 412, 450, 451.
mimulus, 337, 389, 409, 411, 412, 450, 452
minima Ludl , 111, 112
minima Theo (Ficalbia), 105, 108, 116.
minimus Leic , 124
minimus Theo (Megarhinus), 14, 15, 16, 22.
minor Leic (*Culex*), 337, 363, 370, 372, 373 374 .
minor Leic (Topomyia), 45
minor Theo , 420
minuta Theo (*Culicada*), 253
minuta Theo (*Mimomyia*), 116
minutissima Theo. (*Stegomyia*), 232
minutissimus Theo (*Culex*), 337, 362, 368, 365, 360, 389
Mochthogenes, 334, 335, 352, 361, key, 354
modestus, 335, 346.
monetifera, 40
montana, 431, 432, 438, 434, 437
Mucidus, 30, 136, 138, 144, 148
mucidus, 147
multimaculosus, 344
Mycetophylus, 428

navalis, 362
nebulosus, 376
Neoculex, 335, 347, 352, 353, 361
Neodixa, 427, 429
Neomacleaya, 277.
Neomelanocoryn, 376
neozelandica, 436
nepenthicola, 34.
nepenthes, 42
nigerrima, 363.
nigrescens, 355, 358
nigricephala, 398.
nigritarsus, 98, 101.

nigritia, 233
nigropunctatus, 338, 378, 383, 384, 385
nigrorhynchus, 188, 189.
nigrostriatus, 248, 262.
nirotarsus, 283
nigriticus, 389, 418.
nitidiventer, 40
nippom, 253
niveitaniata, 86, 88, 91, 93, 94
niveoides, 140, 210, 211.
niveoscutellum, 248, 251, 252
niveus, 140, 154, 155, 205, 208, 211
nivipes, 56, 65
nivipleura, 61, 76, 77 .
nongpohensis, 317, 318, 319
notoscriptus, 178
novalbopictus, 141, 220, 237.
novobscura, 61, 62, 84.
novochracea, 120, 121, 122
novoniveus, 139, 211.
nummatus, 137, 143, 246, 247, 248, 265.

obscura, 84, 85
obturbans, 310, 312, 313, 314, 316, 317, 318, 319, 320
ocellata, 391
Ochlerotatus, 137, 138, 145, 147, 167, 182, 246, 442.
ochracea Theo (*Grahamia*), 389.
ochracea Theo (*Mansonia*), 120, 122.
ochraceus Theo (*Aedes*), 261, 419
ochrilineata, 431, 434.
Oculeomyia, 387
omissus, 321, 330.
omuscus, 250
Opifex, 445, 440
orbite, 249, 250
O'Reillya, 105
oreophilus, 143, 155, 171, 192, 194, 200
orientalis Barr , 60, 66.
orientalis Edw , 409, 450
Orthopodomyia, 32, 33, 94, 135, keys, 97-98.
ostentatio, 249, 267.

pagei, 267
pallidostriatus, 142, 248, 261, 263, 264
pallidothorax, 338, 378, 380, 381, 382
pallirostris, 155, 190.
pampangensis, 246, 250, 252.
panalektoros, 314

- Panophies*, 118
Paradixa, 430, 433, 436.
Parædes, 446
parascelos, 261
Pardomyia, 144
Pecomyia, 246
pectinatus, 320
Pectinopalpus, 376
pendulus, 320
perezignus, 418, 419
periskeletus, 138, 271, 272.
perplexus Leic (Aedes), 230, 231.
perplexus Leic (Culex), 400
perturbans, 119
pettgrewi, 414
Phagomyia, 153
philippinensis, 34
Philodendromyia, 359
Phoniomyia, 303
pilosa, 302
pipersalatus, 142, 247, 248, 257,
 258, 260, 261
pipiens, 387, 417
plantaginis, 363, 372.
platystyla, 431, 436.
plegipennis, 406
plumosa, 38.
pluvialis, 334, 354, 356.
pocellus, 143, 155, 157, 158
poecilus, 153, 157
Popea, 153
powelli, 34, 37, 39
prominens, 156, 168, 169.
Protomelanoconion, 347
proxima, 40
pseudalbopletus, 141, 220, 235.
Pseudocarroliia, 153
pseudodiurnus, 279, 295.
Pseudograhamia, 246
Pseudograhamia, 45
Pseudohowardina, 267.
pseudolongifurcatus, 378.
pseudomediofasciatus, 279, 280, 286,
 288
pseudomimetacus, 451
pseudoniveus Theo. (Aedes), 205,
 208
pseudoscutellaris, 240, 241.
pseudostenetrus, 255
pseudotæniatus, 143, 154, 157, 177,
 178, 180
Pseudotheobaldia, 86
Psychodidæ, 428.
pulcherrima, 56.
pulchritarsis, 141, 150.
pulchriverter, 143, 154, 155, 187,
 199, 202.
pullatus, 143, 152, 153, 442.
pullus, 383
pulverulentus, 249, 268, 445
punctifemore, 443.
punctipennis, 440
punctipes, 272, 273.
punctissimus, 273
purri, 229, 230
pusillus, 345, 347
pymæus, 120.
quadricinctus, 227, 228.
quasiferox, 25
quasimgritia, 233
quasipipiens, 420
quasisanguinea, 54
Quasistegomyia, 217
queenslandensis, 221, 222.
Rachionotomyia, 34
Radioculex, 105
Ramcia, 438
rami, 279, 295.
raptor, 340, 343, 344, 345, 393
recondita, 61, 77, 83.
rectirostris, 324.
Reedomyia, 246
reesi, 420.
reginæ, 141, 272, 277.
regius, 24
Rhinoskusea, 136, 137, 138, 216.
Rhyncotænia, 118.
rima, 376
rizali, 186
roperi, 72
rubithoracis, 302, 363, 367.
Rumchomyia, 34.
rutherfordi, 60, 68.
samarensis, 233.
sanguinea, 52
sarawaki, 387, 391.
saxicola, 142, 155, 137, 191, 371.
Sayomyia, 440.
scapularis, 147.
scatophagoides, 138, 145, 147.
scutellaris Theo (Aedes), 233, 234.
scutellaris Wlk. (Aedes), 6, 141, 220.
 240.
Scutomyia, 217.
seguini, 127.
seniori, 362, 365.
sepositus, 394
septemguttata, 123, 127.
septempunctata, 129.
sericeus, 420.
serrata, 38, 42, 43.
setulosus, 341.
shebbearei, 328, 367, 378, 380, 381,
 382.

- shallongensis, 317, 318
 shortii, 139, 154, 156, 157, 181, 183.
 sigmoides, 280, 295.
sukkimensis, 24
 similis Leic, 37, 38, 41
 similis Strickl, 303.
 similis, 156, 198.
 simplex Theo (Aëdes), 298.
simplex Theo (*Neomacleaya*), 290
simulatus, 180
 Simulidae, 428, 429
 sinensis, 335, 388, 394, 300
 singularis, 291
 sintoni, 157, 200.
sitiens Theo, 404
 sitiens Wied, 336, 338, 307, 398, 402
Skeironomyia, 34
 Skusea, 273, 275, 277, 281
spathipalpis, 88.
 splendens de Meij, 47
 splendens Wied, 13, 14, 15, 18, 24, 25
 splendens Will, 300
squamipenna Theo, 132
squamipennis L. Arrib, 131, 132
Squamomyia, 34
 Stegomyia, 33, 135, 136, 137, 138, 153, 154, 212, 217, 245, 271, 310, 449, key, 219-220
stegomyia, 151
 stenotrus, 248, 255, 256
 stevensoni, 155, 174.
 stricklandi, 61, 80.
subalbatus, 314
 subalbopictus, 220, 238.
 submediopunctatus, 230, 231.
subniveus, 205
 subnormalis, 72
 subsmilis, 153, 156, 203, 204.
 subulifer, 25
 suffusus, 140, 154, 155, 192, 194, 207
 sugens, 245
suknaensis, 257
 sumatranus, 351
 summorosus, 400
 surellensis, 230, 231.
 sylvestris, 139, 246.
 synthetius, 140, 248, 256.

taeniarostris, 391.
taenata Leic. (Aëdes), 250.
taenata Leic (*Lophoceratomyia*), 366.
taeniorhynchoides, 246, 248, 260.
Taeniorhynchus, 118.
taytayensis, 424
tenax Leic, 394.
tenax de Meij, 42
tenax Theo, 391, 398.
 tenuipalpis, 334, 336, 348, 351, 352, 357, 358, 359
Teromyia, 8
 territans, 347
 testacea, 59, 74, 75
 theileri, 337, 389, 414, 416, 417.
 theobaldi, 313, 319.
 Theobaldia, 4, 31, 33, 38, 98, 135, key, 88
Theobaldomyia, 387.
 thomsoni, 140, 212, 213, 215.
 Tipulidae, 428
tipuliformis Edw., 414
tipuliformis Theo, 416, 417.
 tulliana, 118
 tonga, 240
 Topomyia, 32, 33, 45, 47, 48, 135, 447
Toxorhynchites, 8.
 treubi, 311
Tricholeptomyia, 34
Trichorhynchomyia, 376.
Trichorhynchus, 376
trilineata Leic (*Hulecoetomyia*), 153, 185
trilineata Leic. (*Uranotaenia*), 62
trilineatus Theo. (*Culex*), 263
trimaculatus, 248, 264.
 trinitata, 431, 432, 433, 434.
 Tripteroides, 30, 31, 33, 34, 47, 135, 183, 422, keys, 37-38 Sub-genus, 37
 tripunctatus, 310.
 tritaeniorhynchus, 337, 389, 402, 403, 404, 419, 423, 425.
 tulagensis, 232

 uncus, 291.
 unguiculata, 57, 58, 60, 62, 67.
 unineatus, 143, 155, 170.
uniformis Leic (*Culex*), 424.
uniformis Leic. (*Melanoconion*), 348
uniformis Theo. (Aëdes), 278, 279, 281, 282.
uniformis Theo (*Culex*), 336, 362, 370, 371, 372, 373, 374, 375.
uniformis Theo. (*Mansonia*), 112, 126, 127, 128, 129, 130, 131.
unilineata Leic., 74.
unilineatus Theo., 141, 217, 220, 242.
 ummaculata, 65.
 univittatus, 338, 389, 418, 419, 425.
 Uranotaenia, 3, 32, 33, 52, 56, 111, 116, 135, 142, 178, 447; keys, 59-62.

- vagans*, 338, 389, 416, 422
vallistris, 279, 280, 290.
varrata, 368.
variegatus, 240.
varietas, 277
ventralis, 314
venustipes, 182, 157
Verrallina, 277
versicolor, 151.
vexans, I 139 248, 253, 255, 256,
 257.
vicina, 40
virgatipes, 416
viridis, 301, 306, 307, 308.
viridiventer, 338, 367, 378, 380, 381,
 382, 385
viridis, 291.
vishnu Theo , 6, 336, 384, 389, 398,
 399, 400, 402, 403, 405, 408, 412,
 418
vishnu Theo (♂ only), 404
vittatus, 6, 140, 218, 219, 221, 245,
 443.
vorax, 340, 344, 345, 393

w-albus, 220, 232, 233.
whitel, 337, 389, 402, 405.
whitmorei, 336, 388, 406, 407, 408
Worcesteria, 8
Wyeomyia, 48

yerburyi, 279, 293, 294
yusaf, 279, 282, 443.

zelena, 60, 68, 69, 70
Zengomyia, 447
zeylanica, 431, 434.
zonatipes, 240.

PRINTED BY TAYLOR AND FRANCIS,
RED LION COURT, FLEET STREET, E C 4

PLATE I.

Aedes, subgenus *Finlaya* Head and thorax, showing markings

Fig 1	<i>A. (F.) albolateralis</i>	Theo	♂
2	"	"	♀.
3	"	"	♀ (Variation.)
4	"	<i>deccanus</i> Barr	♀
5	"	<i>gubernatoris</i> Giles	♂
6	"	"	♀
7	"	<i>cacharanus</i> Barr.	♂
8	"	"	♀
9	"	<i>khazani</i> Edw.	♂
10	"	"	♀
11	"	<i>assamensis</i> Theo.	♂.
12	"	"	♀
13	"	<i>unicinctus</i> Edw.	♂
14.	"	"	♀.
15.	"	<i>albotaenatus</i> Leic.	♂
16.	"	"	var. <i>miciranus</i> Edw. ♀.



1



2



3



4



5



6



7



8



9



10




11




12




13



14



15



16

PLATE II.


Aedes, subgenus *Finlaya* Side view, showing scaling of
thorax and abdomen and hind femur

- Fig 1 *A (F) assamensis* Theo ♀
2. „ *khazani* Edw. ♂.
3 „ *cogilli* Edw. ♀.
4 „ *inquinatus* Edw. Type ♂.
5 „ *lophoventralis* Theo. ♀.
6 „ *albotæniatus* Leic. ♂.
7 „ *albolateralis* Theo. ♀.



2

3



4

5



6



7

PLATE III.

Aedes, subgenus *Finlaya*. Head and thorax, showing markings

Fig. 1. *A. (F) elsiea* Barr. ♀.

2. „ *saricola* Edw. ♀.

3. „ *simlensis* Edw. ♀.

4. „ *pulchroventer* Giles. ♀.

5. „ *greeni* (Theo.) var. *kanaranus* Barr. ♂

6. „ *greeni* Theo. (type-form). ♀.

7. „ *pullirostris* Edw. ♀.

8. „ *chrysolineatus* Theo. ♀.

9. „ *oreophilus* Edw. ♀.

10. „ *pseudoteniatus* Giles. ♀.

11. „ *suffusus* Edw. ♀.

12. „ *christophersi* Edw. ♀.

13. „ *oreophilus* Edw. ♂.

14. „ *macdonaldi* Edw. ♀. (Showing pale underside of proboscis.)

15. „ *stevensoni* Barr. ♀.

16. „ *dissimilis* Lelo. ♀. (Showing markings of mid-femora.)



1



2



3




4



5




6



7



8



9



10



11



12



13



14



15



16



1



2



3



4



5



6



7



8



9



10



11



12



13



14



15



16



1



2



3



4



5



6



7

PLATE IV.

Aedes, subgenus *Finlaya* Side view, showing markings.

Fig 1 *A. (F) stevensoni* Barr ♂.

2. „ *harveyi* Barr ♀

3 „ *chrysolineatus* Theo. ♀

4 „ *unicinctus* Edw. ♀

5. „ *albotæniatus* Leic var *mikiranus* Edw. ♀


6. „ *dissimilis* Leic ♂




1




2




3



4



5



6

PLATE V

Aedes, subgenus *Finlaya* Side view, showing markings.

- Fig. 1 *A (F.) chrysolineatus* Theo. ♂
2 ,, *shortti* Barr ♂
3. ,, *oreophilus* Edw ♂.
4 ,, *pseudotæniatus* Giles. ♂.






PLATE VI.

Aedes, subgenera *Stegomyia* and *Christophersomyia*.

Head and thorax, showing markings.

- Fig 1. *A. (S) ægypti* Linn. (*S. fasciata*) ♀
 2. „ *albopictus* Skuse ♀.
 3. „ *unalineatus* Theo ♀
 4. „ *vittatus* Big ♀
 5. „ *edwardsi* Barr ♂
 6. „ *w-albus* Theo. ♀.
 7. „ „ ♀. (Variation)
 8. „ „ ♂
 9. „ *annandalei* Theo ♀.
 10. „ *mediopunctatus* Theo ♀
 11. *A. (C) annulirostris* Theo ♀.
 12. „ *thomsoni* Theo. ♀
 13. *A. (S) ægypti* Linn (*S. fasciata*). ♂.
 14. „ *albopictus* Skuse. ♂
 15. „ *vittatus* Big ♂
 16. *A. (C.) thomsoni* Theo. ♂.



1



2



3



4



5



6



7



8



9



10



11



12



13



14



15



16

PLATE VII


Aedes, subgenus *Stegomyia* Side view, showing markings

Fig 1. *A. (S) ægypti* Linn (*S. fasciata*). ♀.

2. „ *albopictus* Skuse. ♀.

3 „ *edwardsi* Barr. ♀

4 „ *vittatus* Big ♀.




AÉDES (STEGOMYIA)

PLATE VIII.

Aedes, subgenera *Christophersomyia* and *Stegomyia*
Side view, showing markings.

- Fig 1. *A* (*C*) *annulirostris* Theo ♀
2 „ *thomsoni* Theo ♀
3 *A.* (*S*) *w-albus* Theo ♀.
4 „ *mediopunctatus* Theo ♀
5 „ *annandalei* Theo ♀

2687




3



2



4



5

7

1

2

3

4